

INFOGRAPHIC

**SPIDS – simplified predictive intubation difficulty score** ☆




Vinícius Caldeira Quintão <sup>a,\*</sup>, Cláudia Marquez Simões <sup>b</sup>, Maria José Carvalho Carmona <sup>a</sup>

<sup>a</sup> Universidade de São Paulo, Hospital das Clínicas, Faculdade de Medicina, São Paulo, SP, Brazil

<sup>b</sup> Hospital Sírio-Libanês, São Paulo, SP, Brazil

Available online 4 November 2020

**SPIDS - simplified predictive intubation difficulty score**  
153 PATIENTS WHO UNDERWENT HEAD AND NECK SURGERIES



MMT, Modified Mallampati Test; RHN<sub>max</sub>, Maximum range of head and neck motion; TMD, thyromental distance; RHTMD, Ratio of height thyromental distance; NPV, negative predictive value.

DOI of original article:

<https://doi.org/10.1016/j.bjane.2020.09.007>

☆ Article reference: Selvi O, Kahraman ST, Tulgar S, Senturk O, Serifsoy TE, Thomas D, et al. Effectiveness of simplified predictive intubation difficulty score and thyromental height in head and neck surgeries: an observational study. Rev Bras Anesthesiol. 2020;70:595–604. <https://doi.org/10.1016/j.bjane.2020.09.007>

\* Corresponding author.

E-mail: [vinicius.quintao@hc.fm.usp.br](mailto:vinicius.quintao@hc.fm.usp.br) (V.C. Quintão).

<https://doi.org/10.1016/j.bjane.2020.10.008>

© 2020 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).