

Richness and distribution of aquatic macrophytes in Brazilian semi-arid aquatic ecosystems

Riqueza e distribuição de macrófitas aquáticas em ecossistemas aquáticos do semi-árido brasileiro

Gustavo Gonzaga Henry-Silva, Rodrigo Sávio Teixeira de Moura and Luciana Lúcia de Oliveira Dantas

Laboratório de Limnologia e Qualidade de Água do Semi-Árido – LIMNOAQUA,
Departamento de Ciências Animais, Universidade Federal Rural do Semi-Árido – UFERSA,
CEP 59625-900, Mossoró, RN, Brazil
e-mail: gustavo@ufersa.edu.br, rosatemo@bol.com.br, lucianaoli_dantas@yahoo.com.br

Abstract: Aim: The aim of this study was to evaluate the richness and distribution of the aquatic macrophytes in the basin of the Apodi/Mossoró River, in the semi-arid region (caatinga) of Rio Grande do Norte, Brazil; **Methods:** A survey of the floristic composition of the aquatic macrophytes was made at 20 sampling stations in the basin at four seasons (August/2007, November/2007, February/2008, May/2008). Specimens of each species were collected and deposited in the Dárdano de Andrade Lima Herbarium of the Universidade Federal Rural do Semi-Árido; **Results:** We found 40 species of aquatic macrophytes, in 33 genera and 22 families. The families with the most species were Poaceae and Cyperaceae, and the most species-rich genera were *Cyperus* and *Eleocharis*. The most common plant form was amphibian (42.5%), followed by emergent (27.5%), free-floating (12.5%), rooted-submersed (10.0%), and floating-leaved (7.5%). The lowest richness was observed at the estuarine region (3 species), and the highest richness in the upper basin (17 species). The rooted-submersed *Hydrothrix gardneri* Hooker f. and *Ceratophyllum demersum* L. were observed in great abundance and frequency in the Santa Cruz Reservoir of Apodi, especially in areas close to cage farms of Nile tilapia (*Oreochromis niloticus*) (Linnaeus, 1758). The most common free-floating species were *Eichhornia crassipes* (Mart.) Solms., *Pistia stratiotes* L., and *Salvinia auriculata* Aubl., predominantly in stretches that run through urban centers; **Conclusion:** The species richness of aquatic macrophytes in aquatic environments of the caatinga is similar to that observed in other basins of Brazil. Because of the many dams and reservoirs in the semi-arid Northeast, inventory and monitoring of aquatic macrophytes have become essential, especially in basins that will receive water from the diversion of the São Francisco River.

Keywords: aquatic plants, floristic inventory, basin, water diversion, caatinga.

Resumo: Objetivo: Nós objetivamos avaliar a riqueza e a distribuição das macrófitas aquáticas nos ambientes aquáticos da bacia hidrográfica do Rio Apodi/Mossoró, no semi-árido do Rio Grande do Norte; **Métodos:** A investigação da composição florística das macrófitas aquáticas presentes em 20 estações de amostragem da bacia hidrográfica, em quatro épocas do ano (Agosto/2007, Novembro/2007, Fevereiro/2008, Maio/2008). As espécies foram coletadas e depositadas no herbário Dárdano de Andrade Lima da Universidade Federal Rural do Semi-Árido; **Resultados:** Foram identificadas 40 espécies de macrófitas aquáticas distribuídas em 33 gêneros e 22 famílias. As famílias com o maior número de espécies foram Poaceae e Cyperaceae e os gêneros mais representativos foram *Cyperus* e *Eleocharis*. A forma biológica mais freqüente foi a anfíbia (42,5%), seguida pelas emergentes (27,5%); folhas flutuantes (12,5%); submersas enraizadas (10,0%) e com folhas flutuantes (7,5%). A menor riqueza foi observada na região estuarina (três espécies), e a maior riqueza na parte alta da bacia hidrográfica (dezessete espécies). As submersas enraizadas *Hydrothrix gardneri* Hooker f. e *Ceratophyllum demersum* L. foram observadas em grande abundância e freqüência no reservatório de Santa Cruz do Apodi, especialmente em áreas próximas as atividades de criação em tanques rede de tilápia do Nilo (*Oreochromis niloticus*) (Linnaeus, 1758). As flutuantes livres de maior ocorrência foram *Eichhornia crassipes* (Mart.) Solms., *Pistia stratiotes* L., e *Salvinia auriculata* Aubl., ocorrendo predominantemente em trechos que atravessam centros urbanos; **Conclusões:** Nós concluímos que a riqueza de espécies de macrófitas aquáticas em ambientes aquáticos associados ao bioma caatinga é semelhante à riqueza observada em outras bacias hidrográficas do Brasil. Devido a grande quantidade de açudes e reservatórios existentes no semi-árido nordestino, o levantamento e o monitoramento das macrófitas aquáticas tornam-se essenciais, especialmente em bacias hidrográficas que receberão água da transposição do rio São Francisco.

Palavras-chave: plantas aquáticas, levantamento florístico, bacia hidrográfica, transposição de água, caatinga.

1. Introduction

The principal characteristics of the Brazilian semi-arid region are the low rainfall, generally concentrated in certain periods of the year; and the narrow temperature range (Maltchik and Florín, 2002). Studies on aquatic ecosystems of the Brazilian semi-arid region have evidenced the high biodiversity of these environments, including aquatic macrophytes, and have demonstrated that patterns of diversity are related to the hydrological extremes of flood and drought (Medeiros and Maltchik, 1999; Silva-Filho and Maltchik, 2000; Medeiros and Maltchik, 2001; Pedro et al., 2006).

Aquatic macrophytes play an important role in the structure and function of the aquatic environment (Engelhardt and Ritchie, 2001; Chambers et al., 2008; Makkay et al., 2008). Many studies have demonstrated the influence of these plants on the communities of benthic macro-invertebrates (Van den Berg et al., 1997; Takeda et al., 2003), fishes (Agostinho et al., 2003; Pelicice et al., 2005; Sánchez-Botero et al., 2008), periphyton (Pompêo and Moschini-Carlos, 2003), and zooplankton (Lansac-Toha et al., 2003). Floristic inventories have provided essential information for the conservation of biodiversity (Camargo et al., 2003; França et al., 2003; Matias et al., 2003; Paz and Bove, 2007; Rocha et al., 2007; Mora-Olivo and Villaseñor, 2007; Martins et al., 2008; Pivari et al., 2008a).

In recent years, studies of aquatic macrophyte communities have intensified, not only because of the ecological importance of these plants, but also because of the possibility of using them as biological indicators, as well as the proliferation of some species in basins impacted by organic pollution, reservoir construction, and water diversion (Pieterse and Murphy, 1990; Mackay et al., 2003; Thomaz et al., 2003; Camargo et al., 2003; Martins et al., 2008). The two largest drainage basins of the State of Rio Grande do Norte (Piranhas/Açu and Apodi/Mossoró) will receive water from the São Francisco River through the Eixo Norte (North Axis) waterway (ANA, 2007). These basin-integration projects may lead to physical and chemical changes of the water, and will also mix the aquatic biological communities through the introduction of species from one basin to another.

Considering that floristic inventories are important to provide useful information for actions to preserve biodiversity and in the management of weeds, and that studies on aquatic macrophytes in the semi-arid region of Rio Grande do Norte are few,

the present work aimed to evaluate the richness and distribution of the aquatic macrophytes in the basin of the Apodi/Mossoró River, in order to compare and understand possible changes in the assemblages of aquatic macrophytes after artificial diversion of the Sao Francisco River.

2. Material and Methods

2.1. Study area

The basin of the Apodi/Mossoró River is located in the Northeast Middle-Eastern river basin of the State of Rio Grande do Norte ($6^{\circ} 22' 08''$; $4^{\circ} 57' 15''$ S and $38^{\circ} 27' 22''$; $37^{\circ} 8' 11''$ W). The aquatic ecosystems of this region lie in the semi-arid Caatinga, and the rivers are intermittent except in their lower, tidal reaches, and in stretches where reservoirs have been constructed. In most of the basin, the annual mean rainfall is about 700 mm (SEMARTH, 2009). The basin covers an area of 14,276 km², and is the largest drainage basin in the state of Rio Grande do Norte, comprising 26.8% of its area. The basin contains 618 recorded reservoirs (27.4% of the total in the state), all of which may be susceptible to colonization by exotic species. The headwaters of the Apodi/Mossoró River are in the mountains near the city of Luís Gomes in western Rio Grande do Norte, at 831 m altitude. The basin is 210 km long, and its largest reservoir, Santa Cruz, has an area of 2,187.5 km² and a maximum water storage capacity of 600,000,000 m³. Only downstream from the Santa Cruz Reservoir is the Apodi/Mossoró River entirely perennial (SEMARTH, 2009).

2.2. Sampling

In order to evaluate the richness and distribution of the aquatic macrophytes, samples of plant material were obtained at three-month intervals (August/2007, November/2007, February/2008, May/2008) at 20 sampling stations in the Apodi/Mossoró River basin (Figure 1). In the sampling stations, we made random walks up to 2 m from the shoreline, during 30 minutes; at some stations we used a boat. The specimens collected were photographed and placed in exsiccatae for subsequent cataloging in the Dárdano de Andrade Lima Herbarium of the Universidade Federal Rural do Semi-Árido. Species were identified through morphological comparison and consultation of specialized literature (Hoehne, 1979; Joly, 1987; Cook, 1990; Velásquez, 1994; Irgang and Gastal, 1996; Lorenzi, 2000; Pott, VJ. and Pott, A., 2000)




Figure 1. Location and hydrography of the Apodi/Mossoró river basin, RN, Northeast Brazil, with the 20 sampling station. Geographical coordinates: Station 1: 06° 22' 10" S and 038° 27' 39" W; Station 2: 06° 22' 08" S and 038° 27' 27" W; Station 3: 06° 14' 05" S and 038° 14' 54" W; Station 4: 06° 08' 05" S and 038° 11' 35" W; Station 5: 06° 16' 88" S and 038° 15' 70" W; Station 6: 05° 45' 68" S and 037° 48' 15" W; Station 7: 05° 45' 08" S and 037° 47' 41" W; Station 8: 05° 40' 09" S and 037° 47' 59" W; Station 9: 05° 28' 09" S and 037° 31' 29" W; Station 10: 05° 26' 43" S and 037° 31' 09" W; Station 11: 05° 13' 09" S and 037° 21' 46" W; Station 12: 05° 12' 22" S and 037° 21' 10" W; Station 13: 05° 12' 22" S and 037° 20' 25" W; Station 14: 05° 12' 13" S and 037° 20' 44" W; Station 15: 05° 12' 17" S and 037° 20' 13" W; Station 16: 05° 11' 00" S and 037° 20' 15" W; Station 17: 05° 09' 33" S and 037° 17' 02" W; Station 18: 05° 10' 04" S and 037° 14' 32" W; Station 19: 05° 12' 30" S and 037° 11' 01" W; Station 20: 04° 57' 15" S and 037° 08' 11" W.

and with experts. The scientific names used here follow the APGII classification system (2003), and the spelling of names was confirmed by the Missouri Botanical Garden (2010).

To categorize the aquatic macrophytes, we used the definitions of Cook (1996) and Irgang and Gastal (1996). The species were classified as amphibian (Am), emergent (Em), free-floating (FF), rooted-submersed (RS), floating-leaved (FL), and free-submersed (FS). The frequency of occurrence was calculated from the occurrence of each species at the sampling stations at the four periods of the year. Taxa were classified as: Constant = $F > 50\%$, Common = $10\% < F \leq 50\%$, or Rare = $F \leq 10\%$ (Lobo and Leighton, 1986).

3. Results and Discussion

In the Apodi/Mossoró River basin were identified 40 aquatic macrophyte species, members of 33 genera and 22 families (Table 1). The families with the largest numbers of species were Poaceae and Cyperaceae, comprising 30% of the total (Figure 2). The most species-rich genera were

Cyperus and *Eleocharis*, with four and three species, respectively (Figure 3). Both families contain many taxa of aquatic macrophytes (França et al., 2003; Matias et al., 2003; Rocha et al., 2007; Pivari et al., 2008b). Some species of these families can colonize a wide variety of aquatic environments and inhabit ecotone areas (Cook, 1996; Leite et al., 2009). Currently, it is estimated that there are about 10,035 species of Poaceae and 5,000 species of Cyperaceae (Goetghebeur, 1998; Govaerts et al., 2007). Pott, VJ. and Pott, A. (2000), in a study of the distribution of aquatic macrophytes of the Pantanal, identified a total of 273 species, 26 Poaceae and 22 Cyperaceae, together comprising 17.6% of all species found. The importance of these families in aquatic environments is due to their rhizomes, tubers, and stolons, which facilitate vegetative propagation (Pott et al., 1989; Bove et al., 2003; França et al., 2003; Matias et al., 2003).


The most common biological form of macrophyte was the amphibian (42.5%), with 17 species, followed by the emergent (27.5%), free-floating (12.5%), rooted-submersed (10.0%), and floating-

Table 1. List of occurrences of species in hydrographic basin of the Apodi/Mossoró River, RN, Northeast Brazil. (Biological forms: Am = Amphibian, Em = Emergent, FL = Floating leaved, FF = Free-floating, RS = Rooted submersed).

Family/Specie	Biological form	Sampling stations
Acanthaceae		
<i>Ruellia paniculata</i> L.	Am	14, 15
Aizoaceae		
<i>Sesuvium portulacastrum</i> (L.) L.	Am	17, 18, 19, 20
Alismataceae		
<i>Echinodorus grandiflorus</i> (Cham. & Schldl.) Micheli	Em	1, 2, 3, 4
Amaranthaceae		
<i>Blutaparon portulacoides</i> (A. St.-Hil.) Mears	Am	10, 11, 12, 13, 14, 15, 16, 18, 19, 20
<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	Em	4, 5, 7, 8, 9, 11, 13, 15, 16, 17, 18, 19
Araceae		
<i>Pistia stratiotes</i> L.	FF	7, 8, 11, 12, 13, 14, 15, 16, 17
<i>Wolffia brasiliensis</i> Wedd.	FF	7
Asteraceae		
<i>Eclipta alba</i> (L.) Hassk.	Am	3, 4, 6, 11
Ceratophyllaceae		
<i>Ceratophyllum demersum</i> L.	RS	5, 6
<i>Ceratophyllum submersum</i> L.	RS	5, 6
Chenopodiaceae		
<i>Salicornia gaudichaudiana</i> Moq.	Am	19
Convolvulaceae		
<i>Ipomoea fistulosa</i> Mart. ex. Choisy	Am	3, 4, 7, 6, 7, 8, 9, 10, 11, 12, 14, 16, 17, 18
Cyperaceae		
<i>Cyperus esculentus</i> L.	Em	2, 3, 4, 6, 7, 9, 10, 13, 18
<i>Cyperus gardneri</i> Nees	Em	1, 2, 3, 8
<i>Cyperus ligularis</i> L.	Em	4, 6, 10, 13
<i>Cyperus surinamensis</i> Rottb.	Em	3
<i>Eleocharis acutangula</i> (Roxb.) Schult.	Am	3, 4, 7, 8, 12, 18
<i>Eleocharis geniculata</i> (L.) Roem. & Schult.	Am	1, 4, 6, 9, 10, 14, 17, 18
<i>Eleocharis interstincta</i> (Vahl) Roem. & Schult.	Am	3
Fabaceae		
<i>Neptunia plena</i> (L.) Benth.	Am	7, 10, 17, 18
<i>Stylosanthes guianensis</i> (Aubl.) Sw.	Am	3
Hidrophyllaceae		
<i>Hydrolea spinosa</i> L.	Em	1, 2
Lemnaceae		
<i>Lemna valdiviana</i> Phil.	FF	4, 12, 13, 14
Limnocharitaceae		
<i>Limnocharis flava</i> (L.) Buchenau	Em	4
<i>Hydrocleys parviflora</i> Seub.	FL	3
Nymphaeaceae		
<i>Nymphaea alba</i> L.	FL	3, 4
Onagraceae		
<i>Ludwigia helminthorrhiza</i> (Mart.) H. Hara	FL	4, 5, 6, 7, 9, 15, 16
<i>Ludwigia peploides</i> (Kunth) P.H. Raven	Em	8, 10, 11, 12, 13, 14, 15
Poaceae		
<i>Cenchrus echinatus</i> L.	Am	3
<i>Chloris barbata</i> Sw.	Am	6, 12, 13, 14, 15, 18

Table 1. Continued...

Family/Specie	Biological form	Sampling stations
<i>Echinochloa polystachya</i> (Kunth.) Hitchc.	Em	12, 13, 14
<i>Paspalidium paludivagum</i> (Hitchc.& Chase) Parodi	Em	4, 7, 10, 15, 18
<i>Paspalum vaginatum</i> Sw.	Am	11, 12, 13, 14, 15, 16, 17, 18
Pontederiaceae		
<i>Heteranthera seubertiana</i> Solms	Am	3, 4, 8, 9, 10, 11
<i>Eichhornia crassipes</i> (Mart.) Solms	FF	4, 10, 11, 12, 13, 14, 15, 16
<i>Hydrothrix gardneri</i> Hook. f.	RS	6, 9, 10
Potamogetonaceae		
<i>Ruppia maritima</i> L.	RS	20
Rubiaceae		
<i>Borreria alata</i> (Aubl.) DC.	Am	3
Salviniaceae		
<i>Salvinia auriculata</i> Aubl.	FF	4, 6, 7, 10, 14, 15, 16
Scrophulariaceae		
<i>Stemodia maritima</i> L.	Am	1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18

**Figure 2.** Number of occurrence of species, per family, in 20 sampling stations of the basin of the Apodi/Mossoró River, RN, Northeast Brazil.

leaved (7.5%) (Figure 4). No free-submersed species were found. The Common species represented 55.0% of the total of aquatic macrophytes, and the Rare and Constant species represented 37.5% and 7.5%, respectively. Among the Common species, the amphibian was the most frequent (45.5%), followed by the emergent (27.3%), free-floating (18.2%), rooted-submersed (4.5%), and floating-leaved (4.5%). Among the Rare species, the amphibian was the most common (40.0%), followed by the rooted-submersed and emergent

(both with 20.0%), floating-leaved (13.3%), and free-floating (6.7%). Only three biological forms were identified for the Constant species: amphibian (66.7%), and emergent (33.3%). (Table 2).

The 47.5% of the species identified in the Apodi/Mossoró River basin can be considered opportunistic, i.e., they have greater ability to use light, water, nutrients, and carbon dioxide, high seed production, and mechanisms to promote the spread and longevity of seeds, especially through dormancy (Lorenzi, 2000). According to Lorenzi




Figure 3. Number of occurrence of species, per principal genera, in 20 sampling stations of the basin of the Apodi/Mossoró River, RN, Northeast Brazil.




Figure 4. Ranking of biological forms occurring in the basin of the Apodi/Mossoró River, RN, Northeast Brazil. Am = Amphibian; Em = Emergent; FF = Free-floating; RS = Rooted-submersed; FL = Floating-leaved.

Table 2. Classification according to frequency of occurrence of aquatic macrophytes from hydrographic basin of the Apodi/Mossoró River, RN, Northeast Brazil.

Species	Occurrence number	Classification
<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	12	Constant
<i>Blutaparon portulacoides</i> (A. St.-Hil.) Mears	10	Common
<i>Borreria alata</i> (Aubl.) DC.	1	Rare
<i>Cenchrus echinatus</i> L.	1	Rare
<i>Ceratophyllum submersum</i> L.	2	Rare
<i>Ceratophyllum demersum</i> L.	2	Rare
<i>Chloris barbata</i> Sw.	6	Common
<i>Cyperus esculentus</i> L.	9	Common
<i>Cyperus gardneri</i> Nees	4	Common
<i>Cyperus ligularis</i> L.	4	Common
<i>Cyperus surinamensis</i> Rottb.	1	Rare
<i>Echinochloa polystachya</i> (Kunth.) Hitchc.	3	Common
<i>Echinodorus grandiflorus</i> (Cham. & Schldt.) Micheli	4	Common
<i>Eclipta alba</i> (L.) Hassk.	4	Common
<i>Eichhornia crassipes</i> (Mart.) Solms	8	Common
<i>Eleocharis acutangula</i> (Roxb.) Schult.	6	Common
<i>Eleocharis geniculata</i> (L.) Roem. & Schult.	8	Common
<i>Eleocharis interstincta</i> (Vahl) Roem. & Schult	1	Rare
<i>Heteranthera seubertiana</i> Solms	6	Common
<i>Hydrocleys parviflora</i> Seub.	1	Rare
<i>Hydrolea spinosa</i> L.	2	Rare
<i>Hydrothrix gardneri</i> Hook. f.	3	Common
<i>Ipomoea fistulosa</i> Mart. ex. Choisy	14	Constant
<i>Lemna valdiviana</i> Phil.	4	Common
<i>Limncharis flava</i> (L.) Buchenau	1	Rare
<i>Ludwigia helminthorrhiza</i> (Mart.) H. Hara	7	Common
<i>Ludwigia peploides</i> (Kunth) P.H. Raven	7	Common
<i>Neptunia plena</i> (L.) Benth.	4	Common
<i>Nymphaea alba</i> L.	2	Rare
<i>Paspalidium paludivagum</i> (Hitchc. & Chase) Parodi	5	Common
<i>Paspalum vaginatum</i> Sw.	8	Common
<i>Pistia stratiotes</i> L.	9	Common
<i>Ruellia paniculata</i> L.	2	Rare
<i>Ruppia maritima</i> L.	1	Rare
<i>Salicornia gaudichaudiana</i> Moq.	1	Rare
<i>Salvinia auriculata</i> Aubl.	7	Common
<i>Sesuvium portulacastrum</i> (L.) L.	4	Common
<i>Stemodia maritima</i> L.	17	Constant
<i>Stylosanthes guianensis</i> (Aubl.) Sw.	1	Rare
<i>Wolffia brasiliensis</i> Wedd.	1	Rare

(2000), the following species can be considered opportunistic: *Alternanthera philoxeroides*, *Cenchrus echinatus*, *Ceratophyllum demersum*, *Cyperus esculentus*, *C. surinamensis*, *Echinochloa polystachya*, *Echinodorus grandiflorus*, *Eclipta alba*, *Eichhornia crassipes*, *Eleocharis acutangula*, *Hydrolea spinosa*, *Ipomoea fistulosa*, *Lemna valdiviana*, *Limncharis flava*, *Neptunia plena*, *Pistia stratiotes*, *Salvinia auriculata*, *Stylosanthes guianensis*, and *Wolffia brasiliensis*. Bove et al. (2003), in a study of temporarily flooded environments of the coastal plain of northern Rio de Janeiro, observed that opportunist species represented 23% of the total taxa, i.e., about half of the proportion found in the present study. This may indicate that the Apodi/Mossoró River is less preserved, providing favorable conditions for the colonization of potential weed species.

The average richness by sampling station was 9 species, ranging from 3 to 17 species (Figure 5). The stations with the highest richness were 4 and 3, with 17 and 15 species respectively; amphibians (41.7%) and emergents (33.3%) predominated at both stations. Station 20 presented the lowest species richness (*Ruppia maritima*, *Salicornia gaudiachaudiana*, and *Sesuvium portulacastrum*), probably because it is close to the estuary, where high salinities provide unfavorable habitat for most aquatic macrophytes. *Stemodia maritima* was the most common species along the river basin, while the emergent *Ludwigia helminthorrhiza* and *Heteranthera seubertiana* occurred in stretches less impacted by domestic-sewage discharge, as well

as the submersed genus *Ceratophyllum* and the species *Hydrocleys parviflora*. The most common free-floating species were *Eichhornia crassipes*, *Pistia stratiotes*, and *Salvinia auriculata* Aubl, which occurred predominantly in stretches that cross the urban center of the basin.

Most aquatic macrophyte species found in the present survey have wide geographical distributions, and many of them are native to tropical America (Lorenzi, 2000). The rooted submersed *Hydrothrix gardneri* and *Ceratophyllum demersum* were observed in great abundance and frequency in the Santa Cruz Reservoir, especially near fish farms where Nile tilapia (*Oreochromis niloticus*) (Linnaeus, 1758) are raised in cages. The floating-leaved type was less common overall; the species *Ludwigia helminthorrhiza*, found at eight stations, was most prominent. *Nymphaea alba* and *Hydrocleys parviflora* were noted at only one station, just below the Pau dos Ferros Reservoir, in the highest part of the basin.

Compared to other basin-wide inventories of aquatic macrophytes, the present study found species richness similar to that in the Itanhaém River basin (Pereira, 2002) and the Monjolinho River basin (Viana, 2005). However, species richness was relatively low in comparison to other Brazilian wetlands, such as the Banhado do Taim, State of Rio Grande do Sul (Irgang et al., 1984) and the Pantanal (Pott, VJ. and Pott, A., 2000) (Table 3). We found no studies focusing exclusively on hydrophyte communities of the State of Rio Grande do Norte, and in the present study, many temporary




Figure 5. Richness of the species recorded in 20 sampling stations of the basin of the Apodi/Mossoró River, RN, Northeast Brazil.

Table 3. Comparison of the aquatic macrophytes registered in hydrographic basin of the Apodi/Mossoró River, RN, Northeast Brazil, and in others Brazilian aquatic environmental.

Specie	Genus	Family	Sites	Author
242	106	54	Pantanal Mato-grossense - MT	Pott, VJ. and Pott, A. (2000, 2003)
189	104	54	Pernambuco State	Sobral-Leite et al. (2010)
126	95	53	Banhado do Taim - RS	Irgang et al. 1984
121	92	46	Semi-arid Ponds - BA	França et al. (2003)
113	69	40	Temporary swampy environments - RJ	Bove et al. (2003)
89	61	35	Pantanal Lakes - MS	Rocha et al. (2007)
82	62	34	Porto Limão Cárceres Lakes - MT	Silva and Carniello (2007)
66	48	27	Pantanal Floating Mats - MS	Pivari et al.(2008a)
64	42	26	Lobo (Broa) Reservoir - SP	Delello (2008)
56	46	35	Silvana Lake - MG	Pivari et al. (2008b)
53	25	23	Lakes of the Rio Doce Valley - MG	Tavares (2003)
50	29	18	Luís Eduardo Magalhães Reservoir - TO	Lolis (2008)
45	33	18	Jijoca Lake - CE	Matias et al. (2003)
42	38	27	Carapebus Lake - RJ	Paz and Bove (2007)
40	33	22	Apodi/Mossoró Hydrographic Basin - RN	Present study
39	31	21	18 Reservoirs of the São Paulo State	Martins et al. (2008)
35	27	22	Itanhaém Hydrographic Basin - SP	Pereira (2002)
35	28	20	Rosana Reservoir (SP/PR)	Boschilia (2007)
31	24	15	Monjolinho Hydrographic Basin - SP	Viana (2005)
29	24	19	Lagoons in Paraná River floodplain - PR/MS	Santos and Thomaz (2007)
21	33	41	Santana Reservoir - RJ	Pitelli et al.(2008)
15	13	11	Salto Grande Reservoir - SP	Tavares et al. (2004)

environments and reservoirs of the Apodi/Mossoró River basin were not sampled. Thus, the real number of aquatic macrophyte species of this semi-arid basin may be underestimated.

This information indicates that the richness of aquatic macrophytes in the caatinga is similar to that observed in other Brazilian basins. Because of the many dams and reservoirs in the semi-arid Northeast region of Brazil, the inventory and monitoring of aquatic macrophytes supports in evaluating the evolution of these plants, and allows to determine whether some species may become weeds, causing problems for the various uses of these environments. The diversion of water from the São Francisco River to the Apodi/Mossoró River basin may also affect the richness and distribution of the aquatic macrophyte community. Thus, it is essential to monitor the aquatic communities in the lentic and lotic environments of this basin.

Acknowledgements

We are grateful to professors José Iranildo Miranda de Melo, Regina Célia de Oliveira, and Vali Pott for their help in identifying the aquatic macrophytes species and Anibal de Sousa Mascarenhas Filho and Janet W. Reid of the Biological Consulting and Editing Services –USA by revision manuscript into English. We also thank CNPq and Petrobras for financial support.

References

- Agência Nacional de Águas. 2007. *Boletim de Monitoramento dos Reservatórios do Nordeste do Brasil*. Agência Nacional de Águas, Superintendência de Usos Múltiplos. Brasília: ANA.
- AGOSTINHO, AA., GOMES, LC. and JULIO Jr., HF. 2003. Relações entre macrófitas aquáticas e fauna de peixes. In THOMAZ, S. and BINI, LM. (Eds.). *Ecologia e Manejo de Macrófitas Aquáticas*. Maringá: Eduem. p. 261-280.
- Angiosperm Phylogeny Group - APG. 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society*, vol. 141, p. 399-436.
- BOSCHILIA, SM. 2007. *Riqueza, composição e padrões de co-ocorrência da assembléia de macrófitas aquáticas no reservatório de Rosana (SP/PR)*. Maringá: Universidade Estadual de Maringá – UEM. [Master thesis].
- BOVE, CP., GIL, ASB., MOREIRA, CB. and ANJOS, RFB. 2003. Hidrófitas fanerogâmicas de ecossistemas aquáticos temporários da planície costeira do estado do Rio de Janeiro, Brasil. *Acta Botanica Brasílica*, vol. 17, no. 1, p. 119-135.
- CAMARGO, AFM., PEZZATO, MM. and HENRY-SILVA, GG. 2003. Fatores limitantes a produção primária de macrófitas aquáticas. In THOMAZ, SM. and BINI, LM. (Eds.). *Ecologia e Manejo de Macrófitas Aquáticas*. Maringá: EDUEM, 2003. p. 59-84

- CHAMBERS, PA., LACOUL, P., MURPHY, KJ. and THOMAZ, SM. 2008. Global diversity of aquatic macrophytes in freshwater. *Hydrobiologia*, vol. 595, no. 1, p. 9-26.
- COOK, CDK. 1990. *Aquatic Plant Book*. The Hague, The Netherlands: SPB Academic Publishing. 228 p.
- COOK, CDK. 1996. *Water plants of the world*. Amsterdam, The Netherlands: SPB Academic Publishing. 557 p.
- DELELLO, D. 2008. *Composição e distribuição (especial e temporal) de macrófitas aquáticas no reservatório do lobo (Broa), Itrapina/Brotas, SP*. Escola de Engenharia de São Carlos, São Carlos. [Master thesis].
- ENGELHARDT, KAM. and RITCHIE, ME. 2001. Effects of macrophyte species richness on wetland ecosystem functioning and services. *Nature*, vol. 411, p. 687-689.
- FRANÇA, F., MELO, E., GÓES NETO, A., ARAÚJO, D., BEZERRA, MG., RAMOS, HM., CASTRO, I. and GOMES, D. 2003. Flora vascular de açudes de uma região do semi-árido da Bahia, Brasil. *Acta Botanica Brasilica*, vol. 17, p. 549-559.
- GOETGHEBEUR, P. 1998. Cyperaceae. In KUBITZKI, K., HUBER, H., RUDALL, P., STEVENS, P. and STÜTZEL, T. (Eds.). *The families and genera of vascular plants*. Berlin: Spreng-Verlag. p. 141-190.
- GOVAERTS, R., SIMPSON, D., BRUHL, J., EGOROVA, T., GOETGHEBEUR, P. and WILSON, K. 2007. *World checklist of Cyperaceae – Sedges*. Surrey: Kew Gardens. 765 p.
- HOEHNE, FC. 1979. *Plantas aquáticas*. São Paulo: Instituto de Botânica. 168 p.
- IRGANG, BE. and GASTAL Jr., CVS. 1996. *Macrófitas aquáticas da planície costeira do RS*. Porto Alegre: UFRGS. 290 p.
- IRGANG, BE., PEDRALI, G. and WAECHTER, JL. 1984. Macrófitas aquáticas da estação ecológica do Taim, Rio Grande do Sul, Brasil. *Roessleria*, vol. 6, no. 1, p. 395-405.
- JOLY, AB. 1987. *Botânica: introdução à taxonomia vegetal*. São Paulo: Nacional. 777 p.
- LANSAC-TÔHA, FM., VELHO, LFM. and BONECKER, CC. 2003. Influência de macrófitas aquáticas sobre a estrutura da comunidade zooplânctônica. In THOMAZ, SM. and BINI, LM. (Eds.). *Ecologia e Manejo de Macrófitas Aquáticas*. Maringá: Eduem. p. 231-242.
- LEITE, KRB., FRANÇA, F. and SCATENA, VL. 2009. Anatomia de espécies anfíbias de Cyperaceae de lagoas do semi-árido, BA, Brasil. *Acta Botanica Brasilica*, vol. 23, no. 3, p. 786-796.
- LOBO, E. and LEIGHTON, G. 1986. Estructuras de las fitocenosis planctónicas de los sistemas de desembocaduras de río y esteros de la zona central de Chile. *Revista de Biología Marina*, vol. 22, no. 1, p. 143-170.
- LOLIS, SF. 2008. *Macrófitas aquáticas do reservatório Luís Eduardo Magalhães – Lajeado – Tocantins: biomassa, composição da comunidade e riqueza de espécies*. Maringá: Universidade Estadual de Maringá – UEM. [Doctoral thesis].
- LORENZI, H. 2000. *Plantas daninhas do Brasil: aquáticas, parasitas e tóxicas*. Nova Odessa: Instituto Plantarum.
- MACKAY, SJ., ARTHINGTON, AH., KENNARD, MJ. and PUSED, BJ. 2003. Spatial variation in the distribution and abundance of submersed macrophytes in an Australian subtropical river. *Aquatic Botany*, vol. 77, p. 169-186.
- MAKKAY, K., PICK, FR. and GILLESPIE, L. 2008. Predicting diversity versus community composition of aquatic plants at the river scale. *Aquatic Botany*, vol. 88, p. 338-346.
- MALTCHIK, L. and FLORIN, M. 2002. Perspectives of hydrological disturbance as the driving force of Brazilian semiarid stream ecosystems. *Acta Limnologica Brasiliensia*, vol. 14, no. 3, p. 35-41.
- MARTINS, D., COSTA, NV., TERRA, MA. and MARCHI, SR. 2008. Caracterização da comunidade de plantas aquáticas de dezoito reservatórios pertencentes a cinco bacias hidrográficas do estado de São Paulo. *Planta Daninha*, vol. 26, no. 1, p. 17-32.
- MATIAS, LQ., AMADO, ER. and NUNES, EP. 2003. Macrófitas aquáticas da lagoa de Jijoca de Jericoacoara, Ceará, Brasil. *Acta Botanica Brasilica*, vol. 17, no. 4, p. 623-631.
- MEDEIROS, ESF. and MALTCHIK, L. 1999. The effects of hydrological disturbance on the intensity of infestation of *Lernaea cyprinacea* in an stream fish community. *Journal Arid of Environments*, vol. 43, p. 351-356.
- MEDEIROS, ESF. and MALTCHIK, L. 2001. Fish stability and diversity in an intermittent stream from the Brazilian semiarid region. *Austral Ecology*, vol. 26, p. 156-164.
- Missouri Botanical Garden. *Tropicos.org. Missouri Botanical Garden*. Available from: www.tropicos.org. Access in: 12/01/2010.
- MORA-OLIVO, A. and VILLASEÑOR, JL. 2007. Diversidad y distribución de la flora vascular acuática de TAMAULIPAS, México. *Journal of the Botanical Research Institute of Texas*, vol. 1, no. 1, p. 511-527.
- PAZ, J. and BOVE, CP. 2007. Hidrófitas Vasculares da Lagoa de Carapebus, Parque Nacional da Restinga de Jurubatiba, Rio de Janeiro, Brasil. *Revista Brasileira de Biociências*, vol. 5, supl. 2, p. 495-497.
- PEDRO, F., MALTCHIK, L. and BIANCHINI Jr., I. 2006. Hydrologic cycle and dynamics of aquatic macrophytes in two intermittent Rivers of the semi-arid region of Brazil. *Brazilian Journal of Biology*, vol. 66, no. 2b, p. 575-585.
- PELICICE, FM., AGOSTINHO AA. and THOMAZ SM. 2005. Fish assemblages associated with *Egeria* in a tropical reservoir: Investigating the effects of plant

- biomass and diel period. *Acta Oecologica*, vol. 27, p. 9-16.
- PEREIRA, MCT. 2002. *Distribuição e abundância de macrófitas aquáticas em relação às características limnológicas em cinco áreas da bacia do rio Itanhaem, litoral sul do Estado de São Paulo*. Rio Claro: Universidade Estadual Paulista "Júlio de Mesquita Filho" – UNESP. 79 p. [Master thesis].
- PIETERSE, AH. and MURPHY, JK. 1990. *Aquatic weeds*. Oxford: Oxford University Press. 593p.
- PITELLI, RLCM., TOFFANELI, CM., VIEIRA, EA., PITELLI, RA. and VELINI, ED. 2008. Dinâmica da comunidade de macrófitas aquáticas no reservatório de Santana, RJ. *Planta Daninha*, vol. 26, no. 3, p. 473-480.
- PIVARI, MO., POTT, VJ. and POTT, A. 2008a.) Macrófitas aquáticas de ilhas flutuantes (baceiros) nas sub-regiões do Abobral e Miranda, Pantanal, MS, Brasil. *Acta Botanica Brasílica*, vol. 22, no. 2, p. 563-571.
- PIVARI, MOD., SALIMENA, FRG., POTT, VJ. and POTT, A. 2008b. Macrófitas aquáticas da lagoa de Silvana, vale do rio doce, Minas Gerais, Brasil. *Acta Botanica Brasílica*, vol. 22, no. 2, p. 563-571.
- POMPÊO, MLM. and MOSCHINI-CARLOS, V. 2003. *Macrófitas aquáticas e perifiton*. Aspectos ecológicos e metodológicos. São Carlos: Rima. 134 p.
- POTT, VJ. and POTT, A. 2000. *Plantas aquáticas do Pantanal*. Brasília: Embrapa. 404 p.
- POTT, VJ. and POTT, A. 2003. Dinâmica da vegetação aquática do Pantanal. In: THOMAZ, SM. and BINI, LM. (Eds.). *Ecologia e manejo de macrófitas aquáticas*. Maringá: Editora da Universidade Estadual de Maringá. p. 145-162.
- POTT, VJ., BUENO, NC., PEREIRA, RAC., SALES, SM. and VIEIRA, NL. 1989. Distribuição de macrófitas aquáticas numa lagoa na fazenda Nhumirim, Nhecolândia, Pantanal, MS. *Acta Botanica Brasílica*, supl. 3, p. 135-168.
- ROCHA, CG., RESENDE, UM. and SOUZA LUGNANI, J. 2007. Diversidade de macrófitas em Ambientes aquáticos do IPPAN na Fazenda Santa Emília, Aquidauana, MS. *Revista Brasileira de Biociências*, vol. 5, supl. 2, p. 456-458.
- SÁNCHEZ-BOTERO, JL., ARAUJO-LIMA, CARM. and GARCEZ, DS. 2008. Efeitos dos tipos de estandes de macrófitas aquáticas e variações do oxigênio dissolvido e da temperatura na distribuição de peixes em lagos da várzea amazônica. *Acta Limnológica Brasileira*, vol. 20, no. 1, p. 45-54.
- SANTOS, AM. and THOMAZ, SM. 2007. Aquatic macrophytes diversity in lagoons of a tropical floodplain: The role of connectivity and water level. *Austral Ecology*, vol. 32, p. 177-190.
- Secretaria de Estado do Meio Ambiente e dos Recursos Hídricos - SEMARH. *Bacia do rio Apodiú/Mossoró*. 2007. Available from: <http://www.semarh.rn.gov.br>. Access in: 10/2009.
- SILVA, RMM. and CARNIELLO, MA. 2007. Ocorrência de macrófitas em lagoas intermitentes e permanentes em Porto Limão Cáceres-MT. *Revista Brasileira de Biociências*, vol. 5, no. 2, p. 519-521.
- SILVA-FILHO, MI. and MALTCHIK, L. 2000. Stability of macroinvertebrates to hydrological disturbance by flood and drought in a Brazilian semiarid river. *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie*, vol. 27, p. 2661-2466.
- SOBRAL-LEITE, M., CAMPELO, MJA., SIQUEIRA-FILHO, JA. and SILVA, SI. 2010. Checklist das macrófitas vasculares de Pernambuco: riqueza de espécies, formas biológicas e considerações sobre distribuição. In ALBUQUERQUE, UP., MOURA, AN. and ARAÚJO, EL. (Eds.). *Biodiversidade, potencial econômico e processos ecofisiológicos em ecossistemas nordestinos*. vol. II. p. 253-280.
- TAKEDA, AM., SOUZA-FRANCO, GM., MELO, SM. and MONKOLSKI, A. 2003. Invertebrados associados às macrófitas aquáticas da planície de inundação do alto rio Paraná (Brasil). In THOMAZ, S. and BINI, LM. (Eds.). *Ecologia e Manejo de Macrófitas Aquáticas*. Maringá: Eduem. p. 243-260.
- TAVARES, KS. 2003. *A comunidade de macrófitas aquáticas em reservatórios do Médio e Baixo Tietê (SP) e em lagoas da Bacia do Médio Rio Doce (MG)*. São Carlos: Universidade Federal de São Carlos. [Master thesis].
- TAVARES, KS., ROCHA, O., ESPÍNDOLA, ELG. and DORNFELD, CB. 2004. Composição taxonômica de macrófitas aquáticas do reservatório de Salto Grande. (Americana, SP). In ESPÍNDOLA, ELG., LEITE, MA. and DORNFELD, CB. *Reservatório de Salto Grande (Americana, SP): caracterização, Impactos e Propostas de Manejo*. São Carlos: Rima Editora. p. 239-252.
- THOMAZ, SM., BINI, LM. and PAGIORO, TA. Macrófitas aquáticas em Itaipu: ecologia e perspectivas para o manejo. In THOMAZ, SM. & BINI, LM. (Eds.). *Ecologia e manejo de macrófitas aquáticas*. Maringá: Eduem, 2003. p. 319-341.
- VAN DEN BERG, MS., H. COOPS, R. NOORDHUIS, J. VAN SCHIE and J. SIMONS. 1997. Macroinvertebrate communities in relation to submerged vegetation in two *Chara*-dominated lakes. *Hydrobiologia*, vol. 342/343, p. 143-150.
- VELÁSQUEZ, J. 1994. *Plantas acuáticas vasculares de Venezuela*. Caracas: Universidad Central de Venezuela. Consejo de Desarrollo Científico y Humanístico. 992 p.
- VIANA, SM. 2005. *Riqueza e distribuição de macrófitas aquáticas no rio monjolinho e tributários (São Carlos, SP) e análise de sua relação com as variáveis físicas e químicas*. São Carlos: Escola de Engenharia de São Carlos. [Master thesis].