

Phytochemistry and pharmacological insights into *Kalanchoe pinnata*: A brief review

Maduabuchukwu Innocent Nkollo ^{*1}  , Rosemary Nkollo Ngwuede ²  ,
Israel Ofejiro Efejene ³  , Chinedu Henry Olele ⁴  , Ben Chuks Iwelumo ⁵  ,
Christian Chibuogwu ⁶  , and Eromosele Michael Aisuodionoe ⁷  

¹ Department of Optometry, College of Medical and Health Sciences, Novena University, Ogume, Delta State, Nigeria,

² Department of Agronomy and Ecological Management, Faculty of Agriculture, Enugu State University of Science and Technology, Agbani, Enugu State, Nigeria, ³ Department of Pharmacology, Faculty of Basic Medical Sciences, Delta State University of Science and Technology, ⁴ Faculty of Pharmacy, College of Medical and Health Sciences, Novena University, Ogume, Delta state, Nigeria, ⁵ Department of Physiology, College of Medical and Health Sciences, Novena University, Ogume, Delta state, Nigeria, ⁶ Institute for Drug-Herbal Medicines-Excipients Research and Development (ID-HEM-ERD) & Department of Biochemistry, University of Nigeria, Nsukka, Enugu State, Nigeria, and ⁷ Department of Physiology, Faculty of Basic Medical Sciences, Delta State University of Science and Technology, Ozoro, Delta state, Nigeria

^{*} Author to whom correspondence should be addressed

Received: 16-04-2025, Accepted: 08-05-2025, Published online: 09-05-2025



Copyright[©] 2025. This open-access article is distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Nkollo MI, et al. Phytochemistry and pharmacological insights into *Kalanchoe pinnata*: A brief review. *Mediterr J Med Res.* 2025; 2(2): 26-31. [Article number: 6]. <https://doi.org/10.5281/zenodo.15368110>

Keywords: *Kalanchoe pinnata*, medicinal plants, phytochemicals, pharmacological activity, traditional medicine

Abstract: *Kalanchoe pinnata*, commonly known as the “Wonder Plant,” is a tropical succulent widely recognized for its broad spectrum of pharmacological activities and rich phytochemical profile. This review summarizes the phytochemical constituents and therapeutic potentials of *K. pinnata* as documented in various scientific studies. The plant demonstrates significant antimicrobial, anticancer, anti-inflammatory, antidiabetic, and wound-healing properties, largely attributed to its abundance of flavonoids, phenols, alkaloids, vitamins, and other bioactive compounds. These findings highlight its value in traditional medicine and support continued research into its pharmacological applications.

Introduction

Medicinal plants have gained widespread acceptance due to their significant role in traditional medicine, offering remedies for a variety of health concerns. One such medicinal plant is *Kalanchoe pinnata*. *Kalanchoe pinnata*, commonly known as the Wonder Plant, Resurrection Plant, Air Plant, or Africa Never Die, is a perennial shrub found in parts of Africa, the West Indies, Asia, Australia, Bermuda, the Mascarenes, the Galapagos Islands, New Zealand, Brazil, Suriname, Polynesia, and Hawaii. However, it is most commonly found in Madagascar and other regions of Africa [1]. The plant grows up to about 1 meter in height and has over 450 known species. It features cylindrically shaped leaves on a succulent stem. What makes *K. pinnata* unique is that all parts of the plant are medicinally useful. It is easily propagated through stem or leaf cuttings [2]. Originally introduced as an ornamental plant, it is now often found growing wild as a weed in plantation areas. In traditional medicine, *Kalanchoe* species have been used to treat infections, rheumatism, and inflammation, and are known for their immunosuppressive properties [1, 2]. In southeastern Nigeria, for instance, *K. pinnata* plays a role in childbirth practices, where it helps facilitate the detachment of a newborn’s placenta. The leaves, when lightly roasted, are used externally to treat fungal skin infections. Leaf infusions are used internally to reduce fever [3]. Additionally, *K. pinnata* is used to manage pneumonia, intestinal

worms, asthma, and other respiratory tract infections, as well as various inflammatory conditions. It is believed to have antitussive, diuretic, and sedative effects. Other reported uses include the treatment of leg edema, stomach ulcers, and kidney [4]. In Ayurvedic medicine, the plant is valued as an astringent, analgesic, and carminative agent. It is also used to relieve nausea, vomiting, otitis (ear infections), convulsions, headaches, irritation, and general weakness. The leaves are particularly recognized in African traditional medicine for their antifungal properties [1, 3].



Figure 1: *Kalanchoe pinnata* plant displaying characteristic fleshy leaves with marginal plantlets [3].

Methods

This review was conducted using secondary data sourced from peer-reviewed articles accessed via databases such as ScienceDirect, ResearchGate, Google Scholar, Hinari, and Cochrane, as well as physical resources available at Novena University Library.

Phytochemical insights

Kalanchoe pinnata has been found to contain a broad range of phytochemicals including lipids, tannins, alkaloids, flavonoids, phenols, glycosides, vitamins, bufadienolides, cardenolides, minerals, and various organic acids [2]. Among the lipids present are triterpenes such as friedelin, α - and β -amyrins, glutinol, 18- α -oleanane, bryophollone, and taraxerol [5]. Sterols like stigmasterol, kaempferol, dimethoxy flavone-7O- β -D-glucopyranoside, and epigallocatechin-3-O-syringate, as well as fatty acids including stearic, palmitic, arachidic, and behenic acids, have also been identified [6]. The tannins and alkaloids include phenanthrenes, alkanes, alkanols, n-triacontane, and hentriacontane. Flavonoids, phenols, and glycosides such as phosphoenolpyruvate, p-coumaric acid, ferulic acid, protocatechuic acid, cinnamic acid, 4-hydroxybenzoic acid, caffeic acid, and syringic acid are also present [2]. Moreover, compounds like astragalin, 3,8-dimethoxy-4,5,7-trihydroxyflavone, luteolin, rutin, and O-diarabinoside have been identified. The cardenolides, including peposterol, campesterol, isofucosterol, β -sitosterol, clionasterol, 22-dihydrobrassicasterol, bryophyllins A and B, and bufadienolide (noted for its insecticidal properties), contribute to the plant's insecticidal effects [7]. Steroidal glycosides found in the plant include 25-methyl-5 α -ergost-24-en-3- β -ol, ergosta-5,24-dien-3- β -ol, 25-methyl-ergosta-5,24-dien-3- β -ol, 5 α -stigmast-24-en-3- β -ol, (24S)-stigmast-25-en-3- β -ol, (24R)-stigmast-25-en-3- β -ol, patuletin, 3-O-(4-O-acetyl- α -L-rhamnopyranosyl)-7O-(2-O-acetyl- α -L-rhamnopyranoside), and quercetin-3-O- α -L-arabinopyranosyl (1 \rightarrow 2)- α -L-rhamnopyranoside-often used as a marker compound for the plant [7]. Beyond fatty acids, other organic acids such as oxalic acid, citric acid, isocitric acid, oxaloacetate, malic acid, and succinic acid have been detected [2]. The plant is also a source of vitamins including riboflavin, thiamine, niacin, pyridoxine, and amino acids such as glycine, cysteine, methionine, tyrosine, phenylalanine, glutamic acid, as well as protein hydrolysates and casein hydrolysates. Minerals identified include sodium, calcium, potassium, phosphorus, magnesium, manganese, iron, copper, and zinc. The abundance of these

bioactive constituents contributes to the plant's chemoprotective, anti-infective, anticancer, and insecticidal properties. The leaves and bark have also been traditionally used to treat diarrhea and nausea [8]. Molecular formula, Structure description, and functions of key phytochemicals present in Kalanchoe (**Tables 1 and 2**).

Table 1: Triterpenoids

Compound	Molecular Formula	Structure Description	Function
Friedelin	C ₃₀ H ₅₀ O	Pentacyclic triterpenoid with a ketone group	Exhibits anti-inflammatory and hepatoprotective properties [1]
α -Amyrin	C ₃₀ H ₅₀ O	Ursane-type pentacyclic triterpenoid with a hydroxyl group at C-3	The precursor to ursolic acid; possesses anti-inflammatory and analgesic effects [3]
β -Amyrin	C ₃₀ H ₅₀ O	Oleanane-type pentacyclic triterpenoid with a hydroxyl group at C-3	The precursor to oleanolic acid [9] known for anti-inflammatory and hepatoprotective activities [2]
Glutinol	C ₃₀ H ₅₀ O	Pentacyclic triterpenoid with a hydroxyl group	Demonstrates anti-inflammatory and antimicrobial properties [1]
18- α -Oleanane	C ₃₀ H ₅₀	Oleanane-type pentacyclic triterpene hydrocarbon	Serves as a structural backbone for various bioactive triterpenoids [6]
Bryophollone	C ₃₀ H ₄₈ O	Triterpenoid ketone	Exhibits cytotoxic activity against certain cancer cell lines [10]
Taraxerol	C ₃₀ H ₅₀ O	Pentacyclic triterpenoid with a hydroxyl group	Known for anti-inflammatory and hepatoprotective effects [11]

Table 2: Sterols and flavonoids

Compound	Molecular Formula	Structure Description	Function
Stigmasterol	C ₂₉ H ₄₈ O	Plant sterol with a double bond in the side chain	Lowers cholesterol levels; anti-inflammatory and antioxidant properties [3]
Kaempferol	C ₁₅ H ₁₀ O ₆	Flavonoid with hydroxyl groups at positions 3, 5, 7, and 4'	Antioxidant, anti-inflammatory, and anticancer activities [3, 5, 6, 9]
Dimethoxy flavone-7O- β -D-glucopyranoside	C ₂₃ H ₂₄ O ₁₀	Flavone derivative with methoxy groups and a glucose moiety at position 7	Potential antioxidant and anti-inflammatory effects [1, 2]
Epigallocatechin-3-O-syringate	C ₂₂ H ₂₄ O ₁₀	Catechin derivative esterified with syringic acid at the 3-O position	Exhibits strong antioxidant activity [9]

Pharmacological insights

Kalanchoe pinnata has demonstrated a broad spectrum of pharmacological activities supported by in vitro, in vivo, and some clinical studies. These findings provide a scientific basis for many of its traditional medicinal uses.

Anti-inflammatory and analgesic activity: The anti-inflammatory effects of *K. pinnata* have been attributed to the presence of flavonoids, triterpenoids, and other phenolic compounds [12-14]. These constituents inhibit key inflammatory mediators such as prostaglandins, leukotrienes, and cytokines. Animal models have shown significant reductions in inflammation and pain, confirming its analgesic potential [11].

Antimicrobial and antiviral activity: Extracts of *K. pinnata* have demonstrated antimicrobial activity against a variety of pathogens including *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Candida albicans* [15]. The antiviral properties have been particularly noted against herpes simplex virus and influenza strains. These effects are primarily attributed to the flavonoid and bufadienolide content, which disrupt microbial cell walls and inhibit viral replication [8].

Anticancer and cytotoxic effects: Several studies have reported that *K. pinnata* exhibits cytotoxic activity against various cancer cell lines such as HepG2 (liver), MCF-7 (breast), and HeLa (cervical) cells. Bufadienolides, a group of cardiac glycosides present in the plant, induce apoptosis through mitochondrial pathways, inhibit cell proliferation, and alter cellular redox status. These findings support its potential as a source of novel chemotherapeutic agents [10].

Antidiabetic and hypoglycemic activity: Administration of *K. pinnata* extracts in diabetic animal models has shown significant reductions in blood glucose levels. This activity is believed to involve the stimulation of insulin secretion and enhancement of glucose uptake by peripheral tissues, possibly due to the presence of flavonoids and saponins [16].

Antiparasitic and antileishmanial potentials: *Leishmania amazonensis* amastigotes are the causative agents of leishmaniasis, a parasitic disease also known as *uta*, *chiclero ulcer*, or *pian bois*. This condition is characterized by ulcerative lesions on exposed areas of the skin [17]. Several flavonoids present in *Kalanchoe pinnata*, including coumarin and quercetin, have been identified as key contributors to its antileishmanial activity. Specifically, the flavonoids quercitrin, quercetin, and afzelin have demonstrated inhibitory effects against *L. amazonensis* amastigotes in murine models [18].

Antioxidant activity: *K. pinnata* is rich in antioxidant compounds such as quercetin, kaempferol, caffeic acid, and ferulic acid [1]. These compounds scavenge reactive oxygen species (ROS), reduce oxidative stress, and protect biomolecules from oxidative damage [19]. This antioxidant potential contributes to many of the plant's therapeutic effects, including neuroprotection, hepatoprotection, and cardio-protection [3].

Proton pump inhibitory potential: Substances that inhibit gastric acid secretion by targeting the H⁺/K⁺-ATPase enzyme system in the stomach lining are known as proton pump inhibitors, and are effective in the management of peptic ulcers. In traditional medicine, *Kalanchoe pinnata* has long been employed for its anti-ulcer properties. Comparative studies using omeprazole a well-established proton pump inhibitor as a reference drug have demonstrated that *K. pinnata* extracts exhibit significant ulcer-suppressive activity. The leaf and root extracts showed strong anti-inflammatory and anti-ulcer effects in rat models, while the stem extract exhibited comparatively weaker anti-inflammatory activity [19]. The anti-ulcer activity of *K. pinnata* is believed to be mediated through multiple biochemical pathways. Its flavonoid content, particularly quercetin and kaempferol, may inhibit gastric proton pumps (H⁺/K⁺-ATPase), reduce oxidative stress, and promote mucosal defense by enhancing prostaglandin synthesis [3]. Additionally, the antioxidant and free radical-scavenging properties of its phytochemicals help stabilize gastric mucosa, reduce lipid peroxidation, and suppress pro-inflammatory cytokines such as TNF- α and IL-1 β , contributing further to its gastroprotective effects [10].

Wound healing and skin regeneration: Topical application of *K. pinnata* extracts has been shown to accelerate wound healing by promoting collagen synthesis, angiogenesis, and epithelialization. These effects are largely attributed to its anti-inflammatory, antimicrobial, and antioxidant properties [20].

Diuretic and antiurolithiatic effects: Experimental studies have demonstrated increased urinary output and a reduction in kidney stone formation following treatment with *K. pinnata* extracts [4, 16]. The mechanism is believed to involve increased renal clearance and reduced crystallization of oxalate and phosphate [8].

Hepatoprotective activity: *K. pinnata* has shown significant hepatoprotective effects in animal models exposed to hepatotoxins such as carbon tetrachloride and paracetamol. The protective action is thought to be due to its antioxidant and membrane-stabilizing properties [10].

Immunomodulatory effects: Polysaccharides and flavonoids present in the plant have been reported to enhance immune responses by stimulating lymphocyte proliferation, increasing cytokine production, and enhancing macrophage activity [21].

Neuropharmacological effects: Preliminary studies suggest anxiolytic, sedative, and anticonvulsant effects, which may be mediated by modulation of the GABAergic system. This indicates potential applications in anxiety, epilepsy, and related neurological disorders [1].

Antifertility and contraceptive activity: Some studies have indicated antifertility effects, especially in male rats, where extracts reduced sperm count and motility. These effects are considered reversible and are thought to be due to steroid compounds interfering with reproductive hormones [3].

Pesticidal potential: *Kalanchoe pinnata* exhibits notable pesticidal activity when applied in agricultural settings, demonstrating significant efficacy in pest elimination. However, studies have shown that the plant is toxic to cattle when ingested [22]. This toxicity is primarily attributed to the presence of bryotoxins and bufadienolides, which are known to induce cardiac glycoside poisoning in animals [23]. This review has synthesized and evaluated the phytochemical and pharmacological properties of *K. pinnata* as reported in

recent and historical scientific literature. Nonetheless, the current body of knowledge should serve as a foundation rather than a culmination. Continued and rigorous scientific investigation into *K. pinnata*, as well as other medicinal plants employed in traditional medicine, is essential. Moreover, research rooted in traditional practices must be systematically documented and supported, as such efforts are pivotal to the advancement of novel drug discovery and development [4].

Conclusion: The significance of *Kalanchoe pinnata* in traditional medicine is profound and well-recognized. Its wide-ranging therapeutic applications-including the management of systemic diseases, and dermatological conditions, and its roles as an antioxidant and hematinic agent highlight its distinguished status among medicinal plants globally. it is, therefore, fitting that it is locally referred to as the wonder plant.

References

1. Biswas SK, Chowdhury A, Das J. Literature review on pharmacological potentials of *Kalanchoe pinnata* (Crassulaceae). African Journal Pharmacy and Pharmacology. 2011; 5(10): 1456-1461. doi: 10.5897/AJPP11.273
2. Okwu DE, Nnamdi FU. A novel antimicrobial phenanthrene alkaloid from *Bryophyllum pinnatum*. African Journal Pharmacy and Pharmacology. 2011; 5(10): 1258-1262. doi: 10.5897/AJPP11.273
3. Joseph B, Sridhar S, Sankarganesh M, Edwin BT. Rare medicinal plant *Kalanchoe pinnata*. Research Journal Microbiology. 2011; 6 (4): 322-327. doi: 10.3923/jm.2011.322.327
4. Dilip SB, Ramkrushna JA. Research and development of therapeutic herbal tablets from *Kalanchoe pinnata* (Oken) extract. Formulation, Optimization and Evaluation. International Journal of Research and Analytical Reviews. 2024; 11(2): 286-342. doi: Nil.
5. Kamboj A, Saluja A. Microscopical and preliminary phytochemical studies on aerial parts (leaves and stem) of *Bryophyllum pinnatum* Kurz. Pharmacognosy Journal. 2010; 2(9): 254-259. doi: 10.1016/S0975-3575(10)80113-0
6. Toshihiro K, Wemc T, Toshitake C, Taro M. Sterols of *Kalanchoe pinnata*: First report of the isolation of both C-24 epimers of 24-alkyl-A-25-sterol from higher plants. Lipids. 1991; 26(8): 660-665. doi: 10.1007/BF02536432
7. Gaind K, Gupta R. Identification of waxes from leaves of *Kalanchoe pinnata*. Planta Medica. 1974; 52(5): 1071-1079. doi: 10.1055/s-0028-1097931
8. Cryer M, Lane K, Greer M, Cates R, Burt S, Andrus M, et al. Isolation and identification of compounds from *Kalanchoe pinnata* having human alpha herpes virus and vaccinia virus antiviral activity. Pharmaceutical Biology. 2017; 55(1): 1586-1591. doi: 10.1080/13880209.2017.1310907
9. Duraipandiyar V, Ayyanar M, Ignacimuthu S. Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. BMC Complement Alternative Medicine. 2006; 6: 35. doi: 10.1186/1472-6882-6-35
10. Supratman U, Fujita T, Akiyama K, Hayashi H, Murakami A, Sakai H, Koshimizu K, Ohigashi H. Anti-tumor promoting activity of bufadienolides from *Kalanchoe pinnata* and *K. diagremontiana* tubiflora. Bioscience, Biotechnology, and Biochemistry. 2001; 65(4): 947-949. doi: 10.1271/bbb.65.947
11. Igwè SA, Akunyili DN. Analgesic effects of aqueous extracts of the leaves of *Bryophyllum pinnatum*. Pharmaceutical Biology. 2005; 43(8): 658-661. doi: 10.1080/13880200500383108
12. Ojewole JA. Antinociceptive, anti-inflammatory and antidiabetic effects of *Bryophyllum pinnatum* (Crassulaceae) leaf aqueous extract. Journal of Ethnopharmacology. 2005; 99(1): 13-19. doi: 10.1016/j.jep.2005.01.025
13. Akhlaq M, Alum MK, Alam MM. Anti-inflammatory potential of medicinal plants. Mediterranean Journal of Pharmacy and Pharmaceutical Sciences. 2022; 2(1): 13-21. doi: 10.5281/zenodo.6399381
14. Bazine HA, Shlaka MA, Mezogi JS, Alghazeer RO, Sherif FM. Phytochemical and pharmacological studies of *Lycium schweinfurthii* methanolic leaves extract (Solanaceae) in mice. Pharmacy and Pharmacology International Journal. 2022; 10(6): 201-206. doi: 10.15406/ppij.2022.10.00386
15. Mummed B, Abraha A, Feyera T, Nigusse A, Assefa S. In vitro antibacterial activity of selected medicinal plants in the traditional treatment of skin and wound infections in Eastern Ethiopia. Biomedical Research International. 2018; 2018: 1862401. doi: 10.1155/2018/1862401
16. George LA, Radha HR, Somashekaraiah BV. Antidiabetic activity of *Kalanchoe pinnata* in alloxan-induced diabetic rats. Asian Journal of Pharmacy Clinical Research. 2019; 12(3): 241-245. doi: 10.22159/ajpcr.2019.v12i3.30160

17. Muzitano MF, Luzineidi WT, Catherine G, Kaiser CR, Ross-Bergmann B, Costa SS. The antileishmanial activity assessment of unused flavonoids from *Kalanchoe pinnata*. *Phytochemistry*. 2006; 67(18): 2071-2077. doi: 10.1016/j.phytochem.2006.06.027
18. Rocha LG, Almeida JR, Macedo RO, Barbosa-Filho JM. A review of natural products with antileishmanial activity. *Phytomedicine*. 2005; 12(6-7): 514-535. doi: 10.1016/j.phymed.2003.10.006
19. Adesanwo JK, Raji Y, Olaleye SB, Obasanjo SA, Fadare OO, Ige OO, Odusanya, OO. Antiulcer activity of methanolic extract of *Bryophyllum pinnatum* in rats. *Journal of Biological Sciences*. 2007; 7(2): 409-412. doi: 10.3923/jbs.2007.409.412
20. Nayak S, Marshall JR, Isitor G. Wound healing potential of ethanolic extract of *Kalanchoe pinnata* Lam. leaf: A preliminary study. *Indian Journal of Experimental Biology*. 2010; 48(6): 572-576. PMID: 20882759.
21. Umbuzeiro-Valent G, Roubicek DA, Haebisch EM. Mutagenic and antimutagenic evaluation of the juice of the leaves of *Bryophyllum calycinum* (*Kalanchoe pinnata*), a plant with antihistamine activity. *Environmental Molecular Mutagen*. 1999; 33(4): 325-327. doi: 10.1002/(SICI)1098-2280(1999)33:4<325::AID-EM10>3.0.CO;2-E
22. McKenzie RA, Franke FP, Dunster PJ. The toxicity to cattle and bufadienolides content of six *Bryophyllum* species. *Australian Veterinary Journal*. 1987; 64(10): 298-301. doi: 10.1111/j.1751-0813.1987.tb07330.x
23. Reppas GP. *Bryophyllum pinnatum* poisoning of cattle. *Australian Veterinary Journal*. 1995; 72(11): 425-427. doi: 10.1111/j.1751-0813.1995.tb06194.x

Acknowledgments: The authors express their sincere gratitude to the management and staff of Novena University, Ogume Campus Library, for their invaluable support and for providing access to essential resources that contributed significantly to the successful completion of this review.

Author contribution: KSSH conceived, and designed the study, and collected data. KSSH & KTW contributed to data analysis. All authors contributed to data analysis and interpretation of data and drafted and reviewed the manuscript for intellectual context. All authors approved the final version of the manuscript and agreed to be accountable for its contents.

Conflict of interest: The authors declare the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical issues: The authors observed ethical issues including plagiarism, informed consent, data fabrication or falsification, and double publication or submission.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

Author declarations: The authors confirm that they have followed all relevant ethical guidelines and obtained any necessary IRB and/or ethics committee approvals.

الكيمياء النباتية والروى الدوائية في نبات الكالانشو بينات: مراجعة موجزة

مادوابوشوكو إينوست نكولو * 1، روزماري نكولو نغوي 2، إسرائيل أوفيجير و إيفجيني 3، تشينيدو هنري أوليلي 4، بن تشوكس إبويلومو 5، كريستيان تشيبوغوا 6، وإيروموسيل مايكل أيسوديونو 7

1 قسم البصريات، كلية العلوم الطبية والصحية، جامعة نوفينا، أوغومي، ولاية دلتا، نيجيريا

2 قسم الزراعة والإدارة البيئية، كلية الزراعة، جامعة إينوجو الحكومية للعلوم والتكنولوجيا، أغباني، ولاية إينوجو، نيجيريا.

3 قسم علم الأدوية، كلية العلوم الطبية الأساسية، جامعة دلتا الحكومية للعلوم والتكنولوجيا.

4 كلية الصيدلة، كلية العلوم الطبية والصحية، جامعة نوفينا، أوغومي، ولاية دلتا، نيجيريا. 5 قسم علم وظائف الأعضاء، كلية العلوم الطبية والصحية، جامعة نوفينا، أوغومي، ولاية دلتا، نيجيريا

1، 6 معهد أبحاث وتطوير الأدوية العشبية والمواد المساعدة (ID-HEM-ERD) وقسم الكيمياء الحيوية، جامعة نيجيريا، نسوكا، ولاية إينوجو، نيجيريا

7 قسم علم وظائف الأعضاء، كلية العلوم الطبية الأساسية، جامعة ولاية دلتا للعلوم والتكنولوجيا، أوزورو، ولاية دلتا، نيجيريا.

* المؤلف الذي توجه إليه المراسلات

ملخص: الكالانشو بينات، المعروف باسم "النبات العجيب"، نبات عصاري استوائي معروف على نطاق واسع بفعاليته الدوائية الواسعة وتركيبته الكيميائية النباتية الغنية. تلخص هذه المراجعة المكونات الكيميائية النباتية وإمكاناته العلاجية كما هو موثق في دراسات علمية مختلفة. يُظهر النبات خصائص مضادة للميكروبات، ومضادة للسرطان، ومضادة للالتهابات، ومضادة لمرض السكري، ومضادة للجروح، ويعزى ذلك بشكل رئيسي إلى وفرة الفلافونويدات، والفينولات، والقلويات، والفيتامينات، وغيرها من المركبات النشطة بيولوجياً. تُبرز هذه النتائج قيمته في الطب التقليدي، وتدعم الأبحاث المستمرة في تطبيقاته الدوائية.