











SHORT COMMUNICATION article

Analysis of hepatoprotective and antioxidant activities, and determination of the phenolic profile of *Moringa oleifera*

Zaib-un-Nisa¹  , Syed Mubashar Sabir¹  , Zulfiqar Ahmad²  
Izhar Ullah³  , and Muhammad Akhlaq^{4*}  

¹ Department of Chemistry, University of Poonch, Rawalakot, AJK, Pakistan

² Faculty of Veterinary and Animal Sciences, University of Poonch, Rawalakot, AJK, Pakistan

³ Department of Pharmacy, University of Poonch, Rawalakot, AJK, Pakistan

⁴ Department of Pharmacy, Hazara University, Manshara, KP, Pakistan

* Author to whom correspondence should be addressed

Received: 21-01-2026, Accepted: 09-03-2026, Published online: 13-03-2026



Copyright© 2026. This open-access article is distributed under the *Creative Commons Attribution License*, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Nisa et al. Analysis of hepatoprotective and antioxidant activities, and determination of the phenolic profile of *Moringa oleifera*. *Mediterr J Med Res*. 2026; 3(1): 86-93. [Article number: 43]. <https://doi.org/10.5281/zenodo.18912004>

Keywords: Antioxidants, HPLC analysis, hepatoprotective activity, *Moringa oleifera*

Abstracts: *Moringa oleifera*, known as the tree of life, is classified as an herbal plant due to its vast medicinal benefits. Usually, the plant is used to cure diseases. This study investigated the antioxidants and hepatoprotective properties of *Moringa oleifera* grown in different regions of Pakistan. Using assays such as DPPH radical scavenging, iron chelation, and phosphomolybdenum reducing activity, the extract revealed strong antioxidant potential and protection against lipid peroxidation in phospholipid homogenates. It also exhibited significant alpha-amylase inhibition and hepatoprotective effect against paracetamol-induced toxicity, confirmed by biochemical and histopathological analysis. Phytochemical profiling of the leaf extract revealed the presence of multiple phenolic and flavonoid compounds, including various syringic acid, 3-caffeoylquinic acid, 5-p-coumaroylquinic acid, 5-caffeoylquinic acid, 4-caffeoylquinic acid, 3,4-dicaffeoylquinic acid, 3-feruloyl-4-caffeoylquinic acid, dihydromyricetin, apigenin-7-glucoside, quercetin-3-galactoside, quercetin-3-glucoside, quercetin-3-rutinoside-7-glucoside, Isorhamnetin-3-hydroxyferuloyl glucoside-7-glucoside, Quercetin 3-(methoxycaffeoyldiglucoside)-7-glucoside, and quercetin-3-(caffeoyl diglucoside)-7-glucoside. Data suggest that *Moringa oleifera* from Pakistan is rich in bioactive compounds contributing to its antioxidant and liver-protective activities.

Introduction

Free radicals are generated in the oxidation process. These radicals have a damaging effect and cause different diseases. The chain of these reactions causes cell spoilage. The antioxidants neutralize these free radicals and protect against oxidative stress [1]. Excessive Reactive oxygen species (ROS) leads to oxidative cells, tissues, and organs damage, which has been observed to warning many biological functions [2]. Antioxidants prevent oxidation by various processes, mostly by scavenging free radicals from the body like superoxide anion, hydrogen peroxide, nitric oxide, and hydroxyl radical [3]. Several plant species have therapeutic effects due to the presence of phenolic compounds [4, 5]. *Moringa oleifera* (*M. oleifera*) is a plant with many reports on its pharmacological and health benefits. *M. oleifera* is found in India but widely distributed in other parts of the world, including Africa, Europe, and Asia. Moringa is known as 'horseradish tree' or 'drumstick tree' [6]. The most abundant compounds found in *M. oleifera* are Quercetin and kaempferol glycosides [7]. The traditional and nontraditional use of Moringa, its pharmacological effects and their phytopharmaceutical

formulations, clinical studies, toxicity profile, and various other uses have recently recognized [8]. The search of literature has shown that there are few studies on the antioxidant and hepatoprotective potential of *M. oleifera* from Pakistan. The overall aim of this study was to determine the *in vitro* antioxidant activities, hepatoprotective activities, and phenolic profile of aqueous extract being used in traditional formulations.

Materials and methods

Collection of plant: The leaves of *M. oleifera* were collected from Sialkot, Pakistan during 2025, and its identification was done in the Department of Botany, University of Poonch, Rawalakot, AJK, Pakistan.

Antioxidant activities: The DPPH radical activity by *M. oleifera* leaves was checked using the method [9]. The ability to chelate iron was studied following the method [10]. The reduction potential of *M. oleifera* leaves was determined following phosphomolybdenum method [11]. The lipid peroxidation was studied in egg yolk phospholipid by the method reported [3].

Alpha amylase inhibitory activity: The starch iodine method was used for antidiabetic assay with slight modifications [12].

In-vivo hepatoprotective activity: Male albino mice (25-35 g) and aged 2.5 months from the breeding center in CEMB University of Panjab were used for *in vivo* studies. The animals were kept in separate cages with access to food and water under controlled conditions. The mice were randomly divided into four groups, each consisting of five animals, and were fasted overnight before the initiation of the experiment. **Group I** (normal control) given normal saline, and **Group II** (hepatotoxic control) was given paracetamol (70.0 mg/kg). **Groups III & IV** (treated groups) were administered oral doses of aqueous extract of *M. oleifera*, 100 mg/kg and 250 mg/kg, respectively. The treatments were continued for seven days, and the animals were fed orally. In the plant group, the extract was fed to mice after three hours of paracetamol dose. On 8th day, all mice were sacrificed and blood was drawn by heart puncture and level of ALT and AST activities were determined by commercial kits. The livers were collected for biochemical and histopathological analysis. Liver was prepared in normal saline (0.9%), and the level of TBARS, NPSH, and Catalase activities was determined. The TBARS was determined by the method of [13]. Catalase was determined by the Aebi method [14]. While non-protein thiol was determined by using the method [15].


HPLC analysis of phenolics and flavonoids: The dried aqueous extract of *M. oleifera* was filtered and subjected to analysis HPLC system by the method [16]. The unknown compounds were identified from the absorption spectra of compounds in the literature [17].

Ethical approval: All animal procedures were in strict accordance with the NIH guide for the Care and Use of Laboratory Animals and were approved by the University of Poonch Rawal Akot Ethical Committee (UPRAEC-2025).

Statistical analysis: The results were expressed as means \pm SD. Different means were compared by LSD test for the significance of the data.

Results and discussion

Radical scavenging activity: The antioxidant potential of aqueous extracts of *M. oleifera* was evaluated using the DPPH free radical scavenging assay at different concentrations. DPPH radical is the most common radical used to measure antioxidant activities [18-20]. DPPH is violet in color and changes its color to yellow when the extract was added. DPPH is reduced by reaction with extract which provides hydrogen atoms. Normally, phenolics and flavonoids have high reducing activities [21-23]. **Figure 1** shows that the scavenging percentage as obtained by the extract was slightly less than that of standard ascorbic acid. These results are in line with previous reports [5, 24].



Figures 1: DPPH radical scavenging activity of *Moringa oleifera*

The values with different superscripts are statistically significant

Iron chelation assay: When Fe^{2+} reacts with 1,10-phenanthroline, a red colored complex is formed. The addition of extract of natural products to this complex fades its color, and it becomes colorless. This shows the chelating capacity of plant extracts [25]. It was hence concluded that by increasing the concentration of the extract, the chelating activity was also increased. In **Figure 2**, At maximum concentration of extract i.e. 300 $\mu\text{g/ml}$, the highest chelating activity was observed by the leaves extract 30.0% and it was comparable with that of ascorbic acid, 78.0%. **Figure 2** shows that there is a high significant difference among different tested concentrations for metal chelation activities ($p < 0.01$).




Figure 2: Metal chelating activity of *Moringa oleifera*

The values with different superscripts are statistically significant

Reducing activity: The reduction potential of plant extracts as evaluated by the phosphomolybdenum assay involves the formation of a green colored complex as a result of a decrease in oxidation state from VI to V [26]. A direct estimate of the reduction potential of *M. oleifera* leaves extract was obtained by measuring absorbance at 695 nm. **Figure 3** shows that there is a high significant difference among different tested concentrations for reducing activities ($p < 0.01$). **Figure 3** shows that maximum reducing ability was shown by *M. oleifera* leaves extract as the absorbance increases with increasing concentration of the extract. Actually, the molybdenum ion is reduced from Mo (V) to Mo (VI), changing the color of the solution from yellowish to green complex.




Figure 3: Reducing activity of *Moringa oleifera*

The values with different superscripts are statistically significant

Lipid peroxidation in egg yolk: Egg yolk phospholipid was induced with lipid peroxidation using iron sulphate, and the anti-lipid peroxidative property of the extract was determined. The extracts reduced the lipid peroxidation in dose dependent manner (**Figure 4**). The 2-thiobarbituric acid reacts with lipid peroxides and generate a pink color chromogen which is estimated at 532 nm.




Figure 4: Percentage inhibition of lipid peroxidation by *c* aqueous extract

Alpha amylase inhibition assay: The present data confirmed a dose-dependent α -amylase inhibitory effect of plant extract, supporting a clear dose-response relationship. *M. oleifera* leaf extract demonstrated inhibitory effects statistically significantly tested concentrations, suggesting limited efficacy in this model (**Figure 5**). Plant-based enzyme inhibitors are increasingly favored for their safety and added antioxidant benefits over synthetic drugs [27]. The findings highlight *M. oleifera* as the most promising natural α -amylase inhibitors, with potential therapeutic or nutraceutical applications in managing T2DM [28]. However, more pharmacological validation of these species is recommended to support their development as plant-based antidiabetic agents. The glucose-lowering effect of the extract may have been produced by the flavonoids, saponins, and tannins [22, 29, 30].




Figure 5: Alpha amylase inhibitory effect of *Moringa oleifera*

The values with different superscripts are statistically different

In-vivo hepatoprotective activity of *M. oleifera*: Paracetamol administration at a dose of 250 mg/kg resulted in liver damage as shown by biochemical parameters and histopathology (**Table 1**, and **Figure 6**). The treatment with aqueous extracts of all plants at dose of 100 and 250 mg/kg after paracetamol treatment reversed these changes (**Table 1**). Paracetamol is a popular analgesic drug, which does not cause toxicity in therapeutic doses but can result in lethal hepatic necrosis in experimental animals and humans and is known hepatotoxic agent. When the liver is damaged, it causes the leakage of cellular enzymes into the plasma, which are markers of hepatotoxicity [31, 32]. These enzymes are measured in serum to know the extent of hepatocellular damage. As the level of these enzymes were increased on paracetamol treatment, it shows possible toxicity (**Table 1**). Administration of *M. oleifera* aqueous extracts (100 mg/kg and 250 mg/kg, p.o.) brought back the altered levels of ALT and AST showing hepatoprotection. The level of TBARS was significantly increased by treatment with paracetamol, which was decreased by plant extracts (**Table 1**). The plant was not only found effective in decreasing TBARS *in vitro* but also *in vivo*. Catalase activity was significantly decreased in the paracetamol treatment group ($p < 0.05$); however, it was restored to control levels with extract treatment. Glutathione release increases from the liver with paracetamol intoxication [33].

However, treatment with extracts of *M. oleifera* restored the levels of NPSH (**Table 1**). The current data are supported by the previous findings [34]. The findings of histopathology revealed that the control group shows hepatic lobules with typical architecture, central veins, intact hepatocyte cords, and minimal cytoplasmic vacuolation. The treated group showed severe foci of hepatocellular necrosis, ballooning degeneration, sinusoidal dilation, and inflammatory cell infiltration- classical signs of acute hepatotoxicity [35]. However, moderately protective sections of *M. oleifera* treated groups exhibited less necrosis, less severe cytoplasmic vacuolation, and partial lobular reconstruction (**Figure 6**).

Table 1: Phenolic profile of *Moringa oleifera* leaves aqueous extract

Rt (min)	Identity	Absorption spectra (nm)	Mean	STD
1.1	Syringic acid	276	69.1	0.34
4.2	3-caffeoylquinic acid	324, 232	285.1	0.69
5.8	5-p-Coumaroylquinic acid	310, 230	119.0	1.0
6.7	3-Caffeoylquinic Acid	326, 298sh	34.0	0.5
7.3	5-Caffeoylquinic acid	326, 298sh	127.5	0.4
8.3	4-Caffeoylquinic acid	326, 298sh	36.8	1.0
9.3	3,4-Di-caffeoylquinic acid	325, 242	61.0	1.3
10.4	3-Feruloyl-4-caffeoylquinic acid	324, 246	35.9	1.2
12.4	Dihydromyricetin	335, 260	9.11	0.1
14.4	Apigenin-7-glucoside	336	14.9	0.3
15.8	Quercetin-3-galactoside	354, 256	62.5	0.4
16.7	Quercetin-3-glucoside	356, 256	59.8	0.3
18.3	Quercetine-3-rutinoside-7-glucoside	342, 254	6.70	0.3
19.7	Isorhamnetin-3-hydroxyferuloylglucoside-7-glucoside	342, 248	7.57	0.1
24.9	Quercetin 3-(methoxycaffeoyldigluside)-7-glucoside	334, 250	1.72	0.02
27.7	Quercetin-3-(caffeoyldigluside)-7-glucoside	335, 253	2.54	0.02




Figure 6: Histopathology showing hepatoprotective action of *Moringa oleifera* aqueous extract

High Performance Liquid Chromatography (HPLC) analysis: Herein, we report the phenolic profile of *M. oleifera* leaves aqueous extract (**Figure 7**). The phenolic profile showed many phenol containing compounds identified by their retention times. By HPLC analysis syringic acid, 3-caffeoylquinic acid, 5-p-coumaroylquinic acid, 5-caffeoylquinic acid, 4-caffeoylquinic acid, 3,4-di-caffeoylquinic acid, 3-feruloyl-4-caffeoylquinic acid, dihydromyricetin, apigenin-7-glucoside, quercetin-3-galactoside, quercetin-3-glucoside, quercetine-3-rutino-side-7-glucoside, isorhamnetin-3-hydroxyferuloylglucoside-7-glucoside, quercetin 3-

(methoxycaffeoyldiglucoside)-7-glucoside and quercetin-3-(caffeoyldiglucoside)-7-glucosidewere identified in the leaves of *M. oleifera*. The retention times and amounts of these compounds are shown in **Table 1**. The earlier studies have reported that leaves contained free phenolics: flavonols (mainly quercetin and Kaempferol glycoside), hydroxycinnamic acid derivatives, and a small amount of lignan (isolariciresinol isomers) [36]. This study reported some new compounds in the extract of *M. oleifera*, highlighting the importance of the study.




Figure 7: HPLC fingerprint of *Moringa oleifera* aqueous extract obtained from leaves showing the presence of different phenolics

Conclusion: *Moringa oleifera* leaves contain important phytochemicals and valuable antioxidant activities. The hepatoprotective activity of *Moringa oleifera* against paracetamol justifies the popular use of the plant against liver disease. This makes the *Moringa oleifera* a suitable candidate for its use in food and pharmaceuticals.

References

1. Pourmorad F, Hosseinimehr SJ, Shahabimajd N. Antioxidant activity of phenol and flavonoid contents of some Iranian medicinal plants. *African Journal of Biotechnology*. 2006; 5(11): 1142-1145. doi: 10.4314/AJB.V5I11.42999
2. Ansari WA, Srivastava K, Nasibullah M, Khan MF. Reactive oxygen species (ROS): Sources, generation, disease pathophysiology, and antioxidants. *Discover Chemistry*. 2025; 2(1): 191. doi: 10.1007/s44371-025-00275-z
3. Sabir SM, Khalid R. HPLC analysis, antioxidant and inhibitory effect of *Ficus carica* fruits and leaves extracts. *Herba Polonica*. 2023; 69(4): 23-32. doi: 10.5604/01.3001.0054.0917
4. Sun W, Shahrajabian MH. Therapeutic potential of phenolic compounds in medicinal plants-natural health products for human health. *Molecules*. 2023; 28(4): 1845. doi: 10.3390/molecules28041845
5. Munayr MS, Alshreef ZAB. Phytochemical characterization and radical-scavenging activity of solvent fractions from corn silk. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(3): 83-93. doi: 10.5281/zenodo.17094971
6. Kou X, B. B, Olayanju JB, Drake JM, Chen N. Nutraceutical or pharmacological potential of *Moringa oleifera* Lam. *Nutrients*. 2018; 10(3): 343. doi: 10.3390/nu10030343
7. Saini RK, Sivanesan I, Keum Y-S. Phytochemicals of *Moringa oleifera*: A review of their nutritional, therapeutic and industrial significance. *Three Biotech*. 2016; 6(2): 203. doi: 10.1007/s13205-016-0526
8. Pareek A, Pant M, Gupta MM, Kashania P, Ratan Y, Jain V, Pareek A, Chaturgoon AA. *Moringa oleifera*: An updated comprehensive review of its pharmacological activities, ethnomedicinal, phytopharma-ceutical formulation, clinical, phytochemical, and toxicological aspects. *International Journal of Molecular Sciences*. 2023; 24(3): 2098. doi: 10.3390/ijms24032098
9. Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *International Journal of Biochemistry and Cellular Biology*. 2007; 39(1): 44-84. doi: 10.1016/j.biocel.2006.07.001
10. Omoba OS, Olagunju AI, Salawu SO, Boligon AA. HPLC-DAD phenolic profiling and *in vitro* antioxidant activities of three prominent nigerian spices. *Preventive Nutrition and Food Science*. 2019; 24(2): 179-186. doi: 10.3746/pnf.2019.24.2.179

11. Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Analytical Biochemistry*. 1999; 269: 337-341. doi: 10.1006/abio.1999.4019
12. Ononamadu C, Ezeigwe O, Tajudeen O, Ihegboro GO, Lawal TA, Salawu K, et al. Starch-iodine assay method underestimates α -amylase inhibitory potential of antioxidative compounds and extracts. *BioTechnologia*. 2020; 101: 45-54. doi: 10.5114/bta.2020.93103
13. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry*. 1979; 95(2): 351-358. doi: 10.1016/0003-2697(79)90738-3
14. Aebi H. Catalase in vitro. *Methods in Enzymology*. 1984; 105: 121-126. doi: 10.1016/s0076-6879(84) 05016-3
15. Jollow DJ, Mitchell JR, Zampaglione N, Gillette JR. Bromobenzene-induced liver necrosis: Protective role of glutathione and evidence for 3,4-bromobenzene oxide as the hepatotoxic metabolite. *Pharmacology*. 1974; 11(3): 151-169. doi: 10.1159/000136485
16. Zeb A. A reversed phase HPLC-DAD method for the determination of phenolic compounds in plant leaves. *Analytical Methods*. 2015; 7(18): 7753-7757. doi: 10.1039/C5AY01402F
17. Taniguchi M, Lindsey JS. Absorption and fluorescence spectra of organic compounds from 40 sources: Archives, repositories, databases, and literature search engines. *Reporters, Markers, Dyes, Nanoparticles, and Molecular Probes for Biomedical Applications XII*, SPIE. 2020; 42-61. doi: 10.1117/12.2542859
18. Baliyan S, Mukherjee R, Priyadarshini A, Vibhuti A, Gupta A, Pandey RP, Chang CM. Determination of antioxidants by DPPH radical scavenging activity and quantitative phytochemical analysis of *Ficus religiosa*. *Molecules*. 2022; 27(4): 1326. doi: 10.3390/molecules27041326
19. Han KSS, Win KT. Chemical compositions, nutritional values, total phenolic content, and antioxidant activity from *Eleocharis dulcis*. *Mediterranean Journal of Medicine and Medical Sciences*. 2026; 2(1): 28-35. doi: 10.5281/zenodo.18636582
20. Ogbeide OK, Omorodion S, Akhidenor FI, Orazulike OJ, Igbinsosa MO, Otoro D, et al. Comparative study on the phytochemical composition, amino acid profile, antioxidant, in vitro anti-inflammatory, and in vitro anti-diabetic activities on the leaf and stem bark of *Acalypha indica*. *Mediterranean Journal of Medical Research*. 2025; 2(2): 55-64. doi: 10.5281/zenodo.15579785
21. Kaurinovic B, Vastag D. Flavonoids and phenolic acids as potential natural antioxidants. *Antioxidants*. 2019; 2(1): 1-14. doi: 10.5772/intechopen.83731
22. Nizamuddin SFS. Polyphenol-rich black chokeberry (*Aronia melanocarpa*) and its therapeutic potential in type 2 diabetes mellitus: A comprehensive review. *Mediterranean Journal of Medicine and Medical Sciences*. 2025; 1(3): 31-42. doi: 10.5281/zenodo.17619107
23. Adeleke PA, Okubena O, Okubena A, Ajayi AM, Okubena O, Jegede FB, et al. Anti-aging potentials of a polyphenol-rich supplement from African Sorghum bicolor leaf sheaths: A narrative review. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(4): 58-77. doi: 10.5281/zenodo.17742769
24. Lu Y, Bao T, Mo J, Ni J, Chen W. Research advances in bioactive components and health benefits of jujube (*Ziziphus jujuba* Mill.) fruit. *Journal of Zhejiang University Sciences*. 2021; 22(6): 431-449. doi: 10.1631/jzus. B2000594
25. Husain FM, Ahmad I, Khan MS, Ahmad E, Tahseen Q, Khan MS, Alshabib NA. Sub-MICs of *Mentha piperita* essential oil and menthol inhibits AHL mediated quorum sensing and biofilm of Gram-negative bacteria. *Frontiers in Microbiology* 2015; 6: 420. doi: 10.3389/fmicb.2015.00420
26. Abu Bakar FI, Abu Bakar MF, Abdullah N, Endrini S, Fatmawati S. Optimization of extraction conditions of phytochemical compounds and anti-gout activity of *Euphorbia hirta* L. (Ara Tanah) using response surface methodology and Liquid Chromatography-Mass Spectrometry (LC-MS) analysis. *Evidence-Based Complementary and Alternative Medicine*. 2020; 2020: 4501261. doi: 10.1155/2020/4501261
27. Bashary R, Vyas M, Nayak SK, Suttee A, Verma S, Narang R, Khatik GL. An insight of alpha-amylase inhibitors as a valuable tool in the management of type diabetes mellitus. *Current Diabetes Reviews*. 2020; 16(2): 117-136. doi: 10.2174/1573399815666190618093315
28. Ahmed R, Khandaker MS. Natural products as of nutraceuticals treatment for neurological disorders: An overview. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(2): 62-69. doi: 10.5281/zenodo.15226021
29. Mthiyane FT, Dlodla PV, Ziqubu K, Mthembu SXH, Muvhulawa N, Hlengwa N, et al. A review on the antidiabetic properties of *Moringa oleifera* extracts: Focusing on oxidative stress and inflammation as main therapeutic targets. *Frontiers in Pharmacology*. 2022; 13: 940572. doi: 10.3389/fphar.2022.940572
30. Le Ahmed R, Uddin MM, Hoque M. Nutraceuticals: Food-based therapeutics and health benefits. *Mediterr J Med Med Sci*. 2025; 1(1): 22-30. doi: 10.5281/zenodo.15771921
31. wis JH, Seeff LB. The Origins of the modern-day study of drug hepatotoxicity: Focus on Hyman J. Zimmerman. *Clinical Liver Diseases*. 2020; 15(Suppl 1). S25-S36. doi: 10.1002/cld.856

32. Moukhtar KM, Douma AD. *Urtica dioica* L. ameliorative efficiency against CCl₄-induced hepatic fibrosis in rats. Mediterranean Journal of Medical Research. 2026; 3(1): 1-13. doi: 10.5281/zenodo.18120112
33. Rotundo L, Pysopoulos N. Liver injury induced by paracetamol and challenges associated with intentional and unintentional use. World Journal of Hepatology. 2020; 12(4): 125-136. doi: 10.4254/wjh.v12.i4.125
34. Roy N, Laskar RA, Sk I, Kumari D, Ghosh T, Begum NA. A detailed study on the antioxidant activity of the stem bark of *Dalbergia sissoo* Roxb., an Indian medicinal plant. Food Chemistry. 2011; 126(3): 1115-1121. doi: 10.1016/j.foodchem.2010.11.143
35. Momo DO, Iyekowa O, Owolabi OJ. Acute and sub-acute toxicity studies of the methanol extract of *Oecophylla longinoda* by oral administration in mice. Mediterranean Journal of Medical Research. 2025; 2(2): 45-54. doi: 10.5281/zenodo.15475213
36. Starzyńska-Janiszewska A, Stodolak B, Fernández-Fernández C, Mickowska B, Verardo V, Gómez-Caravaca AM. Phenolic profile, antioxidant activity and amino acid composition of *Moringa* leaves fermented with edible fungal strains. Foods. 2022; 1(23): 3762. doi: 10.3390/foods11233762

Author's contribution: ZUN, SMS, MA conceived and designed the study. ZUN, SMS, ZA & IU collected data. ZUN, SMS, IH & MA contributed to data analysis. ZUN, SMS & ZA performed data analysis and data interpretation. ZUN, SMS, IZ & MA drafted the manuscript drafting. All authors read and approved the final manuscript.

Conflict of interest: The authors declare the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical issues: The authors completely observed ethical issues, including plagiarism, informed consent, data fabrication or falsification, and double publication or submission.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

Author declarations: The authors confirm that they have followed all relevant ethical guidelines and obtained any necessary IRB and/or ethics committee approvals.

Generative AI disclosure: No Generative AI was used in the preparation of this manuscript.

تحليل الخصائص الوقائية للكبد والمضادة للأكسدة، وتحديد التركيب الفينولي لنبات المورينجا أوليفيرا

زيب النساء¹، سيد مبشر صابر¹، ذو الفقار أحمد²
إظهار الله³، ومحمد أخلاق^{4*}

- 1 قسم الكيمياء، جامعة بونش، روالاكوت، آزاد جامو وكشمير، باكستان
 - 2 كلية العلوم البيطرية والحيوانية، جامعة بونش، روالاكوت، آزاد جامو وكشمير، باكستان
 - 3 قسم الصيدلة، جامعة بونش، روالاكوت، آزاد جامو وكشمير، باكستان
 - 4 قسم الصيدلة، جامعة هزارة، مانسهر، خيبر بختونخوا، باكستان
- * المؤلف المسؤول عن المراسلات

الملخص: تُصنف المورينغا أوليفيرا، المعروفة بشجرة الحياة، ضمن النباتات الطبية نظرًا لفوائدها العلاجية العديدة. وعادةً ما تُستخدم هذه النبتة لعلاج الأمراض. هدفت هذه الدراسة إلى بحث خصائص مضادات الأكسدة وحماية الكبد في المورينغا أوليفيرا المزروعة في مناطق مختلفة من باكستان. وباستخدام اختبارات مثل إزالة جذور DPPH الحرة، واستخلاق الحديد، ونشاط اختزال الفوسفوموليبيديوم، أظهر المستخلص قدرةً قويةً كمضاد للأكسدة وحمايةً من بيروكسدة الدهون في متجانسات الفوسفوليبيد. كما أظهر تثبيطاً ملحوظاً لإنزيم ألفا-أميليز وتأثيراً وقائياً للكبد ضد السمية الناجمة عن الباراسيتامول، وهو ما أكدته التحليلات البيوكيميائية والنسجية المرضية. كشف التحليل الكيميائي النباتي لمستخلص الأوراق عن وجود مركبات فينولية وفلافونويدية متعددة، بما في ذلك حمض السيرينجيك، وحمض 3-كافويل كينيك، وحمض 5-بارا-كومارويل كينيك، وحمض 5-كافويل كينيك، وحمض 4-كافويل كينيك، وحمض 4،3-ثنائي كافويل كينيك، وحمض 3-فيروليل-4-كافويل كينيك، وثنائي هيدروميريسيتين، وأبيجينين-7-جلوكوزيد، وكيرسيتين-3-جالاكتوزيد، وكيرسيتين-3-جلوكوزيد، وكيرسيتين-3-روتينوزيد-7-جلوكوزيد، وإيزورامينيتين-3-هيدروكسي فيروليل جلوكوزيد-7-جلوكوزيد، وكيرسيتين 3-(ميثوكسي كافويل ثنائي جلوكوزيد)-7-جلوكوزيد، وكيرسيتين-3-(كافويل (ديغلوكونيد))-7-جلوكوزيد. تشير البيانات إلى أن نبات المورينجا أوليفيرا من باكستان غني بالمركبات النشطة بيولوجياً التي تساهم في خصائصه المضادة للأكسدة والوقائية للكبد.