

Bioactive, antioxidant, and inhibitory potential against amylase and glucosidase of medicinal plants used in the treatment of infants

Kayode O. Karigidi^{1*}  , Abiola O. Ilori²  , Funmilola M. Ojo³  , and Oluwadamilola T. Aturu³  

¹ Department of Biochemistry, Olusegun Agagu University of Science and Technology, Okitipupa, Ondo State, Nigeria

² Department of Physical Sciences, Olusegun Agagu University of Science and Technology, Okitipupa, Ondo State, Nigeria

³ Department of Biological Sciences, Olusegun Agagu University of Science and Technology, Okitipupa, Ondo State, Nigeria

* Author to whom correspondence should be addressed

Received: 19-01-2026, Accepted: 20-03-2026, Published online: 23-03-2026



Copyright© 2026. This open-access article is distributed under the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Kargidi et al. Bioactive, antioxidant, and inhibitory potential against amylase and glucosidase of medicinal plants used in the treatment of infants. *Mediterr J Med Res.* 2026; 3(1): 127-136. [Article number: 47]. <https://doi.org/10.5281/zenodo.19154488>

Keywords: Antioxidant, bioactive, infant, inhibitory properties, medicinal plants

Abstract: Medicinal plants have been a significant part of health care since time immemorial. They are used in folk medicine to treat and manage infant health. In this study, bioactive, antioxidant, and inhibitory potential against amylase and glucosidase of nine (*Carica papaya L*, *Mangifera indica L*, *Ocimum gratissimum*, *Vernonia amygdalina Schreb*, *Azadirachta indica A.juss*, *Cymbopogon citrate (DC.) Stapf*, *Psidium guajava L*, *Jatropha curcas L*, and *Momordica charantia L*) medicinal plants were sought. The bioactive potential was evaluated using total phenolics and total flavonoid assays. Total antioxidant capacity (TAC), ferric reducing antioxidant potential (FRAP), 1-diphenyl-2-picrylhydrazyl (DPPH) scavenging activity, and 2-azobis-3-ethylbenzothiazoline-6-sulfonate radical scavenging ability (ABTS) were used to determine antioxidant capacity. In contrast, the inhibitory potential was evaluated against amylase (AMY) and glucosidase (GLU) activities. All assays were conducted using standard procedures. The results showed that *Azadirachta indica A.juss* has the highest total phenolic contents (82.82 ± 1.57 mg GAE/100 g) and total flavonoid (71.95 ± 1.19 mg QUE/100 g), while *Vernonia amygdalina Schreb* (54.08 ± 1.69 mg AAE/100 g) and *Mangifera indica L* (68.83 ± 0.09 mg Fe²⁺E/100 g) were highest for TAC and FRAP, respectively. All extracts exhibited concentration-dependent inhibition for DPPH, ABTS, AMY, and GLU. Inhibition coefficient (IC₅₀) was calculated; *Mangifera indica L* exhibited the lowest IC₅₀ for DPPH (28.50 ± 0.57 µg/mL) and ABTS (34.50 ± 0.88 µg/mL), while *Jatropha curcas L* exhibited the lowest IC₅₀ for AMY (140.40 ± 1.08 µg/mL) and GLU (85.50 ± 1.17 µg/mL) inhibitions. The results show that medicinal plants possess bioactive, antioxidant, and inhibitory potentials, justifying their use in infants.

Introduction

Infant is a term used to describe children during the earliest period of their life, before they can walk and talk. Infants are susceptible to infections and deficiency diseases during this period due to the undeveloped immune system [1]. In 2022, about five million infant deaths were recorded globally, while the infant mortality rate is reported as 72 per 1000 live births [2, 3]. In developing countries, infant health is managed by using medicinal plants due to their low cost and availability [4]. Medicinal plants have been the mainstay of health systems since time immemorial due to their ability to modulate physiological functions and inhibit infection. Because of their

phytochemical compounds with diverse biological activities, medicinal plants are used to treat diseases in infants, such as cough, fever, cold, and rashes [5, 6]. Several common medicinal plants are used in Nigeria for infants with various diseases, as described below.

Carica papaya L. (CPL), known as Papaya, is a member of the Caricaceae allied to the Passifloraceae family. The decoction made from fresh Papaya leaves is mixed with tea to treat malaria, high blood glucose, and hypertension in Nigeria [7]. Fresh young leaves are used empirically as food for humans and animals [8, 9].

Mango *Mangifera indica* L. (MIL) is named Mango, belongs to the Anacardiaceae family. Its leaf is used in ethnomedicine to treat diarrhea [10]. Leaves of MIL were used to treat respiratory diseases/infections [11]. It is reported by Ediriweera and others [12] to possess antimicrobial properties over antiplasmodial, antibacterial, antifungal, antiviral, and antimalarial activities.

Ocimum gratissimum L. (OCM), called scent leaf, belongs to the Lamiaceae family and is well-cultivated in Africa, Asia, and South America. It is used to treat diseases such as cough, anemia, diarrhea, pneumonia, pains, and infections [13, 14]. Its leaf has been reported to possess antidiabetic, antimicrobial, and anticancer properties [15]. OCM is considered a prominent source of phytochemicals/secondary metabolites [16]. Over its use in traditional medicine, the leaves are used in various soups [17].

Vernonia amygdalina (VA), referred to as the bitter leaf, is a member of the Compositae family. VA is consumed as a vegetable or used as an herb to treat diabetes and malaria in traditional medicine [18]. In ethnomedicine, the leaves and roots are used for fever, hiccups, kidney problems, and stomach discomfort [19]. The antihelminthic, antimalarial, antitumourigenic, hypoglycemic, and hypolipidaemic properties of VA have been reported [20]. VA is also well used in insect control as an insect repellent against bean weevil [21].

Azadirachta indica A.Juss (AI), referred to as Neem, belongs to the family of Meliaceae. AI is used in malaria, diabetes mellitus, and as an insect repellent [22]. Various parts of Neem have been widely used to treat human illnesses for centuries. Neem twigs are used for cleaning teeth and are regarded as an efficient kind of dental care in traditional medicine [23]. The flower of the Neem is used for the prevention and treatment of bile diseases [19]. The antidiabetic, antimalarial, anticancer, and antimicrobial properties have been reported [24, 25].

Cymbopogon citrate straph (CC), known as lemon grass, is a member of Poaceae family. CC is well used for gastrointestinal problems and antispasmodic effects in folk medicine [26]. The antiviral, antibacterial, antifungal, and anticancer activities of CC have been documented [27]. CC is used as a seasoning or tea substitute for food. Its tea is frequently used in some Saharan countries, such as India and Cuba, to lower blood glucose concentration [28]. In Ayurveda, CC essential oil is used to treat hypertension, fever, stomach disease, and inflammation related to rheumatism, colds, and flu [28]. The CC leaf extract and decoction have various therapeutic effects, including anti-inflammatory, cough-relieving, digestive, anti-influenza, antipyretic, anti-diabetes, and antimalarial [28].

Psidium guajava L. (PG) is referred to as guava. PG belongs to the Myrtaceae family, and the extract from its leaves is used for diarrhea, rheumatism, digestive problems, malaria, and bacterial infections [29]. Guava leaves are used to manage health complications, including blood sugar levels, healthy bowel movements, and constipation [30]. It is documented that PG possesses pharmacological potentials such as anti-inflammatory, analgesic, antibacterial, antioxidant, and antitumor activities [31].

Jatropha curcas (JC), known as the purging nut, belongs to the family of Euphorbiaceae. JC is used in ethnomedicine for diarrhea, wounds, and diabetes mellitus [32, 33]. It is well used in the treatment of malaria infection [34]. Studies of JC roots, stems, bark, leaves, seeds, and fruits have shown numerous biological effects,

such as coagulative, antioxidant, antimalarial, anticancer, wound-healing, antimicrobial, and anti-inflammatory activities [35-37].

Momordica charantia L. (MC) belongs to the family of Cucurbitaceae and is referred to as either bitter melon or bitter gourd. In ethnomedicine, its leaves and fruit are used to treat diseases such as rheumatism, gout, worms, colic, spleen, and liver disorders [38]. MC is documented for the treatment of malaria, microbial and parasitic infections, diabetes, inflammation, hypertension, and strengthening of immunity in the Benin Republic [39, 40]. Numerous studies have stated its ability to lower blood glucose levels in induced diabetic acid [32, 41, 42].

Therefore, despite the use of these medicinal leaves in treating infant diseases. More information is needed on their mechanism of action. Hence, this study was designed to evaluate the polyphenol, antioxidant, and inhibitory potentials against amylase and glucosidase of commonly used medicinal plants in the treatment of infants.

Materials and methods

Sourcing of medicinal plants: The plants were sourced from the Ilaje Local Government Area (LGA), Ondo State, Nigeria, in 2025. The plants were identified and authenticated at the herbarium unit of the Department of Biological Sciences (Table 1). The leaves were dried at room temperature for 42 days and milled into a powder form.

Table 1: Common medicinal plants identifications

| Botanical | Code | Herbarium number |
|---|------|------------------|
| <i>Carica papaya</i> L. | CPL | OAUSTECH/H/00674 |
| <i>Mangifera indica</i> L. | MIL | OAUSTECH/H/00690 |
| <i>Ocimum gratissimum</i> | OCM | OAUSTECH/H/00333 |
| <i>Vernonia amygdalina</i> Schreb | VA | OAUSTECH/H/00680 |
| <i>Azadirachta indica</i> A.juss | AI | OAUSTECH/H/00278 |
| <i>Cymbopogon citratus</i> (DC.) straph | CC | OAUSTECH/H/00691 |
| <i>Psidium guajava</i> L. | PG | OAUSTECH/H/00692 |
| <i>Jatropha curcas</i> L. | JC | OAUSTECH/H/00671 |
| <i>Momordica charantia</i> L. | MC | OAUSTECH/H/00657 |

Preparation of the extracts: The extracts were prepared according to the previous method [43]. 1.0 g of pulverized leaves was solvated in 10 ml of distilled water for 6 hrs. They were centrifuged at 5000 rpm, and the supernatant was used for analysis immediately.

Determination of Total Phenolic Content (TPC): A method of Kim et al. [44], as modified [43], was used to assay TPC. 0.5 ml was added to 0.5 ml of 10.0% of Folin-Ciocalteu phenol reagent. After 5 min, 2.5 ml of 7.0% Na₂CO₃ and 2.5 ml of distilled water were mixed thoroughly. The solution was kept in the dark at ambient temperature for 1 hr. and 30 min and a spectrophotometric reading was taken at 750 nm. TPC was calculated from the Gallic acid standard curve and expressed as Gallic acid equivalent (mg GAE/100 g).

Determination of Total Flavonoid Content (TFC): The method of Park et al. [45] was used to assay TFC. Thus, 0.3 ml of the sample, 3.4 ml of 30.0% of methanol, 0.15 ml of 0.5 M NaNO₂, and 0.15 ml of 0.3 M AlCl₃·6H₂O were added successively. After 5 min, 1.0 ml of 1.0M NaOH was added, and the absorbance was measured at 506 nm. TFC was calculated from the standard curve and presented as quercetin equivalent (mg QUE/100 g).

Total Antioxidant Capacity phosphomolybdate assay (TAC): The method of Prieto et al. [46] was used. 0.2 ml of the sample and 2.0 ml of phosphomolybdate reagent were added and shaken to mix. The solution was capped and put in a boiling water bath at 95 °C for 1 hr. and 30 min. The solution was then cooled, and a spectrophotometric

reading was taken at 695 nm. The total antioxidant capacity was calculated from the ascorbic acid calibration curve and expressed as ascorbic acid equivalent (mg AAE/100 g).

Determination of Ferric-Reducing Antioxidant Potential (FRAP): The method by Benzie and Strain [47] was used. 0.2 ml of the sample and 2.80 ml of the FRAP working reagent were added, shaken, and incubated at ambient temperature for 30 min. Spectrophotometric readings were taken at 593 nm, and the FRAP equivalent was calculated from the ferrous sulphate standard curve as (mg Fe²⁺E/100 g).

Determination of 1, 1-diphenyl-2-picrylhydrazyl scavenging activity(DPPH): The procedure of Gyamfi et al. [48] was used. Appropriate dilutions of 1.0 ml (0.05 - 0.20 mg/ml) were mixed with (40 mg/l) DPPH solution (4.0 ml) solvated in methanol. The solution was mixed, left in the dark for 30 min, and a spectrophotometric measurement was taken at 520 nm.

Determination of 2-azobis-3-ethylbenzothiazoline-6-sulfonate radical scavenging ability (ABTS): The method of Re et al. [49] was used. Appropriate dilutions of 1.0 ml (0.05-0.20 mg/ml) were added to 2.0 ml of ABTS reagent and shaken. Absorbance spectrophotometric measurements were taken at 734 nm, and the percentage inhibition was calculated.

Determination of amylase inhibition assay (AI): The method of Worthington et al. [50] was used. Appropriate dilutions of 1.0 ml (0.05-0.20 mg/ml) and 500 µl (0.02 M) of sodium phosphate buffer (pH 6.9 with 0.006M NaCl) containing 0.5 mg/ml α-amylase solution were added together and permitted to stay at ambient temperature for 10 min. Then, 500 µl (1.0%) of starch solution prepared with 0.02M sodium phosphate buffer (pH 6.9 with 0.006M sodium chloride) was added. The solution was then incubated at 25 °C for 10 min. The reaction was halted with 1.0 ml (96 mM) of dinitrosalicylic acid reagent. The test tubes were then incubated in a boiling water bath for 5 min. The spectrophotometric assessments were taken at 540 nm, and the inhibition percentage was calculated.

Determination of the glucosidase inhibition assay (GI): The method of Apostolidis et al. [51] was used. Appropriate dilutions of 1.0 ml (0.05-0.20 mg/ml) were added to 1000 µl α-glucosidase solution (1.0 U/l) prepared in 0.1M phosphate buffer (pH 6.9) and pre-incubated for 10 min at 25° C. After pre-incubation, 500 µl of 5.0 mM nitrophenyl-glucopyranoside solution prepared in 0.1M phosphate buffer (pH 6.9) was added. The reaction mixture was incubated at 25°C for 5 min. The absorbance of the reaction mixture was measured at 405 nm. The percentage inhibition was calculated.

Statistical analysis: Data were presented as the mean ± S.D. of triplicate readings. Differences among the groups were tested by a one-way analysis of variance (ANOVA). Statistical differences among the means were considered significant at p<0.05.

Results

In this study, the results of the medicinal plants of TPC, TFC, TAC, and FRAP in Nigeria are presented in **Table 2**. Thus, the TPC was statistically significantly higher in AI and MIL than in other medicinal plants (p<0.05). In addition, TFC was significantly higher in AI (p<0.05), while TAC was significantly higher (p<0.05) in VA in the medicinal plants. The FRAP is significantly higher in MIL (p<0.05) than in the other medicinal plants. **Figure 1** shows that the ABTS and DPPH scavenging capacities were concentration-dependent, while **Figure 2** shows the medicinal plants' potential to inhibit AMY and GLU. The IC₅₀ of the medicinal plants for ABTS, DPPH, AMY, and GLU are presented in **Table 3**. The MIL has the lowest IC₅₀ for ABTS (28.50±0.57 µg/ml) and DPPH (34.50±0.88 µg/ml). The lowest IC₅₀ for AMY (140.40±1.08 µg/ml) and GLU (85.50±1.17 µg/ml) was recorded in JC.

Table 2: TPC, TFC, TAC, and FRAP of the medicinal plants

| Medicinal plant | Total Phenolic content (mg GAE/100 g) | Total Flavonoid content (mg QUE/100 g) | Total antioxidant capacity (mg AAE/100 g) | Ferric-reducing antioxidant potential (mg Fe ²⁺ E/100 g) |
|-----------------|---------------------------------------|--|---|---|
| CPL | 75.17±1.58 ^b | 15.92±2.28 ^f | 19.46±1.10 ^d | 67.56±0.24 ^b |
| MIL | 81.22±1.05 ^a | 36.25±1.25 ^c | 22.43±0.76 ^c | 68.83±0.09 ^a |
| OCM | 76.49±1.44 ^b | 58.99±2.34 ^b | 26.12±1.63 ^b | 65.50±0.30 ^c |
| VA | 71.33±0.54 ^c | 11.30±0.91 ^g | 54.08±1.69 ^a | 63.16±1.55 ^d |
| AI | 82.82±1.57 ^a | 71.95±1.19 ^a | 25.86±1.79 ^b | 67.93±0.36 ^b |
| CC | 52.19±0.49 ^f | 12.34±0.48 ^g | 19.59±1.85 ^d | 47.16±1.15 ^f |
| PG | 64.12±0.87 ^e | 17.47±0.09 ^e | 20.39±1.07 ^d | 58.64±0.36 ^e |
| JC | 68.77±0.34 ^d | 24.26±0.77 ^d | 21.10±0.63 ^d | 63.88±0.34 ^d |
| MC | 53.33±0.44 ^f | 11.82±1.33 ^g | 19.14±1.77 ^d | 47.60±0.18 ^f |

Data are expressed as Mean ± SD. Figures with different superscripts are significantly different (p<0.05) *Carica papaya L* (CPL), *Mangifera indica L* (MIL), *Ocimum gratissimum* (OCM), *Vernonia amygdalina Schreb* (VA), *Azadirachta indica A. juss* (AI), *Cymbopogon citratus (DC) straph* (CC), *Psidium guajava L* (PG), *Jatropha curcas L* (JC), and *Momordica charantia L* (MC)




Figure 1: ABTS and DPPH scavenging activities of the medicinal plants

Carica papaya L (CPL), *Mangifera indica L* (MIL), *Ocimum gratissimum* (OCM), *Vernonia amygdalina Schreb* (VA), *Azadirachta indica A. juss* (AI), *Cymbopogon citratus (DC) straph* (CC), *Psidium guajava L* (PG), *Jatropha curcas L* (JC), and *Momordica charantia L* (MC)




Figure 2: Amylase and glucosidase inhibitory activities of the medicinal plants

Carica papaya L (CPL), *Mangifera indica L* (MIL), *Ocimum gratissimum* (OCM), *Vernonia amygdalina Schreb* (VA), *Azadirachta indica A. juss* (AI), *Cymbopogon citratus (DC) straph* (CC), *Psidium guajava L* (PG), *Jatropha curcas L* (JC), and *Momordica charantia L* (MC)

Table 3: IC₅₀ (µg/ml) of the medicinal plants against ABTS, DPPH, AMY, and GLU inhibitions

| Plant | ABTS | DPPH | AMY | GLU |
|-------|--------------------------|--------------------------|--------------------------|--------------------------|
| CPL | 52.25±0.45 ^c | 78.56±0.56 ^b | 357.28±2.11 ^g | 201.96±1.09 ^c |
| MIL | 28.50±0.57 ^a | 34.50±0.88 ^a | 252.50±2.01 ^f | 240.05±2.26 ^e |
| OCM | 84.40±1.02 ^d | 200.05±2.68 ^c | 228.40±1.57 ^e | 200.50±1.08 ^c |
| VA | 215.40±1.21 ^h | 208.57±1.99 ^e | 180.40±2.84 ^b | 199.40±2.10 ^c |
| AI | 34.38±0.24 ^b | 204.55±1.57 ^d | 218.50±1.44 ^d | 200.15±1.89 ^c |
| CC | 169.78±1.08 ^f | 333.50±2.54 ^g | 141.80±1.98 ^a | 206.50±2.08 ^d |
| PG | 197.34±1.11 ^g | 203.28±1.08 ^d | 202.10±2.57 ^e | 181.23±1.68 ^b |
| JC | 124.32±1.28 ^e | 199.04±1.45 ^e | 140.40±1.08 ^a | 85.50±1.17 ^a |
| MC | 188.87±2.45 ^g | 274.20±1.25 ^f | 185.40±1.50 ^b | 212.50±2.88 ^d |

Results are expressed as Mean ± SD. Figures with different superscripts are significantly different (p<0.05) *Carica papaya L* (CPL), *Mangifera indica L* (MIL), *Ocimum gratissimum* (OCM), *Vernonia amygdalina Schreb* (VA), *Azadirachta indica A. juss* (AI), *Cymbopogon citratus (DC) straph* (CC), *Psidium guajava L* (PG), *Jatropha curcas L* (JC), and *Momordica charantia L* (MC)

Discussion

Since ancient times, medicinal plants have been used in infants to treat diseases and improve children's health in traditional medicine [52]. Despite advances in the development of drugs, they are still used to date, especially in developing countries, because of their availability and marginal costs. Thus, this study investigated the bioactive, antioxidant, and inhibitory potentials of nine medicinal plants used in folk medicine to treat children. Medicinal plants are sources of many bioactive compounds, such as phenolics and flavonoids [53-55]. These compounds have been reported to exhibit biological activities: antiviral, antibacterial, anticancer, antidiabetic, and anti-inflammatory [5, 56]. These secondary metabolites are believed to be safer when derived from medicinal plants than when synthesized in the laboratory [57]. This study has confirmed the presence and abundance of phenolics and flavonoids in the medicinal plants studied. These phytochemicals conferred many therapeutic functions on the medicinal plants. The values obtained for TPC and TFC were comparable to those of other medicinal plants [58].

Fetal growth restriction has been reported to affect up to 25.0% of live births in low- and middle-income countries, and this has been linked to oxidative stress, which might lead to neurodegenerative diseases in the future [59]. Antioxidants are chemical entities that halt the activities of free radicals, thereby inhibiting oxidative stress that can lead to chronic and neurodegenerative diseases [60, 61]. The current study has confirmed that these medicinal plants have potent antioxidant power, which can mitigate oxidative stress and prevent the development of chronic and neurodegenerative diseases in infants. This enormous antioxidant potential might serve as the basis for using medicinal plants in infants. The antioxidant value of these medicinal plants ranked well compared with values reported for other medicinal plants by other researchers [62]. Diabetes mellitus is an autoimmune disease that affects children around the globe and has continued to be on the rise [63, 64]. Genetics alone cannot explain this incidence; therefore, the lifestyle and diet of expecting and nursing mothers might also be implicated [65]. One of the treatment mechanisms for this disease is the inhibition of amylase and glucosidase. These break large polysaccharides, such as starch, into smaller monosaccharide units such as glucose. The plants could inhibit these enzymes, thereby conferring certain antidiabetic properties. This could be the reason for its use in children's health. The IC₅₀ values of these medicinal plants ranked well compared with other plants [66].

Conclusion: The nine medicinal plants included in this study used for the treatment of infants possess bioactive, antioxidant, and inhibitory properties against amylase and glucosidase. These properties justify their use in infant health cases.

References

1. Kollmann TR, Kampmann B, Mazmanian SK, Marchant A, Levy O. Protecting the newborn and young infant from infectious diseases: Lessons from immune ontogeny. *Immunity*. 2017; 46(3): 350-363. doi: 10.1016/j.immuni.2017.03.009
2. Qamar K, Essar MY, Siddiqui JA, Salman A, Salman Y, Head MG. Infant and child mortality in Afghanistan: A scoping review. *Health Scientific Reports*. 2024; 7(7): e2224. doi: 10.1002/hsr2.2224
3. Nwanze LD, Siuliman A, Ibrahim N. Factors associated with infant mortality in Nigeria: A scoping review. *PLoS ONE*. 2023; 18(11): e0294434. doi: 10.1371/journal.pone.0294434
4. Elmansuri NO, Mhani LA, Elhaddar SE, Shushani MA. Libyan mothers' awareness of natural products among infants. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2022; 2(2): 38-43. doi: 10.5281/zenodo.6780482
5. Choudhary S, Zehra A, Mukarram M, Wani KI, Naeem M, Hakeem KR, et al. Potential uses of bioactive compounds of medicinal plants and their mode of action in several human diseases. In: Aftab T, Hakeem KR, Eds. *Medicinal and Aromatic Plants*. Springer, Cham. 2021; doi: 10.1007/978-3-030-58975-2_5
6. Rafi IK. Cough care with medicinal herbs: Safe and natural alternatives. *Mediterranean Journal of Medicine and Medical Sciences*. 2025; 1(2): 17-22. doi: 10.5281/zenodo.16763823
7. Abdelrahim LA, Hiblo HF. Effect of *Carica papaya* Linn (Caricaceae) leaf extracts on reactive oxygen species production. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(3): 52-56. doi: 10.5281/zenodo.16739978
8. Sharma A, Sharma R, Sharma M, Kumar M, Barbhai MD, Lorenzo JM, et al. *Carica papaya* L. leaves: Deciphering its antioxidant bioactives, biological activities, innovative products, and safety aspects. *Oxidative Medicine Cellular Longevity*. 2022; 2022: 331-320. doi: 10.1155/2022/2451733
9. Lim X, Chan J, Japri N, Lee J, Tan T. *Carica papaya* L. Leaf: A systematic scoping review on biological safety and herb-drug interactions. *Evidence-Based Complementary and Alternative Medicine*. 2021; 21: 5511221
10. Yakubu MT, Salimon SS. Antidiarrhoeal activity of aqueous extract of *Mangifera indica* L. leaves in female albino rats. *Journal of Ethnopharmacology*. 2015; 163: 135-141. doi: 10.1016/j.jep.2014.12.060
11. Zhang Y, Chen Q, Liu MY, Ruan JY, Yu HY, Li J, Wang T. Effects of benzophenones from mango leaves on lipid metabolism. *Chemical and Pharmaceutical Bulletin*. 2019; 67(7): 634-639. doi: 10.1248/cpb.c18-00905
12. Ediriweera MK, Tennekoon KH, Samarakoon SR. A review on ethnopharmacological applications, pharmacological activities, and bioactive compounds of *Mangifera indica* (Mango). *Evidence-Based Complementary and Alternative Medicine*. 2017; 2017: 6949835. doi: 10.1155/2017/6949835
13. Akara EU, Okezie E, Ude VC. *Ocimum gratissimum* leaf extract ameliorates phenylhydrazine-induced anemia and toxicity in Wistar rats. *Drug Metabolism and Personalized Therapy*. 2021; 36(4): 311-320. doi: 10.1515/dmpt-2020-0185
14. Nwaneri MGU, Chukwunwejim RC, Ezejiegu CK, Nedum CH, Egbuna RN, Ebenebe IN. Evaluation of the antifungal efficacy of *Mitracarpus scaber* extracts and *Ocimum gratissimum* oil against clinical isolates of fungi. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(4): 29-37. doi: 10.5281/zenodo.17455781
15. Eid AM, Jaradat N, Shraim N, Hawash M, Issa L, Shakhsher M, et al. Assessment of anticancer, antimicrobial, antidiabetic, anti-obesity, and antioxidant activity of *Ocimum basilicum* seeds essential oil from Palestine. *BMC Complementary Medicine and Therapy*. 2023; 23(1): 221. doi: 10.1186/s12906-023-04058-w
16. Neeharika B, Vijayalaxmi KG, Suvarna VC. Antioxidant, antidiabetic, and antimicrobial efficacy of germinated *Ocimum gratissimum* and *Ocimum basilicum* seed. *Journal of the Sciences of Food and Agriculture*. 2025; 105(7): 3843-3856. doi: 10.1002/jsfa.14141
17. Ugbogu OC, Emmanuel O, Agi GO, Ibe C, Ekweogu CN, Ude VC, et al. A review on the traditional uses, phytochemistry, and pharmacological activities of clove basil (*Ocimum gratissimum* L.). *Heliyon*. 2021; 7(11): e08404. doi: 10.1016/j.heliyon.2021.e08404
18. Yeap SK, Ho WY, Beh BK, San Liang W, Ky H, Yousr AHN et al. *Vernonia amygdalina*, an ethnoveterinary and ethnomedical used green vegetable with multiple bioactivities. *Journal of Medicinal Plants Research*. 2010; 25: 2787-2812. doi: Nil.
19. Evbuomwan L, Chukwuka EP, Obazenu EI, Ilevbare L. Antibacterial activity of *Vernonia amygdalina* leaf extracts against multidrug-resistant bacterial isolates. *Journal of Applied Sciences and Environmental Management*. 2018; 22(1): 17-21. doi: 10.4314/jasem.v22i1.4

20. Kaur D, Kaur N, Chopra A. A comprehensive review on phytochemistry and pharmacological activities of *Vernonia amygdalina*. Journal of Pharmacognosy and Phytochemistry. 2019; 8(3): 2629-2636. Corpus ID: 231400427.
21. Obiefule IE, Nwankwo EN, Ogbuefi EO, Uzochukwu CU. Toxicity of *Vernonia amygdalina* leaf powder against *Callosobruchus maculatus* (F.) (Coleoptera: Chrysomelidae) on stored cowpea (*Vigna unguiculata* L. Walp). Acta Entomology and Zoology. 2025; 6(1): 200-205. doi: 10.33545/27080013.2025.v6.i1c.210
22. Atawodi SE, Atawodi JC. *Azadirachta indica* (Neem): A plant of multiple biological and pharmacological activities. Phytochemistry Reviews. 2009; 8: 601-620. doi: 10.1007/s11101-009-9144-6
23. Gupta SC, Prasad S, Tyagi AK, Kunnumakkara AB, Aggarwal BB. Neem (*Azadirachta indica*): An Indian traditional panacea with modern molecular basis. Phytomedicine. 2017; 34: 14-20. doi: 10.1016/j.phymed.2017.07.
24. Wylie MR, Merrell DS. The antimicrobial potential of the Neem tree *Azadirachta indica*. Frontiers in Pharmacology. 2022; 13: 891535. doi: 10.3389/fphar.2022.891535
25. Ezin V, Chabi IB. *Azadirachta indica*: Its biological, pharmacological, antidiabetic potential, and omics applications. In: Antidiabetic plants for drug discovery. 2022; (1-22). Apple Academic Press, New York, USA. ISBN: 9781003282938.
26. Negrelle RR, Gomes EC. *Cymbopogon citratus* (D.C.) Stapf: Chemical composition and biological activities [J]. Revista Brasileira de Plantas Medicinaias. 2007; 9(1): 80-97. doi: Nil.
27. Kiełtyka-Dadasiewicz A, Esteban J, Jabłońska-Trypuć A. Antiviral, antibacterial, antifungal, and anticancer activity of plant materials derived from *Cymbopogon citratus* (DC.) Stapf species. Pharmaceuticals. 2024; 17(6): 705. doi: 10.3390/ph17060705
28. Du X, Zhang M, Wang S, Li J, Zhang J, Liu D. Ethnopharmacology, chemical composition and functions of *Cymbopogon citratus*. Chinese Herbal Medicines. 2024; 16(3): 358-374. doi: 10.1016/j.chmed.2023.07.002
29. Ugbogu EA, Emmanuel O, Ebubechi UM, Dike DE, Chukwuebuka OB, Ibe C, et al. The ethnobotanical, phytochemistry, and pharmacological activities of *Psidium guajava* L. Arabian Journal of Chemistry. 2022; 15(5): 103759. doi: 10.1016/j.arabjc.2022.103759
30. Tousif MI, Nazir M, Saleem M, Tauseef S, Shafiq N, Hassan L, et al. *Psidium guajava* L. An Incalculable but Underexplored Food Crop: Its Phytochemistry, Ethnopharmacology, and Industrial Applications. Molecules. 2022; 27(20): 7016. doi: 10.3390/molecules27207016
31. Liu C, Jullian V, Chassagne F. Ethnobotany, phytochemistry, and biological activities of *Psidium guajava* in the treatment of diarrhea: A review. Frontiers in Pharmacology. 2024; 15: 1459066. doi: 10.3389/fphar.2024.1459066
32. Sharma AK, Gangwar M, Tilak R, Nath G, Sinha ASK, Tripathi YB, et al. Comparative in-vitro antimicrobial and phytochemical evaluation of methanolic extract of root, stem, and leaf of *Jatropha curcas* Linn. Pharmacognosy Journal. 2012; 4(30): 34-40. doi: 10.5530/pj.2012.30.7
33. Ale AO, Odesanmi OS, Magbagbeola OA. Effects of *Jatropha curcas* root extracts on the metabolic profile of mature female rats. Mediterranean Journal of Pharmacy and Pharmaceutical Sciences. 2024; 4(2): 23-29. doi: 10.5281/zenodo.11173840
34. Dégbé M, Tété-Bénissan A, Debierre-Grockiego F, Toubaté BB, Akakpo AMW, Débare H, et al. Evaluation of Antitoxoplasmic Potential of Selected Antimalarial Medicinal Plants from Togo: *Artemisia annua* L., *Jatropha curcas* L., *Moringa oleifera* Lam., and *Sarcocephalus latifolius* (Smith) Bruce. International Journal of Biochemistry and Research Review. 2025; 34(3): 134-153. doi: 10.9734/ijbrr/2025/v34i3993
35. Salim MN, Masyitha D, Harris A, Balqis U, Iskandar CD, Hambal M. Anti-inflammatory activity of *Jatropha curcas* Linn. latex in cream formulation on CD68 expression in mice skin wound. Veterinary World. 2018; 11(2): 99-103. doi: 10.14202/vetworld.2018
36. Sulaimon LA, Ola-Mudathir FK, Abdullahi BA, Mukhtar AI, Obuotor EM. A comparative study on antioxidant and cytotoxic potentials of methanol extract of *Jatropha curcas* seeds and leaves. Journal of Drug Delivery and Therapeutics. 2021; 11(6-S): 43-48. doi: 10.22270/jddt.v11i6-S.5220
37. Saleh ZM, Abdel Azeiz AZ, Mehany A, El-Swaify ZA. Anticancer and antimicrobial activity of *Jatropha*'s leaves extracts. Egyptian Journal of Botany. 2023; 63(2): 621-634. doi: 10.21608/EJBO.2023.162736.2137
38. Chekka SV, Mantipelly NK. *Momordica charantia*: A natural medicinal plant. GSC Biological and Pharmaceutical Sciences. 2020; 12(2): 129-135. doi: 10.30574/gscbps.2020.12.2.0251
39. Dah-Nouvlessounon D, Chokki M, Noumavo ADP, Cârâc G, Furdui B, Sina H, et al. Ethnopharmacological value and biological activities via antioxidant and anti-protein denaturation activity of *Morinda lucida benth* and *Momordica charantia* L. leaves extracts from Benin. Plants. 2023; 12(6): 1228. doi: 10.3390/plants12061228
40. Iyekowa O, Ukpebor JE, Ehigimeto E. Physicochemical analysis of *Terminalia catappa* (Almond) seed nuts grown in Benin City, Nigeria. Mediterranean Journal of Pharmacy and Pharmaceutical Sciences. 2023; 3(1): 6-11. doi: 10.5281/zenodo.7771322

41. Oyelere SF, Ajayi OH, Ayoade TE, Pereira GBS, Owoyemi BCD, Ilesanmi AO, Akinyemi OA. A detailed review on the phytochemical profiles and anti-diabetic mechanisms of *Momordica charantia*. *Heliyon*. 2022; 8(4): e09253. doi: 10.1016/j.heliyon.2022.e09253
42. Richter E, Geetha T, Burnett D, Broderick TL, Babu JR. The effects of *Momordica charantia* on type 2 diabetes mellitus and Alzheimer's disease. *International Journal of Molecular Science*. 2023; 24(5): 4643. doi: 10.3390/ijms24054643
43. Karigidi KO, Ojebode ME, Olaiya CO. Effect of cooking on antioxidant and enzyme activity linked to carbohydrate metabolism and lipid peroxidation of eggplant (*Solanum melongena*). *Pertanika Journal of Tropical Agricultural Science*. 2018; 41(4): 1717-1730. Corpus ID: 208228494.
44. Kim DO, Jeong SW, Lee CY. Antioxidant capacity of phenolic phytochemicals from various cultivars of plums. *Food Chemistry*. 2003; 81(3): 321-326. doi: 10.1016/S0308-8146(02)00423-5
45. Park Y-S, Jung S-T, Kang S-G, Heo BK, Arancibia-Avila P, Toledo F, et al. Antioxidants and proteins in ethylene-treated kiwifruits. *Food Chemistry*. 2008; 107: 640-648. doi: 10.1016/J.FOODCHEM.2007.08.070
46. Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Analytical Biochemistry*. 1999; 26(9): 337-341. doi: 10.1006/abio.1999.4019
47. Benzie F, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*. 1996; 239(1): 70-76. doi: 10.1006/abio.1996.0292
48. Gyamfi M, Yonamine M, Aniya Y. Free radical scavenging action of medicinal herbs from Ghana: *Thonningia sanguine* on experimentally induced liver injuries. *Genetic Pharmacology*. 1999; 32 (6): 661-667. doi: 10.1016/s0306-3623(98)00238-9
49. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free radical biology and medicine*. 1999; 26(9-10): 1231-1237. doi: 10.1016/s0891-5849(98)00315-3
50. Worthington V. Alpha amylase. *Worthington enzyme manual* (36-41). New Jersey, United States: Worthington Biochemical Corporation. 1993; CAS: 9000-85-5. EC: 3.2.1.1.
51. Apostolidis E, Kwon YI, Shetty K. Inhibitory potential of herb, fruit, and fungal-enriched cheese against key enzymes linked to type-2 diabetes and hypertension. *Innovative Food Sciences and Emerging Technologies*. 2007; 8(1): 46-54. doi: 10.1016/j.ifset.2006.06.001
52. Tugume P, Nyakoojo C. Ethno-pharmacological survey of herbal remedies used in the treatment of pediatric diseases in Buhunga parish, Rukungiri District, Uganda. *BMC Complementary and Alternative Medicine*. 2019; 19: 353. doi: 10.1186/s12906-019-2763-6
53. Obel MA, Asfoor TAT. The potential of endemic medicinal plants in the central regions of Abyan Governorate, Yemen, for sustainable pharmaceutical applications. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(3): 46-51. doi: 10.5281/zenodo.16537815
54. Nisa Z, Sabir SM, Ahmad Z, Akhaq M. Analysis of hepatoprotective and antioxidant activities, and determination of the phenolic profile of *Moringa oleifera*. *Mediterranean Journal of Medical Research*. 2026; 3(1): 86-93. doi: 10.5281/zenodo.18912004
55. Han KSS, Win KT. Chemical compositions, nutritional values, total phenolic content, and antioxidant activity from *Eleocharis dulcis*. *Mediterranean Journal of Medicine and Medical Sciences*. 2026; 2(1): 28-35. doi: 10.5281/zenodo.18636582
56. Nkollo MI, Ngwuode RN, Efejene IO, Olele CH, Iwelumo BC, Chibuogwu C, Aisuodionoe EM. Phytochemistry and pharmacological insights into *Kalanchoe pinnata*: A brief review. *Mediterranean Journal of Medical Research*. 2025; 2(1): 26-31. doi: 10.5281/zenodo.15368110
57. Narayanan Z, Glick BR. Secondary metabolites are produced by plant growth-promoting bacterial endophytes. *Microorganisms*. 2008; 10(10): 2008. doi: 10.3390/microorganisms10102008
58. Mehmood A, Javid S, Khan MF, Ahmad KS, Mustafa A. *In vitro* total phenolics, total flavonoids, antioxidant and antibacterial activities of selected medicinal plants using different solvent systems. *BMC Chemistry*. 2022; 16: 64. doi: 10.1186/s13065-022-00858-2
59. Sebastiani G, Navarro-Tapia E, Almeida-Toledano L, Serra-Delgado M, Paltrinieri AL, García-Algar Ó, Andreu-Fernández V. Effects of antioxidant intake on fetal development and maternal/neonatal health during pregnancy. *Antioxidants*. 2022; 11(4): 648. doi: 10.3390/antiox11040648
60. Karigidi KO, Akintimehin ES, Omoboyowa DA, Adetuyi FO. Phytochemical profiling and antioxidant activities of *Monodora myristica* and *Dennettia tripetala* against lipid peroxidation in rat heart. *Acta Pharmaceutica Scientia*. 2022; 60(1): 49-63. doi: 10.23893/1307-2080.APS.6004

61. Adeleke PA, Annafi OS, Ajayi AM, Ben-Azu B, Okubena O, Umukoro S. Sorghum bicolor-based supplement reduces oxidative stress and pro-inflammatory cytokines to mitigate rotenone-induced Parkinsonian-like motor dysfunctions in rats. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2024; 4 (3): 15-26. doi: 10.5281/zenodo.13309953
62. Mwamatope B, Tembo D, Chikowe I, Kampira E, Nyirenda C. Total phenolic contents and antioxidant activity of *Senna singueana*, *Melia azedarach*, *Moringa oleifera*, and *Lannea discolor* herbal plants. *Scientific African*. 2020; 9: e00481. doi: 10.1016/j.sciaf.2020.e00481
63. Chiang JL, Maahs DM, Garvey KC, Hood KK, Laffel LM, Weinzimer SA, et al. Type 1 Diabetes in children and adolescents: A position statement by the American Diabetes Association. *Diabetes Care*. 2018; 41(9): 2026-2044. doi: 10.2337/dci18-0023
64. Jebiril AO, Rbeida OA, Younis HM. Alloxan blood level and the risk of diabetes mellitus in children. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2025; 5(2): 1-7. doi: 10.5281/zenodo.15116795
65. Katsarou A, Gudbjörnsdóttir S, Rawshani A, Dabelea D, Bonifacio E, Anderson BJ, et al. Type 1 diabetes mellitus. *Nature Reviews, Disease Primers*. 2017; 3: 17016. doi: 10.1038/nrdp.2017.16.
66. Shrestha D, Sharma P, Adhikari A, Mandal AK, Verma A. A review on Nepalese medicinal plants used traditionally as alpha-amylase and alpha-glucosidase inhibitors against diabetes mellitus. *Current Traditional Medicine*. 2021; 7(5): e02072119443. doi: 10.2174/2215083807666210702144103

Acknowledgments: The study received funding from the Tertiary Education Trust Fund (TETFUND) under the 2022 IBR grant.
Author's contribution: KOK, AOI & FMO conceived and designed the study. All authors contributed to the data collection and analysis. KOK, AOI & FMO contributed to data analysis and interpretation and drafted the manuscript. All authors approved the final version of the manuscript and agreed to be accountable for its contents.
Conflict of interest: The authors declare the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
Ethical issues: The authors have considered ethical issues, including plagiarism, informed consent, data fabrication or falsification, and double publication or submission.
Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.
Generative AI disclosure: No Generative AI was used in the preparation of this manuscript.

الفعالية البيولوجية، ومضادات الأكسدة، والقدرة التثبيطية للنباتات الطبية المستخدمة في علاج الرضع ضد إنزيمي الأميليز والجلوكوزيداز

كاويدي أو. كاريدي 1*، أبيولا أو. إيلوري 2، فونميلولا م. أوجو 3، وألوداميلولات. أتورو 3

- 1 قسم الكيمياء الحيوية، جامعة أولوسيجون أغاجو للعلوم والتكنولوجيا، أوكيتيوبا، ولاية أوندو، نيجيريا
 - 2 قسم العلوم الفيزيائية، جامعة أولوسيجون أغاجو للعلوم والتكنولوجيا، أوكيتيوبا، ولاية أوندو، نيجيريا
 - 3 قسم العلوم البيولوجية، جامعة أولوسيجون أغاجو للعلوم والتكنولوجيا، أوكيتيوبا، ولاية أوندو، نيجيريا
- * المؤلف المسؤول عن المراسلات

المخلص: لطالما شكلت النباتات الطبية جزءًا هامًا من الرعاية الصحية منذ القدم. وتستخدم في الطب الشعبي لعلاج صحة الرضع والحفاظ عليها. في هذه الدراسة، تم البحث عن النشاط الحيوي، ومضادات الأكسدة، والقدرة التثبيطية لإنزيمي الأميليز والجلوكوزيداز في تسعة نباتات طبية (كاريا بابايا، مانجيفيرا إنديكا، أوسيموم غراتيسيموم، فيرنونيا أميغدالينا، أذيراختا إنديكا، سيمبوجون سيترات، بسيدوم جوافة، جاتروفا كركاس، ومومورديكا شارانتيا). تم تقييم النشاط الحيوي باستخدام فحوصات الفينولات الكلية والفلافونويدات الكلية. استخدمت السعة الكلية لمضادات الأكسدة (TAC)، وقدرة اختزال الحديد المضادة للأكسدة (FRAP)، ونشاط إزالة جذور 1-ثنائي فينيل-2-بيكريل هيدرازيل (DPPH)، وقدرة إزالة جذور 2-أزوبيس-3-إيثيل بنزوثيرازولين-6-سلفونات (ABTS) لتحديد القدرة المضادة للأكسدة. في المقابل، قُيِّمت القدرة التثبيطية ضد نشاطي الأميليز (AMY) والجلوكوزيداز (GLU). أجريت جميع الاختبارات وفقًا للإجراءات القياسية. أظهرت النتائج أن مستخلص النيم (*Azadirachta indica* A.juss) يحتوي على أعلى نسبة من المركبات الفينولية الكلية (82.82 ± 1.57 ملغ مكافئ حمض الغاليك/100 غرام) وأعلى نسبة من الفلافونويدات الكلية (71.95 ± 1.19 ملغ كيرسيتين/100 غرام)، بينما كان مستخلص نبات فيرنونيا أميغدالين (*Vernonia amygdaline* Schreb) (54.08 ± 1.69 ملغ مكافئ حمض الأسكوربيك/100 غرام) ومستخلص المانجو (*Mangifera indica* L) (68.83 ± 0.09 ملغ مكافئ الحديد الثنائي/100 غرام) الأعلى من حيث القدرة الكلية المضادة للأكسدة (TAC) وقدرة اختزال الحديد (FRAP)، على التوالي. وأظهرت جميع المستخلصات تثبيطًا يعتمد على التركيز لكل من DPPH و ABTS و AMY و GLU. وتم حساب معامل التثبيط (IC50). أظهر نبات المانجو (*Mangifera indica* L) أقل قيمة IC50 لتثبيط DPPH (28.50 ± 0.57 ميكروغرام/مل) و ABTS (34.50 ± 0.88 ميكروغرام/مل)، بينما أظهر نبات الجاتروفا (*Jatropha curcas* L) أقل قيمة IC50 لتثبيط AMY (140.40 ± 1.08 ميكروغرام/مل) و GLU (85.50 ± 1.17 ميكروغرام/مل). تشير هذه النتائج إلى أن النباتات الطبية تمتلك خصائص حيوية ومضادة للأكسدة ومثبطة، مما يبرر استخدامها لدى الرضع.