



**IBEROAMERICAN
JOURNAL OF
MEDICINE**

iberoamericanjm

Journal homepage: www.iberoamericanjm.tk

Review

Factors affecting SARS-CoV-2 (COVID-19) Pandemic, including Zoonotic, Human Transmission and Chain of Infection. Reducing Public health Risk by Serum Antibody Testing, Avoiding Screening in Unhygienic Places and False PCR Reporting. A Scientific Review

Kamran Mahmood Ahmed Aziz^{a,}, Abdullah Othman^b, Waleed Abdullah Alqahtani^c, Sumaiya Azhar^d*

^a *Consultant Diabetologist and Research Scientist, Aseer Endocrine and Diabetes Center, Aseer Central Hospital, MOH, Saudi Arabia*

^b *Consultant Pediatric Endocrinologist and Director Aseer Endocrine and Diabetes Center, Aseer Central Hospital, MOH, Saudi Arabia*

^c *Consultant Adult Endocrinologist and Medical Director, Aseer Central Hospital, MOH, Saudi Arabia*

^d *Pharmacists and Medical Researcher, B. Clinical Pharmacy, University of Karachi, Pakistan*

* *Corresponding author.*

E-mail address: drkamran9999@yahoo.com

ISSN: 2695-5075 / © 2021 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

<http://doi.org/10.5281/zenodo.4599718>

ARTICLE INFO

Article history:

Received 10 February 2021

Received in revised form

04 March 2021

Accepted 10 March 2021

Keywords:

SARS-CoV-2

RT-PCR

Antibody

Zoonotic

Animal transmission

Genomics

Asymptomatic fraction

Herd immunity

ABSTRACT

During December 2019, a rapid increase in the number of SARS-CoV-2 (COVID-19) cases was reported worldwide. We investigated several factors for rapid increase in SARS-COV-2. Genomic sequence reveals that domestic and wild animals were likely ancestors and zoonotic source for SARS-CoVs, MERS-CoVs, and SARS-CoV-2; these viruses replicated in animals and humans during past several decades, exhibiting diverse mutations and self-limiting diseases except during outbreaks. SARS-CoV-2 has been retrospectively isolated in different studies in August 2019, several months before Wuhan reported. Hence, there is a possibility that viruses went undetected and infecting sub-clinically, in past several years, and SARS-CoV-2 antigens and neutralizing antibodies may have been present in humans since years. All SARS-CoVs are basically respiratory viruses, spread by droplets, hence droplet precautions are essential. Furthermore, silent phase of transmission (asymptomatic/subclinical) can be beneficial for humans. Lack of symptoms eventually lessen virus transmission and reduce the pathogen's long-term survival and provide strong humoral herd immunity (with seropositivity and diverse antibodies) up to several years and during epidemics. RT-PCR has low sensitivity and specificity, carries a high risk of handling live virus antigens, and requires difficult protocols. As viral load also sharply declines after few days of onset of infection, this technique might overlook infection. Furthermore, SARS-CoV-2 infection may be present in blood when oropharyngeal swabs are negative by RT-PCR. Conversely, antibodies against SARS-CoVs develop robustly in by reduced amount of antigens and ELISA for diagnosing antibodies demonstrates 100% specificity and 100% sensitivity, even in clinically asymptomatic individuals. These antibodies can be used for serologic surveys, monitoring and screening. Furthermore, screening tests for SARS-COV-2 should be avoided in unhygienic public places by nasopharyngeal swabs, which carry a high risk of further transmission, co-infection or super-infection. If above mentioned factors and Infection control policy is followed, SARS-CoV-2 pandemic can be controlled effectively.

© 2021 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE: Ahmed Aziz KM, Othman A, Alqahtani WA, Azhar S. Factors affecting SARS-CoV-2 (COVID-19) Pandemic, including Zoonotic, Human Transmission and Chain of Infection. Reducing Public health Risk by Serum Antibody Testing, Avoiding Screening in Unhygienic Places and False PCR Reporting. A Scientific Review. Iberoam J Med. 2021;3(2):138-160. doi: 10.5281/zenodo.4599718.

1. INTRODUCTION

During December 2019, one of the novel coronavirus infections SARS-CoV-2 (COVID-19) outbreak occurred in Wuhan, China. After that, several research papers were published, identifying the characteristics of this virus with nearly one thousand publications worldwide. However, most of these publications are based on the information from one aspect, not encompassing all the causative factors which are leading to the increasing frequency of the current pandemic.

Previous publications have not collected all reports with systemic review; and several biological transmission facts were missed. To limit SARS-CoV-2 (COVID-19) infections and pandemic emergencies, and to understand all scientific facts, here we have summarized different aspects of this disease with systemic review.

2. METHODS

To retrieve the research papers, the software of Reference Manager (Version 12) was used for internet search of literature and electronic databases, including PubMed/Medline, ISI Web of Knowledge, Z.39.50 gateway (international standard client-server communication protocols, and includes gateway to library catalogues). Furthermore, Google Scholar was also included in the search methodology. All the published papers were reviewed thoroughly, in details and the recent information regarding SARS-CoV-2 was extracted.

3. MOLECULAR BIOLOGY, IMMUNOLOGY AND GENETICS

3.1. HISTORY AND CORRECT NOMENCLATURE OF THE VIRUS FOR SCIENTIFIC STUDIES

Discovery of coronavirus took place in 1930s after domestic chickens were infected, resulting in acute respiratory infection, and demonstrated to be caused by a virus known as avian infectious bronchitis virus (IBV); later labelled as the coronavirus of the fowl or chicken (*Gallus gallus*). This coronavirus replicates in epithelium of upper, lower respiratory tracts and several other organs of the chicken, and is one of the important causes of economic loss for poultry industry. Virus was detectable in respiratory secretions and feces. This coronavirus was labelled 229-E. Now there is increasing evidence that coronavirus also infects species of birds other than chickens, such as bats, with astonishing molecular diversity [1-6]. Hence, it was then well known that coronaviruses are causing disease in several other animals including pigs, cows, chickens, dogs, and cats; for example Transmissible Gastroenteritis Virus (TGEV) and Porcine Epidemic Diarrhea Virus (PEDV). Recently, a novel coronavirus, called SW1, was identified in a deceased Beluga whale, which caused respiratory and liver failure [7].

The first human coronaviruses (HCoV) were discovered in the 1960s with demonstration that colds could be induced by nasal washing which did not contain rhinoviruses. Later in vitro experiments demonstrated the presence of coronaviruses with similar morphology as IBV [8-11]. Furthermore, it has been shown by scientist that human coronavirus can cause diseases beyond respiratory system, affecting multiple organ systems. This has been demonstrated in studies conducted on patients with multiple sclerosis, where coronavirus RNAs, and antigens were detected from the isolate of brain tissue. Additionally, scientists have proved human coronavirus gene expression in the brains of multiple sclerosis patients and labelled this phenomenon as "Infectious causes of multiple sclerosis" [12-18]. Current research literature demonstrates that all human coronaviruses (CoVs) are thought to originate from animal reservoirs with changes which have occurred in the genome during the course of propagation at different geographical locales. SARS-CoV (from bats via masked palm civet cats in China) and MERS-CoV (camels in the Middle East, seasonal especially during camel birthing) being prominent recent examples [19, 20].

Furthermore, and in fact, it has been demonstrated that

novel bat CoVs are likely ancestors for SARS-CoV and MERS-CoV, with discovery of hundreds of coronaviruses over the past decades [21]. It has been demonstrated in research that SARS-CoV originated in Chinese horseshoe bats which contain same genetic sequences of SARS-related CoVs with serologic evidence of a prior infection with a related CoV [22, 23].

Hence, it is very clear that coronavirus already existed and replicated in humans during past several decades, in multiple organs exhibiting its antigens during testing. This was also evident from the fact that diagnosis of coronaviruses was unnecessary as the disease was self-limiting and naturally completing its course with the exceptions during the outbreaks.

Initially, W.H.O (World Health Organization, an agency of the United Nations) named this virus as 2019-nCoV. On Feb 11, 2020, WHO renamed the disease as coronavirus disease 2019 (COVID-19), a rather unspecific name, not a scientific name, but a traditional approach. However, on the basis of phylogenetic analysis (genome sequence), and using a computational framework of comparative genomics, the Coronavirus Study Group (CSG) of the International Committee on Virus Taxonomy designated the scientific name to this naturally occurring virus as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), a universal nomenclature approach [24-28]. Coronaviridae Study Group of the International Committee on Taxonomy of Viruses has clearly stated that "The Study Groups quantify and partition the variation in the most conserved replicative proteins encoded in open reading frames 1a and 1b (ORF1a/1b) of the coronavirus genome" and that "Although these viruses were isolated at different times and locations from different human and animal hosts (with and without causing clinical disease), they all belong to the species *Severe acute respiratory syndrome-related coronavirus*, and their relationship parallels that between human individuals and the species *Homo sapiens*". Hence, SARS-CoV-2 should be the recommended name.

3.2. TAXONOMY AND MOLECULAR BIOLOGY OF SARS-COV-2

Coronaviruses (CoVs; order *Nidovirales*, family *Coronaviridae*, subfamily *Orthocoronavirinae*), are the largest group of enveloped positive-sense single stranded RNA viruses, belonging to the *Nidovirales* order, characterized by club-shaped spikes projecting from the cell surface of the virion, with unusually large complex

RNA genome [29-31]. Regarding genomic organization, Coronaviruses contain a non-segmented, positive-sense RNA genome of ~30 kilobase (kb). The genome contains a 5' cap structure along with a 3' poly (A) tail, allowing it to act as an mRNA for translation of the replicase polyproteins. The organization of the coronavirus genome is 5'-leader-UTR- replicase-S (Spike)-E (Envelope)-M (Membrane)-N (Nucleocapsid)-3' UTR-poly (A) (UTR=tail untranslated region). Considering Virion Structure, Coronavirus virions are spherical with diameters of approximately 125 nm. Virus particles consist of four main structural proteins that is, the spike (S), membrane (M), envelope (E), and nucleocapsid (N) proteins, all of which are encoded within the 3' end of the viral genome [32]. The *Coronaviridae* are further subdivided into four genera, the alpha, beta, gamma, and delta coronaviruses. Seven types of coronaviruses are known to infect humans, causing respiratory symptoms of various severities. HCoV-229E (α -CoV), HCoV-OC43 (β -CoV), and HCoV-NL63 (α -CoV) cause common cold. Since 2000, there has been three major worldwide health emergencies and severe crises, namely the 2003 SARS (severe acute respiratory syndrome; SARS-CoV, a β -CoV) outbreak, the 2012 MERS (Middle East respiratory syndrome; MERS-CoV; a β -CoV) outbreak, and the SARS-CoV-2 (2019-nCoV; a β -CoV) outbreak. Genomic structures of SARS-CoV and MERS-CoV are well studied previously. Moreover, since 2000, no vaccine has been developed so far. Virus initially attaches to the host cell receptor by S protein (the receptor-binding domain, RBD), and this interaction is responsible for further actions and disease process. Aminopeptidase N (APN) receptors are used by several α -coronaviruses. However, SARS-CoV and HCoV-NL63 target angiotensin converting enzyme 2 (ACE2) as their host cell receptor; while MERS-CoV enters into human cells via attaching to dipeptidyl-peptidase 4 (DPP4) receptor, then ultimately leading to translation of the replicase gene from the virion genomic RNA [33, 34].

3.3. EXISTENCE OF CORONAVIRUSES INCLUDING SARS-COV-2 IN HUMANS FOR THE PAST SEVERAL DECADES BY COMPARATIVE ANALYSIS OF GENOMIC DATA

Since the origin of SAR-CoV-2 or HCoV-19 (COVID-19), debates have arrived for its real origin and its historical existence with human's in the past. The fact that this virus is known to humans come from the evidence that SARS-CoV-2 is optimized and showed

high affinity to human ACE-2 receptors. In the history, SARS-CoVs also share same ancestor [35]. Coronavirus spike (S) glycoproteins facilitates virus entry into cells and that they are the main target of potentially neutralizing polyclonal antibodies produced by humans. Strikingly, structural similarities were found between SARS-CoV-2 S and SARS-CoV S glycoproteins confirming close relation between them, and both of them recognize ACE-2 to enter target cells. This phenomenon is the most likely due to natural selection and decades of their associations [36-38]. Current evidence also demonstrate that the RNA genomic sequence of SARS-CoV-2 (COVID-19) shows 89% nucleotide homology with bat SARS-CoV and 82% with human SARS-CoV, indicating that bat is the probable zoonotic source and this virus might have been existed since long time in animals and humans [39]. Past evidence also give proof that there may be a zoonotic reservoir or source (a *Rhinolophus affinis* bat and Malayan pangolins or *Manis javanica*), and identification of a potential intermediate host of SARS-CoV-2 will be of clinical significance [40, 41]. Furthermore, during its zoonotic course, SARS-CoV-2 has acquired mutations (recombination), making it more complex and difficult to identify. Studying zoonotic infectious pathology of pangolin genome aids in understanding that how SARS-CoV-2 has jumped into humans. [42, 43].

Furthermore, researchers have also compared the affinity of different SARS-CoVs. They have come to a conclusion that "2019-nCoV likely uses human ACE-2 less efficiently than human SARS-CoV (year 2002) but more efficiently than human SARS-CoV (year 2003). Because ACE-2-binding affinity has been shown to be one of the most important determinants of SARS-CoV infectivity; 2019-nCoV has evolved the capability to infect humans and some capability to transmit among humans. Alarmingly, our data predict that a single N501T mutation (corresponding to the S487T mutation in SARS-CoV) may significantly enhance the binding affinity between 2019-nCoV RBD and human ACE-2. Thus, 2019-nCoV evolution in patients should be closely monitored for the emergence of novel mutations at the 501 position (to a lesser extent, also the 494 position)" [44]. These findings, including with additional research literature [27, 28] suggest that the SARS-CoV-2 gradually evolved via mutations and was associated with humans since decades. Hence SARS-CoV-2 antigens and neutralizing antibodies may have been present in humans since long time and this may be the reason that

RT-PCR (reverse transcriptase-polymerase chain reaction) and other diagnostic techniques are detecting previous antibodies and antigens (cross reactivity) which is resulting in an increasing number of positive patients daily in spite of strict lock down policy globally.

This astonishing fact was revealed in one of the study where positive antibody reactivity with western blot analysis was observed [45]. It was demonstrated that a strong cross-reactivity of antibodies was observed between SARS-CoV-positive human plasma and SARS-CoV-2 rNP with 45 kD specific band observation. The study was successful in confirming this cross-reactivity via ELISAs (enzyme linked immunosorbent assay). It was concluded that these patients were infected previously with SARS-CoVs in 2002 epidemic. Hence, patients previously infected and positive for SARS-CoVs might exhibit SARS-CoV-2 antibodies for long periods (seropositivity) and also positive with RT-PCR (false positive), in the absence of marked clinical infection and symptoms. Hence, RT-PCR methods should be very precise, sensitive and interpreted cautiously that the patient is really positive enough to spread the infection.

The fact of evolutionary relationships is further strengthened by the other studies which reveal that this virus was genetically closely related (89.1% nucleotide similarity) to a group of SARS-like coronaviruses (genus Betacoronavirus, subgenus Sarbecovirus) that had previously been found in bats in China [46, 47]. The RBD of its spike protein were also found to be almost identical (73.8-74.9% amino acid identity). Furthermore, it was found that SARS-CoV-2 exhibited 100% amino acid similarity to bat SL-CoVZC45 in the nsp7 and E proteins, suggesting that bats act as natural reservoir for these coronaviruses and using same receptor for cell entry, the ACE-2. Furthermore, the amino acid sequences of the seven conserved replicase domains in "ORF1ab" were also 94.4% identical, with an overall genomic sequence identity of 96.2%, suggesting that the two viruses belong to the same species, *the SARS-related coronaviruses* [8, 48].

Interestingly, retrospective investigative studies done in France have shown that SARS-CoV-2 was present in northern France since before Wuhan (China) reported (in December 2019), in a patient with no history of travel to China. The study has reported independent SARS-CoV-2 introductions without local transmission [49]. According to the authors "we used the newly generated genomes to investigate the origins of SARS-CoV-2 lineages circulating in northern France". They have reported that clinical characteristics of SARS-CoV-2 patients were

different; a unique V367F (G22661T) mutation in the receptor binding domain of the Spike protein was observed, which was not observed in other genomes. The study has concluded that SARS-CoV-2 was present earlier in France. Similarly, in another study there is a report of a 42 year old male patient with a history of type-2 diabetes and asthma, traveling record to Algeria in August 2019, absence of travel history to China, was admitted in intensive care unit (ICU) in France with hemoptysis, cough, chest pain, headache and fever, evolving for four days with no etiological diagnosis [50]. With initial examination unremarkable, sputum negative for pathogens, lymphopenia, and the chest computed tomography (CT) imaging demonstrated bilateral pulmonary ground-glass opacities in the inferior lobes. He was discharged after antibiotic treatment. Of interest, one of his children presented with influenza like symptoms. According to hospital policy, all respiratory samples were stored and frozen at -80 °C to be used in future investigations. Medical records and respiratory sample of this patient was again investigated retrospectively for RT-PCR and which was proven to be positive for SARS-CoV-2. Hence, it can be concluded that the disease was already present previously in France and was circulating in asymptomatic individuals.

These observations clearly show that virus might be freely moving in different parts of the world, went undetected, with sub-clinical infections. If studies and screening could have been conducted in the past few years, or perhaps four to five years back, it may be possible to identify this virus earlier. Surprisingly, no increased death rates have been reported in the past and it may be possible that humans have already developed immunity and antibodies against this virus (seropositivity). Again, as discussed above, this is the reason for more emerging cases of SARS-CoV-2 (COVID-19) as these subjects were not screened previously. Further studies are required to investigate respiratory samples retrospectively for the past few years (or perhaps several years) to rule out the emergence, propagation and clinical characteristics of SARS-CoV-2 at multicenter level.

3.4. GENOMIC SEQUENCE, GENETIC DIVERSITY AND EVOLUTION OF HUMAN MERS AND SARS COVS

On January 5th, 2020, the first whole-genome sequence of the SARS-CoV-2 virus (Wuhan-Hu-1) was submitted in the U.S. National Center for Biotechnology Information (NCBI) GenBank. After that, hundreds of researchers explored the details of the genetic sequence

of this RNA virus with several hundreds of publications. Most of recent published papers are still investigating the evolutionary history of SARS-CoV-2. Thousands of global SARS-CoV-2 whole-genome sequences are now available on the data sharing platform hosted by the Global Initiative on Sharing All Influenza Data (GISAID). Virus evolution can be now studied or explored by data visualization tools like Nextstrain and CoV-GLUE. Further genomic epidemiology can be explored at GISAID (Global Initiative on Sharing All Influenza Data) database [51].

Nucleotide sequence substitution is one of the most important mechanisms of all viral evolution in nature, responsible for high mutation rate, rapid evolution of RNS viruses and resistance to most vaccines and antiviral drugs. Viruses encode enzymes responsible for replicating their DNA or RNA genomes. Viral RNA polymerases exhibits low fidelity (the intrinsic error rate) approximately 10^{-4} mutations per nucleotide copied, a rate higher than DNA viruses, a phenomenon of virulence, which is also explained by quasispecies [52-58].

Study done by Shen Z et al. have reported that "metatranscriptome sequencing on bronchoalveolar lavage fluid (BALF) samples from eight patients with SARS-CoV-2 found that the number of intrahost variants ranged from 0 to 51 with a median number of suggesting a high evolution rate of the virus" [59].

Other studies performed on genomic analysis for diversity and mutations have found eighty-six complete or near-complete genomes of SARS-CoV-2 and has revealed several mutations and deletions on coding and non-coding regions with missense mutations (a point mutation in which a single nucleotide change results in a codon that codes for a different amino acid) including ORF1ab polyprotein, 3' end of the genome, spike surface glycoprotein, matrix protein, nucleocapsid protein, and three mutations (D354, Y364, and F367) located in the spike surface glycoprotein receptor-binding domain. Moreover, it was reported that because spike surface glycoprotein has an important role in binding to host cell receptors, this leads to variable "antigenicity" and rapid mutations with genetic diversity of coronavirus SARS-CoV-2 [60, 61].

In fact, it was found that the mutation rate of SARS-CoV-2 was also comparable to Ebola virus. The mutation rate in the SARS-CoV genome was observed to be $0.80 - 2.38 \times 10^{-3}$ nucleotide substitution per site per year, similar and comparable to other RNA corona viruses, not unusual, and not higher than HIV (human

immunodeficiency virus). It should be noted that moderate to high mutation results in a high-level of intra-host RNA genome variants [62-67]. As mentioned above by a review of retrospective investigative studies on respiratory samples for SARS-CoV-2 [49, 50], the genomic diversity have been underestimated, and virus was present in different regions, went undetected in asymptomatic individuals (infecting sub-clinically). In other words, SARS-CoV-2 virus was changing genomic sequences, perhaps for several years, with development of immunity in humans without causing increased number of deaths. It might be possible that deaths caused by usual pneumonia or other respiratory illnesses were missed, went undetected by RT-PCR for SARS-CoV-2 for the past years. Cross reactivity between antibodies and antigens proves this scenario that patients currently positive with SARS-CoV-2 might be positive previously with other SARS-CoVs in the past decades [45].

4. HUMAN TO HUMAN TRANSMISSION, ANIMAL RESERVOIRS AND WILDLIFE ZONOTIC TRANSMISSIONS. A NEED FOR SERIOUS INTERVENTIONS TO STOP RECURRENT PANDEMICS

While considering the infectious diseases, it is important to know the exact mechanism of the disease transfer, animal reservoirs, zoonosis and vectors. If these factors are not studied very carefully, then factors considered for limiting the disease transmission cannot be guaranteed. After pandemic of SARS-CoV-2, several hundreds of news was spreading in social media regarding human to human transmissions by touching. While carefully reading the published research literature, some other conclusions also arise. So far, there are no conclusive case controls or randomized trials which demonstrate definitive human to human transmission in pandemics by usual touching each other. Shen Z et al., have reported that "By investigating a possible person-to-person spread event, we found no evidence for the transmission of intra-host variants" [59]. Moreover, most of the previous information on human-to-human transmission of influenza comes from studies of human inoculation with influenza virus and observational studies. In contrast, to influenza and SARS-CoVs, several remarkable human studies of rhinovirus and respiratory syncytial virus transmission are found in the literature and we can come up to a definitive conclusion on their transmission pattern [60].

Some other studies have raised the suspicion of human to

human transmission of SARS-CoV-2. One of the papers published in *New England Journal of Medicine* have raised such suspicion when they investigated one family (patient, his wife and a son) with travel history from Wuhan and returned to Vietnam. Father and son were positive for SARS-CoV-2 while wife was negative, although they were travelling together. Son was positive for SARS-CoV-2 because he shared the bed with his father. This raises the suspicion of droplet infection which he might acquire from his father in a small closed room, while wife remained negative. The authors have reported travelling history to four cities across Vietnam using various forms of transportation systems (including planes, trains, and taxis) with a total of 28 close contacts. However, all of them remain free of infection, which raises the concern regarding human-to-human transmission by usual touching [68].

Although it is still unclear whether the transmission through person-to-person contact occurs via large respiratory droplets, due to coughing and sneezing, as in SARS, or via fomites [69], however, precautionary measures are recommended. During the outbreak of MERS-CoV, The Fifth Meeting of the International Health Regulations Emergency Committee concerning MERS-CoV concluded that although the subject of transmission was serious, there was no evidence of sustained human-to-human transmission; and increased number of cases was due systemic weaknesses in infection prevention and control principles [70]. It was mentioned above regarding interesting retrospective investigation done on respiratory samples by Gambaro et al., and others [49, 50] that, even before Wuhan reporting, SARS-CoV-2 strains were already introduced and present in France in patients without history of travel to China and without local human to human transmission, perhaps in significant number of asymptomatic individuals for several months, and SARS-CoV-2 went undetected.

It is well known that strains of human coronavirus OC43 and 229E cause approximately 33 % of common colds and hospital-acquired upper respiratory tract HCoV infections (nosocomial infections) in premature newborns. Studies have also demonstrated detectable minor infectivity of HCoV-229E for up to three hours on various hospital surfaces and also on sterile latex surgical gloves, which are significant vector sources for hospital-acquired viral infections. However, rapid loss of virus infectivity was observed after disinfecting, drying or cleaning. These and other studies have shown that health care facilities are the source infection transmission, and

proper infection control protocols should be followed [71, 72].

Although there is considerable conflicting and limited data, it has been also shown that SARS and MERS are usually transmitted by large respiratory droplets ($\geq 10 \mu\text{m}$ in diameter) during coughing and sneezing [73]. From the past research experiences of nCoVs (novel coronaviruses such as SARS and MERS outbreaks), there is sufficient evidence of droplet transmission of infection, but no or limited evidence of sustained human-to-human transmission (through routes such as contact) [74]. Moreover, our current knowledge on transmission is based on a small number of cases and further clinical case control or randomized trials are required to understand transmission physics of nCoVs. Studies which have done intensive follow-up of close contacts with index cases have also failed to demonstrate onward transmission due to the limited evidence [75, 76]. Similarly, research reports from SARS-CoV-2 outbreaks also have shown that high reporting of cases was due to health care associated transmission, overcrowding in hospitals with lack of diagnostic tools, which raised the bias and high incidence reporting was notoriously unreliable. Hence, alone contact transmission may not be as important as influenza virus survives on hand for less than five minutes. Although hand washing has been demonstrated to reduce transmission of respiratory illness, but there is no specific scientifically based evidence which shows clear benefits of repeated hand washing for several times a day during epidemics [77, 78]. However, as a precautionary tool, especially in health care facilities or hospital, it is advisable for health care staff to wash their hand while dealing with sick patients. Hence, it can be concluded that droplet spread is the sole or primary route of transmission for all influenza viruses, including SARS-CoVs.

Finally, if current SARS-CoV-2 transmission pattern was a new, based on its mutations, then it can be considered to have a unique transmission pattern from human to human. However, this was not the case and currently there is no evidence to validate such theory in reality. However, and conversely, a recent study has demonstrated that SARS-CoV-2 genomic mutations did not increase the virus transmission capability. They conducted research from the data of 75 countries and did virus genomic analysis from 15,000 SARS-CoV-2 patients by a phylogenetic index. Instead, they concluded that these mutations are neutral primarily induced by RNA editing by human immune system, and this high rate of mutations do not appear to be benefiting the virus.

Currently there is no evidence that more transmissible lineages of SARS-CoV-2 will emerge [79]. Hence, these mutations appear to be a natural process and pathogen to human interactions and adaptations, occurring since long time, with no effect on current virus transmissibility.

In summary, it can be concluded that SARS-CoV-2 is not a new true contagious disease and which, is not transmitted "*solely*" by direct contact with an infected people, for example hand shake, etc., unlike other contagious diseases (e.g., chickenpox, smallpox, measles, leprosy, ringworm, gonorrhoea syphilis, etc.). However, studies do confirm SARS-CoV-2 transmission via Droplet infection (though coughing, sneezing and spitting of infected people) like other respiratory diseases, e.g., pneumonia diphtheria, influenza, tuberculosis, common cold, whooping cough, etc., which are more common in more crowded living conditions. In conclusion, mode of transmission of such diseases including influenza is via droplet [80]. Similarly, nosocomial infections and opportunistic airborne transmission (which may occur through fine particle aerosols) should also be considered. It was also reported that human-to-human transmission is usually documented in health care facilities during outbreaks or pandemics due to poor infection control measures [81-84].

The nosocomial spread in tertiary care hospitals is because of the fact that receptor for the SARS S-glycoprotein, ACE-2, is found in the lower respiratory tract and, while admitted in hospital, procedures such as intubation contribute to nosocomial spread [85]. Under all these conditions, precautionary measures are essential in hospital settings and intensive care units to avoid spread of droplet infections (droplet precaution), such as wearing mask, gloves, gown, keeping distances between beds at least eight meters, and avoiding poor hygienic conditions [69, 86-89]. Moreover, and interestingly, it has been shown that most of the patients with SARS-CoVs infection had a milder clinical course of the disease and usually do not transmit the virus because of insufficient adaptation of the civet cat virus RBD to human ACE-2 receptors, which is required for propagation of the infection of human cell lines [90, 91]. During pandemics, it has also been also shown that, as commonly thought, quarantine is of unproven reality. There are no research based trials on quarantine regarding influenza like viruses and previous experience with influenza and SARS-CoVs epidemics suggests that large scale quarantine of a given population is logistically and technically very difficult [92].

Interestingly, and conversely, one study has found that neither contact tracing nor isolation of infected people would prevent an epidemic of certain viruses, including the coronaviruses [93]. Hence, quarantine will not be necessarily must in diseases which are not contagious and there are no recommended guidelines published so far for such activities, but only precautions and infection control measures with quality assurance will be appropriate for health care workers working inside the hospitals [73]. It should also be noted that effectiveness of quarantine or isolation will be of limited value under poor isolation facilities, failure of infection control procedures and protocols for isolation ward and for isolate patients. Moreover, the direct and indirect costs involved in quarantine measures are of considerable importance which has economic, international trade, and environmental effects [94-97].

It has been observed that some quarantine and isolation centers and quarantine camps are so small or narrow that health care workers are frequently affected by droplet infections and if seronegative for SARS-CoV-2 subjects are kept there for temporarily monitoring, these normal individuals also may get infection during the isolation period in these isolation centers or camps because of poor or substandard infection control procedures which is alarming and dangerous. Nowadays, on commercial basis, five star hotels are also being offered which is not cost effective and unaffordable for the majority. It is the prime duty of governmental and international health agencies to look into this subject and to provide standard infectious control protocols and facilities for their citizens of all classes, without bias because SARS-CoV-2 is currently considered a pandemic health issue.

Although quarantining is carried out for the sake of caution or safety, however, current evidence does not support this concept because studies have demonstrated that subclinical infection in SARS-CoVs is almost nonexistent with almost absence of mildly symptomatic cases, without any serious health effects. Study done by Leung et al. have done serologic survey for immunoglobulin IgG against SARS-CoV in a representative sample of close contacts of all SARS patients by viral lysate enzyme linked immunosorbent assay (ELISA) and Positive results were confirmed with immunofluorescence assay (IFA) and neutralization tests (sensitivity of 100% and specificity of 95%). They reported that " of 1,068 samples analyzed, 2 (0.19%, exact 95% CI 0.02%–0.67%) had a positive titer (1:25 to 1:50 on IFA compared to at least 1:100 in most recovered SARS cases) for SARS-CoV IgG antibody.

Neither participant with a positive sample reported a chronic medical condition or being sick with febrile or respiratory illness from February to August. Both seropositive participants arose from two superspreading events in Hong Kong, i.e., Prince of Wales Hospital nosocomial outbreak and Amoy Gardens environmental point source community outbreak" [98-100]. They have discussed further about interspecies transmission from animals to humans that asymptomatic infection was observed in Guangdong animal traders (palm civets), who demonstrated an animal SARS-CoVs seropositivity rate of 72.7% (95% CI 49.8%–89.3%) in the absence of prior clinical disease [101]. They have concluded that "SARS rarely manifests as a subclinical infection, and at present, wild animal species are the only important natural reservoirs of the virus". In fact, animal reservoirs and zoonotic transmission are well known for SARS-CoVs since decades [102].

There is an evident and strong risk of pathogens emerging from animal reservoirs (including wild and domestic animals) that have a serious potential to infect humans. This risk is increased when humans are in close and continuous contact with animals. On the other hand, Insect vectors are also another source of infections for several other infectious diseases. It is also well known that coronavirus (CoVs) are long having been associated with animals, infecting them and causing serious respiratory disease. MERS-CoV severely infects Rhesus macaques (*Macaca mulatta*), and symptoms of this animal are similar to human infections [103-105]. Further evidence for the origin of SARS-CoV-2 from the bats comes from the fact that this virus uses the same cell surface receptor, the ACE2, as discussed above [40, 59]. Recent studies have confirmed and isolated coronavirus from several other animals as well, including domestic rabbits [106-109]. Moreover, in a recent study it has been shown that novel SARS-CoV-2 (COVID-19) has a 96 % genomic resemblance with a bat *Rhinolophus affinis*, and considered the origin of epidemic due to animal to human zoonotic transmission [110].

Studies published early in 2020 have given strong evidence that now domestic animals, such as ferrets and cats are prone to develop SARS-CoV-2 infection with rapid transmission. Their nose, throat, and lower respiratory secretions demonstrate high viral titers and neutralizing antibodies (against s-protein) are detectable in their sera [111-113]. This alarming fact should be considered by global health authorities. This might be the reason of increased number of SARS-CoV-2 cases in the community.

Hence, to control transmission to humans, animal reservoirs must also be investigated. Further extensive research at multicenter level is required to know more about coronavirus' animal reservoir and to explore the exact mechanism of zoonotic transmission and chain of infection to prevent the spread of SARS-CoVs to humans.

It is one of the great responsibilities of international health organizations to conduct research and to make decisions or guidelines only on published randomized controlled trials for patients' safety, security and minimizing the health risk. For example, statement for direct contact transmission was based on one of a research letter (not the original research), which was done only for 10 days on three patients [114, 115]. The authors of this study themselves have stated in their letter that "This study has several limitations. First, viral culture was not done to demonstrate viability. Second, due to operational limitations during an outbreak, methodology was inconsistent and sample size was small." Most of such studies conducted are assumed or presumed, which leads to further spreading of false news in social media which poses public health risks due to malpractice. Hence, international health agencies should conduct original research trials in their research facilities worldwide and not to rely on pilot studies or letters. Also, they should do research to explore more about animal reservoirs and mechanisms of coronavirus zoonotic transmission to humans.

5. DEVELOPMENT OF ANTIBODIES AGAINST SARS-CVOS IN HUMANS AND THEIR SEROLOGIC DETECTION FOR DIAGNOSIS WITH HIGH SPECIFICITY AND SENSITIVITY WITHOUT THE RISK OF HANDLING LIVE VIRUS

Neutralizing antibodies (IgG, IgM, and IgA) develop rapidly with high-level of efficacy and safety, against SARS-CoVs in human population conferring immunity [116-118]. It has been shown that the enzyme-linked immunosorbent assay (ELISA) can be used for serosurveillance of coronavirus related SARS-CoVs which can detect antibodies in clinically asymptomatic individuals with 100% specificity. This is highly specific and can detect SARS-CoVs antibodies even when blood is infected by other organisms.

The ELISA is developed by epitope mapping, using synthetic peptides from the spike, membrane, and nucleocapsid (S, M, and N) proteins of the virus protein sequences of SARS-associated coronavirus. As compared to ELISA, RT-PCR (reverse transcriptase-

polymerase chain reaction) will be inferior in such cases because viral load sharply declines after nine days of disease onset and RT-PCR carries a high risk of handling infectious live virus. RT-PCR may also be exhausting as it requires strict criteria for confirmation of positive results, the test has to be done on at least two different samples, with controls (positive and negative), have to be repeated at least twice and confirmed with different laboratories if the result is positive. Furthermore, RT-PCR requires sophisticated instruments or equipment, highly trained laboratory personals, with very high laboratory quality-control standards and which may be not possible when mass screening for population is considered.

The techniques for identification of seroconversion to SARS-CoVs by ELISA is one of the definitive and accurate criteria, is preferred and standard for retrospective detection of infection by SARS-CoVs. Immunoassay for SARS-CoVs includes ELISA or western blot (with virus whole antigen or recombinant antigens), IFA (immunofluorescence assay; using whole virus), and detection of specific neutralizing antibodies by precise methods. All SARS-CoVs patients demonstrate increase in IgG (immunoglobulin G) within few days and anti-SARS-CoVs IgG persistently remain by more than three to six months. [119-130]. In research studies, diagnostic sensitivity of ELISA has been shown to be 100% with specificity of 100% with no cross-reactivity detected with common non-coronavirus respiratory pathogens, such as OC43 and 229E. Immunological studies and serological assays have demonstrated that detection of antibodies in plasma or serum via ELISA allow for screening and diagnosis of SARS-CoV-2 (COVID-19) as early as two days after onset of symptoms, without the risk of handling live infectious virus. Moreover, research trials have demonstrated that serological assays or studies helps study the immune responses to SARS-CoV-2 in a qualitative and quantitative manner and can be used for cohort or population studies or screening to determine seroprevalence or previous exposure in a given population [119, 131]. Researchers have shown that such serosurveys are essentially diagnostic and determine the precise rate of infection in an affected region and determine that how many people are immune to SARS-CoV-2 in affected area, which may be an important strategy for health care and hospital workers.

It should be noted that immune responses to SARS-CoV-2 are identical to SARS-CoVs and MERS-CoVs. In fact, it has been shown that individuals infected with coronaviruses or SARS-CoVs specifically demonstrate neutralizing antibodies in the serum with protection from reinfection and acquired immunity up to years [128, 132-134]. Hence, by detecting antibodies in a given population, we can safely determine that such people in a given community have acquired immunity against SARS-CoV-2 and "can go safely back to their work and normal routine". In fact, antibody response to SARS-

CoVs and MERS-CoVs is usually robust, healthy and persistent, significantly correlating with viral or antigen load [134, 135]. Similarly, Antibodies (IgG and IgM) to the new SARS-CoV-2 has also been demonstrated to be persistence for long periods [136]. Moreover, a reduced amount of viral antigen is sufficient to trigger a humoral antibody response, and is not blunted further by antibody response or reaction [137]. Additionally, studies have shown that S-proteins (spike proteins) of SARS-CoVs are the main target of Neutralizing antibodies and that sensitivity of S-protein based ELISA were higher in detecting antibodies and are optimally recommended as screening test.

Furthermore, neutralizing activity was detectable in 89% of patients recovered at 36 months, and even at that time S-protein antibody detection rate was 100%, indicating sustainable neutralizing activity. Hence, antibodies confers not only persistent immunity, but also helpful in screening and diagnosing the SARS-CoVs [138]. Additionally, current research studies have demonstrated that SARS-CoV-2 infection may be present in blood when oral swabs are negative for SARS-CoV-2 and the researchers have recommended serological testing for confirmation and future epidemiological testing at large scale.

Serological assays (microneutralization assays) with specificity and sensitivity without cross reactivity are now developed and approved in the USA for screening antibodies to SARS-CoV-2 and distributed to 200 laboratories worldwide [131, 139-143]. Further research trails are also required to study and confirm the details of above-mentioned research trails. Hence, understanding the basic mechanisms of acquired immunity is an urgent issue which will make future strategic planning and relaxation in social distancing protocols [144].

The passive immunization or the technique of using polyclonal or monoclonal antibodies, which have neutralizing activity against SARS-CoVs, via convalescent serum (convalescent plasma therapy) from SARS patients to control the severe disease and its transmission, have possible benefits. Passive transfer of immune serum has been tested in experimental mouse model of SARS-CoV infection. Furthermore, commercial production of SARS antiserum containing antibodies has also been applied in clinical practice [23, 145-151]. However, for mAb (monoclonal antibody) prophylaxis against SAR-CoVs or SARS-CoV-2 has conditions which must be met before initiating this therapy. First, it is a time-consuming method and availability of neutralizing antibodies (Nabs) in the market is difficult and challenging. Secondly, within the group of SARS-CoVs, RBD demonstrates highest variability isolated from animals [6, 23]. Single mAb may not be enough to protect against all genomic strains of the SARS-CoVs, and genotyping of SARS-CoVs is recommended for selecting optimal or appropriate neutralizing mAbs when a new outbreak is detected. It is, therefore, advisable to investigate prior whether SARS-

CoV quasispecies generation in vitro cell culture will affect the outcome of virus neutralization antibodies and assays. [152-155].

Moreover, effect of passive antibody treatment and its immune response while protecting against SARS-CoVs is debatable and controversial because patients who died of SARS demonstrated acute antibody and immune response. As compared to active immunization and until vaccines are available, great caution is required in case of plasma therapy because neutralizing antibodies are associated with strong antibody agglutination reaction with hypercytokinemia, pulmonary pro inflammatory accumulation and fatal acute lung injury (ALI) [156]. In summary, however, human antibodies to SARS-CoVs are protective for both donors and patients, provided the procedure is done according to standard recommendations. If vaccines are developed to confer permanent immunity, SARS-CoV-2 infection can be eliminated [157].

6. ASYMPTOMATIC TRANSMISSION FROM HUMAN TO HUMAN, THE ASYMPTOMATIC FRACTION, SELF-LIMITING COURSE: INFLUENZA IN CONTEXT

Here the literature and significant studies from 1800 onward were reviewed. Influenza was and still considered one of the most important causes of morbidity and mortality in elderly patients. In a period of 1957-1960, over 86,000 deaths occurred in the United States as a result of influenza. Patients above the age of 65 with associated diseases were at higher risk [158, 159]. The death rate from influenza in England and Wales were relatively low during the past decades because of general use of antibiotics. After the first outbreak in 1847-1848 with English death rate of 459 per million, the rate declined after 1850s. In the pandemic of influenza 1918-1919 during World War 1, the crude death rate of 3129 per million were observed.

Worldwide, this Spanish flu (La Grippe) was one of the deadliest pandemic in human history and about 500 million people (approximately one third of the world population) were infected with H1N1 influenza virus. In one year, more people died of influenza than four year plaque from 1347 to 1351. The observed death rate was about 100 million worldwide. 675,000 deaths occurred only in the United States. After 1936, it has been remarkably constant [160, 161]. Hence, decline in attack episodes can be observed in any pandemic infectious disease, due to several factors, including humoral immunity, and treatments (antibiotics or vaccines) [162]. The protection may be greater if influenza virus strains and genomic sequence of RNAs match closely the

vaccine strain. Globally, it has been reported that there is a higher mortality rate from seasonal influenza associated severe respiratory illness (H3N2 and H1N1 viruses) with a death rate of 290 000-650 000 [163-165]. Currently, during influenza epidemics in the United States, annually more than 36,000 death and 114,000 hospitalizations are observed [166].

As organizations relied more on vaccinations and mass screening was not done, there is possibility that people infected previously might harbor same antigenic genomic sequence as that of SARS-CoV-2 with humoral immunity or antibodies. In fact, and as mentioned above, in different studies it was demonstrated that SARS-CoV-2 existed in France before the reports came from Wuhan, China [49, 50]. Recently not much is known about influenza and therapies to prevent it. Currently, still new strategies are required for mitigating the severity of influenza pandemic and are now top health priority. Research and data from influenza pandemics has demonstrated that preventive measure during the peak of pandemic can significantly reduce the attacks, but has little effect on overall pandemic rates. Prophylaxis or vaccines can reduce influenza rates up to 75%.

Reviewing details by medical literature, it can be concluded that clinical characteristics of pandemic of SARS-CoVs or SARS-CoV-2 is similar in nature to previous influenza pandemics or attacks, requiring same precautions. Future pandemic strains of virus and pandemic transmission are uncertain in both cases, until further details are available [167-171]. In the absence of vaccination, silent transmission or asymptomatic transmission has an importance regarding the community health stability. An infection without any signs or symptoms is called asymptomatic fraction, asymptomatic or subclinical disease [172].

A study done by researchers at Princeton has found that this asymptomatic or silent phase of transmission can be a successful evolutionary strategy for viruses and SARS-CoV-2 (COVID-19). A silent infection has definitive short-term advantages. The study has provided some insights of silent transmission on the virus' long-term survival and how it could be beneficial to humans. In other words, lack of symptoms eventually lessen virus transmission and reduce the pathogen's long-term survival, which are called "host-parasite interactions" [173].

Hence, silent infection has advantages of providing humoral immunity, in spite of applying other strategies such as identification, contact tracing and quarantine which have economic impact and difficult

implementation. An asymptomatic individual can go to his normal work or routine and may come in contact with many susceptible people, giving a chance to develop immunity. Interestingly, a study conducted in South Korea has used a unique model of epidemiology to study the dynamics of the pandemic by using SIR model (S, susceptible; I, infectious; R, recovered). They concluded that Public information disclosure targeting SARS-CoV-2 positive cases were more effective in limiting transmission than lockdowns in the region with 50 percent lower economic losses [174]. Similarly, strict lock down and quarantine policy should be revised as there are further studies demonstrating high percentage of asymptomatic individuals. A study conducted on cruise ship passengers has described 81% asymptomatic and the researchers have emphasized to ease lockdown restriction as some other strategy have to be adapted in future [175].

In contrast, a symptomatic person with fever and cough can self-isolate himself by staying at home for few days until recovery. By this way overall transmission can be reduced over time, and in fact, evolution favors such behavior. This methodology is of immense importance in epidemics where the virus is constantly changing its genomic characteristics, virulence and antigenicity with treatment complexities and vaccine development difficulties; in this scenario, only very sick people can isolate themselves in their homes and the remaining healthy subjects (asymptomatic, subclinical or asymptomatic fraction) can continue their work, with precautions, conferring wide variety of immunity to the community with seropositivity of diverse antibodies, and hence, limiting the epidemics. This technique may also substitute convalescent plasma therapy on large scale, which is costly, a time consuming complex laboratory process, and impossible to apply when a large population or nation is considered with the difficulty that single monoclonal antibody will not be enough to protect against all genomic strains of the SARS-CoVs [145, 148]. However, passive immunization or convalescent plasma therapy by neutralizing antibodies (NAb) may be reserved for neutralizing antigens and viremia load in severely ill SARS-CoV-2 patients admitted to ICU and considered or requiring mechanical ventilation [176, 177].

In another study, researchers have shown that there are several factors which affect early transmission by a newly infected host. Infection can persist longer in individuals who are less symptomatic, with reduced transmission. Three infectious stages are defined: fully

asymptomatic, less symptomatic, or fully symptomatic which are possible results of evolution. Hence, biostability can be achieved with time when humoral immunity develops or even by active vaccine immunizations. Furthermore, and as mentioned above, by a review of further intensive literature details, no definitive experimental studies can be found regarding person-to-person spread of flu by touching. In fact, not much is known about influenza in this context. Under such conditions, and as has been seen in past few decades, asymptomatic carriers play an important role in human adaptation for viral diseases and development of acquired immunity. It has been observed that 50%-77% of infections with seasonal flu are asymptomatic (subclinical, with mild temperature rise), which may be due to pre-existing partial immunity [166, 178, 179]. The influenza virus shedding can occur in asymptomatic individuals, and disease transmission may also occur, but not at the same rate as observed in symptomatic patients. This phenomenon is practically observed since even with high infectious control measures, epidemic flu progression may slow down, but cannot stop. However, it is advisable to take droplet precaution during working in public places and hospitals.

It should be noted that any viral disease, if become seasonal, will be less virulent as most of the people in the community will developed variety of divergent antibodies against diverse antigens. Studies have reported that SARS-CoV-2 is the 7th coronavirus which may become seasonal [144]. The other six were: four human coronaviruses causing seasonal common colds (alphacoronaviruses NL63 and 229E; betacoronaviruses HKU1 and OC43); the remaining two, SARS-CoV and MERS-CoV, from zoonotic reservoir, yet not become endemic so far. However, as we discussed above, SARS-CoV-2 might be circulating worldwide, unnoticed in previous years. There is a high possibility that SARS-CoV-2 caused some sort of seasonal flu or perhaps outbreak in certain communities or regions, and went undetected. There is a need to study the humoral response to SARS-CoV-2 if it becomes seasonal.

There are reports of influenza outbreak in the past several decades with several reports repeatedly. However, their previous events were clinically controlled and stable. As soon as the people of the community develop antibodies or herd immunity, even by vaccines, there is a less chance of influenza outbreak or becoming pandemic. One study has revealed that 65% and 75% of the children between the age group two to four years were seropositive for 229E and NL63, respectively.

Moreover, by six years almost all children were seropositive. This may be the result of repeated re-infection. This also supports the evidence that children possess seroprotection against homologous virus genotypes. Furthermore, In future, if adult from such population is infected with more severity, then this indicates lack of cross-protective antibodies or antigen exposure. Hence, herd immunity from a variety of antigens with resopositivity with divergent antibodies is the most important factor controlling outbreaks in community. However, older people with comorbid conditions are usually susceptible for any type of influenza infections and special care must be given with vaccines to reduce influenza morbidity and mortality. [180-182].

Current evidence suggests that in most of the patients, SARS-CoVs and other coronaviruses follows a self-limiting course. Severe disease is the most likely among older patients and those with comorbidities. Currently no specific antiviral therapies are available and studies have also shown that antiviral therapies did not accelerate clinical improvement and were proven unsuccessful [183-185]. In the current pandemic, cellular and humoral immunity with specific monoclonal antibodies developed by humans are the only natural source which can prevent infection spread as early as hours to two days and will remain up to several months to years, conferring immunity [156, 186]

With the passage of time when more humans develop immunity against SARS-CoVs including SARS-CoV-2, more resistance will be demonstrated by the herd immunity (either by acquired immunity or active immunization by vaccines). Similarly, if repeated viral infections occur in Humans, they can be combat later by memory B-cells, a response from immune system [187-191].

7. RT-PCR IN THE DIAGNOSIS OF SARS-COV-2, SAMPLE SPECIMENS AND PRACTICAL REALITIES AND POSSIBILITIES OF FALSE POSITIVE OR FALSE NEGATIVE, WITH HIGH RISK OF HANDLING LIVE VIRUS ANTIGENS

Before in detail of RT-PCR (or RT-qPCR), it is important to discuss the sample specimens which are being considered for RT-PCR. Because of respiratory complication of SARS-CoV-2 that is pneumonia, suggesting that mainly lower respiratory tract is infected and affected. Recently, trials have been published

regarding accuracy of different respiratory specimens, including broncho-alveolar lavage fluid (BALF), sputum and nasal swabs. In the current epidemic, nasopharyngeal swabs are being considered the first choice for taking samples for RT-PCR. This is because it is easy (although sometimes painful to reach to pharynx via nose) and cheap. BALF is practically difficult, require a trained staff, suction device, and is more painful at the same time and cannot be performed at large scale. On the other hand, studies have shown that (apart from BALF) sputum samples demonstrated highest positive (90%) rate at all clinical stages of SARS-CoV-2, followed by nasal swabs (73%) and then the throat swabs (61%). However, BALF showed 100% positivity [192]. Hence, studies have not recommended throat swabs for the virus detection and considered sputum as most accurate for laboratory diagnosis. Currently, RT-PCR method is used also for diagnosing SARS-CoVs from nasopharyngeal or oropharyngeal swabs.

Although RT-PCR (reverse transcriptase-polymerase chain reaction) has improved ability to detect pathogen and efforts are done to develop PCR for pan-species, Genus and family (which is not practical). However, still with highly variable RNA viruses it is unsuccessful and not possible for PCR to detect the genomic sequences and is highly "biased" with which only known pathogens or antigens can be detected. This is because the primers chosen will alter nucleic acids which will be amplified. Moreover, RNA viruses exhibit such an extensive genomic diversity that it is very difficult to design primer sets for PCR with high sensitivity and specificity. Because of inadequacy of any single RNA primer pair to detect all human viruses (or range of viruses), several RNA primer sets are chosen to run sample analysis; but this increases the cost of the diagnostic methodology with complexity of the assay with difficulties of comparing results among different laboratories.[193-195].

In contrast, NGS (next generation sequencing) allows greater, unbiased and massive sequencing of genetic material. Furthermore, serologic assays are more sensitive and important in scenarios where genome (RNA) is difficult to isolate or is no longer present, and for cohort epidemiological studies [196-200]. As discussed above, RT-PCR has some other drawbacks (as viral load sharply declines after nine days of disease onset) and difficulties, requires re-confirmation, strict quality control, and carries a risk of handling infectious live virus. Mutation rates are also high among RNA

viruses and make them difficult to identify.

One of the most important issues with RT-PCR is the reporting of false negative and false positive results; as viral RNA genome in sufficient quantity is required to be detected by RT-PCR. Other important factor regarding false results is that the RT-PCR use primers, which can be affected by the variances or genomic diversity of viral RNA of SARS-CoV-2 (COVID-19). False negative results occur due to mutations in the primer and probe target regions in SARS-CoV-2 genome. Furthermore, currently several types of SARS-CoV-2 RT-PCR kits have been developed rapidly and available in the market with different quality. They were brought into the market without strict quality control measures. In fact, the sensitivity and specificity of real-time RT-PCR test is not very accurate or 100% like that of serology test [201, 202].

On February 12th 2020, a sudden rises in new cases of SARS-CoV-2 was observed. Researchers have shown that such increase in numbers was due to change in the diagnostic methodology, and indeed at that time, more than seven types of SARS-CoV-2 test kits were developed and approved rapidly and there were several factors attributed to their false negative rate (FNR). As an example, using primers in the ORF1ab gene and N genes are affected by the variation in the viral RNA genomic sequences. Additionally, regarding the natural history and collection of specimens, sampling techniques also contribute to high FNR. It was also demonstrated that in one testing scenario, FNR from one testing was as high as 30% to 50% in SARS-CoV-2 patients. It has been also reported that one of the patient was not confirmed SARS-CoV-2 infection by RT-PCR three times within three weeks until BALF was done and next generation sequencing (NGS) with RT-PCR were positive for SARS-CoV-2. This will not only increase the cost of testing, but also underestimate the true positive cases; and during the outbreak or pandemic, it is a serious issue [203, 204]. Moreover, there are events of false positive reports when tested on animals such as goat [205]. Such reports and testing kits should be further investigated to eliminate the public health risk. Hence, it is a serious issue and urgent call to rapidly improve the quality and standard of testing kits with their operative techniques for the accurate diagnosis of SARS-CoV-2.

High false positive rates are also a public health risk because such people will self isolates themselves and their office work will be affected. It is now well-known that sensitivity and specificity of RT-PCR are very poor

and false results are common. Recently, a study was conducted on RT-PCR false predictively among high risk patients who were exposed in inpatient departments and also health care workers to rule out SARS-CoV-2 infections [206]. They have reported false negative probability of 67% at initial days of infection and that precautions must be taken while interpreting the results. Hence, if suspicion is high, the RT-PCR should not be used alone for ruling out the diagnosis, but rather serological testing (especially IgG, which is produced rapidly after infection), should also be carried out and considered with clinical course of the disease [207].

If RT-PCR is considered for testing, then lower respiratory tract will be recommended (sputum, BALF), if possible, instead of nasopharyngeal swab because of genomic sequence analysis of SARS-CoV-2 and ACE2 viral receptors. Correct specimen selection is an important during pandemics. Blood specimens with serology testing are preferable over nasopharyngeal swabs. There are further reports globally that nasopharyngeal swabs (upper respiratory tract) were negative for pandemic (H1N1; novel swine-origin influenza A) 2009 influenza virus, but proved to be positive on BALF and some patients requiring admission to intensive care units (ICU) and mechanical ventilation were positive by bronchoscopic specimens [192, 208, 209]. These studies have important practical clinical implications that patient with influenza like illness with pneumonia and unexplained diagnosis should undergo bronchoscopic specimens.

An interesting recent study done on 1014 patients' cohort with SARS-CoV-2 (COVID-19) in radiology and image processing department has demonstrated that positive rates of RT-PCR throat swab assay were 59% while 88% for chest CT imaging with 97% sensitivity for the diagnosis of suspected SARS-CoV-2 cases. This study has concluded that CT imaging is more reliable and practical than RT-PCR in pandemic areas; RT-PCR showed low positive rates, lack of sensitivity, long processing time with unpredictable stability [210-214]. Similarly, a study discussed above on cruise ship passengers also reported a significant false-negative rate with RT-PCR testing [;Error! No se encuentra el origen de la referencia.].

Hence, serum based testing methods for SARSCoV-2-specific immunoglobulin-M will be more sensitive and specific. It was demonstrated that serologic test are gold standard for coronavirus testing during epidemics and showed a sensitivity of 0.96 (95% CI, 0.91 to 0.98) and a specificity of 0.96 (CI, 0.92 to 0.97) which was based on

SARS-CoVs disease clinical course, with the time from symptom onset to attending the clinic was 3.3 ± 2.6 days [204].

Studies have reported that patients showing symptoms of SARS-CoVs clinically and radiologically were further confirmed by RT-PCR with positivity of 68.2 %, while serology confirmation was 100% [215]. It is also worth noting that viral titers significantly diminish within few days after development of clinically significant disease both in humans and animals, indicating that serum antibody detection will be appropriate. Hence, according to these observations, serological testing further improves clinical case detection and must be used in combination with other detection methods [216-220].

Moreover, it has been demonstrated that in some exposed individuals RT-PCR may be negative and at the same time can have serologic evidence of antibodies against viruses. The study has shown that there was a 4-fold rise in antibody titer without signs and symptoms (subclinical infection) with RT-PCR negative (false negative) [172]. As discussed above, serological diagnosis of coronavirus infection is a good and recommended option. An indirect immunofluorescence assay (IFA), and enzyme linked immunosorbent assay (ELISA) are used to measure developing antibodies during the infection. The most appropriate test is immunochromatographic assay (ICA) [221-223]. With technological advancements in molecular biology and recombinant engineering, it is now possible to calculate and quantify virus particles with high accuracy in a given sample. These techniques include flow cytometry, dynamic light scattering, and quantitative capillary electrophoresis [224-227].

Furthermore, diagnosing coronavirus infection on RT-PCR is also not as simple as is thought because these human coronaviruses (such as SARS-CoV, MERS-CoV and SARS-CoV-2) are huge (around 30 kb); and production or generation of CoV infectious clones are hampered due to their huge size genome and the toxicity of some CoV replicase gene sequences during its propagation in bacteria. These problems can be overcome by bacterial artificial chromosomes (BACs) in vitro ligation of cDNA fragments, and using vaccinia virus as a vector for the propagation of CoV full-length cDNAs [228-231]. Moreover, usually 3-4 days are required for the final results, and then again test is repeated to confirm. The accessories and supplies (swabs, testing kits, and testing media) become short a while doing mass screening. Guide RNA can also recognize other interspersed RNS genomic sequences,

false positive results appear with loss of specificity. Inadequate sampling with low virus quantities, timing and site of collection (upper or lower respiratory tract), poor collection or handling, performance of kits (or substandard kits) are other issues which affect PCR results.

In conclusion, RT-PCR results must be cautiously interpreted. RT-PCR is also a complex, expensive and time-consuming technique if followed properly, specifically, with quality control measures, which is not possible at mass screening level.

8. TESTING SWARS-COV-2 IN UNHYGENIC, UNSTERILIZED PUBLIC PLACES UNDER, A RISK FACTOR FOR OF HUMAN TO HUMAN TRANSMISSION

For the past approximately five months, it has been observed that SARS-CoV-2 (COVID-19) testing is being done in public places, which are unhygienic and unsterilized. If it is believed that SARS-CoV-2 is highly infectious and transmittable, then precautions are necessary. A highly infectious virus should be isolated and tested in a highly sterilized laboratory. If not tested in sterilized laboratories, then the infection control principles are violated [232-234]. However, for the past few months the tendency of testing was observed to be increased in crowded public places, on the roads such as in motorcycles and cars, drive thru, field camps, markets which are unhygienic and source of reinfection. Taking nasopharyngeal samples in unsterilized conditions or public places there is also a serious risk of co-infection or superinfection by other viruses and bacteria [235-239].

Recently, it was observed that use of gloves by public was not according to recommendations. Usually, health care professionals and public assume that after wearing gloves they are fully protected. They usually touch objects freely, ignoring that they themselves will be infected by touching objects, their face and others as well. If a viral disease is of a serious concern and spreads by touching, then the use of gloves in such a manner will be dangerous. In this way, the gloves will be inoculated by several viruses and bacteria, and will further spread infections by ignorant people in the community. In such as case, hand sanitizing will be more appropriate. There are reports globally that if gloves are not used in an appropriate way, then they will be the source of infection and its further transmission [240-245].

It was also observed that the health care personals take

precautions for themselves only and not for the public. Health staff is touching cars and other materials during testing, while at the same time not following sterilization and infection control principles, changing their disposable gloves and gowns which is a dangerous and serious issue. While testing, if a person is assumed to be positive, then the health care staff will transmit the infection to the negative one if strict infection control principles are not followed. This is difficult to achieve in crowded public places, on the roads and small camps. Similarly, collecting and transferring the sample to test tube in open air may inoculate airborne viruses, causing cross reactivity with other viruses and false positive results. All these may be the other reasons of increasing frequency of infections all over the world, despite severe lockdown. While searching on the internet, several famous organizations, institutions and newspaper websites were discovered with more than 500 pictures showing testing being done in public places, outdoor camps which were unsterilized [246]. It is the urgent responsibility of international health agencies, health care authorities, officials and policy makers to understand and control such unhealthy and unethical behaviors. Furthermore, it was also observed that after severe lockdown of several hours, in some countries few hours are given to the citizens to purchase food, during which more crowding is observed. This is again a cause of increasing numbers of SARS-CoV-2, despite of precautions [247]. These situations and problems should be balanced.

9. CONCLUSIONS AND RECOMMENDATIONS

By the review of the genomic sequences of SARS-CoVs, it is concluded that these viruses existed in humans and animals since decades. Studies have demonstrated that SARS-CoV-2 infection existed in other regions several months before Wuhan announced, in August 2019. Hence, there is possibility that SARS-CoV-2 circulated in different parts of the world, infecting subclinically, and went undetected. This may be a reason of increasing number of cases all over the world, despite severe lockdown; as in previous years, mass screening was not done to detect this virus. Regarding SARS-CoV-2, animal sources cannot be excluded and animals might be a potential reservoir and can play active role in virus transmission. This is a second reason of increasing cases in certain communities. Hence, active research is required to investigate animal transmission of SARS-

CoVs and SARS-CoV-2. Similarly, Randomized controlled trials are required to investigate human to human transmission by touching, as there are very few studies with limited evidence and conflicting results. As all SARS-CoVs are basically respiratory viruses, droplet precautions and infection control measures are essential, especially for hospitals and health care staff. However, if a subject sneezes on his hand, sanitization of hand is mandatory or washing it by soap. Asymptomatic transmission from human to human usually occurs in communities. This asymptomatic or silent phase of transmission can be a successful evolutionary strategy for viruses including SARS-CoV-2. Lack of symptoms eventually lessen virus transmission and reduce the pathogen's long-term survival and silent infection has advantages of providing humoral immunity. Strategies such as identification, contact tracing and quarantine have economic impact and difficult to implement. Under such circumstances, asymptomatic people can continue their work with droplet precautions, and infection control measures while symptomatic or sick people can isolate themselves in their homes until clinical recovery. RT-PCR has low sensitivity and specificity, carries a high risk of handling live virus antigens, usually gives false negative and false positive results and must be interpreted cautiously. Furthermore, by handling live viral antigens in public places for testing, there is a high risk of virus spread in a community. This may be a third reason of increasing number of cases by false positive RT-PCR reporting. Conversely, highly specific antibodies are developed against SARS-CoVs and SARS-CoV-2, confer immunity, and can be used for serologic surveys, monitoring and screening with high specificity and sensitivity. However, testing and screening of SARS-COV-2 should be avoided in unhygienic public places by nasopharyngeal swabs, which carries a high risk of further transmission, and such highly infectious virus must be isolated and tested in highly sterilized laboratory.

10. REFERENCES

- Ambali AG, Jones RC. Early pathogenesis in chicks of infection with an enterotropic strain of infectious bronchitis virus. *Avian Dis.* 1990;34(4):809-17.
- Cavanagh D. Coronaviruses in poultry and other birds. *Avian Pathol.* 2005;34(6):439-48. doi: 10.1080/03079450500367682.
- Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, et al. Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science.* 2020;368(6494):1016-20. doi: 10.1126/science.abb7015.
- Poon LL, Chu DK, Chan KH, Wong OK, Ellis TM, Leung YH, et al. Identification of a novel coronavirus in bats. *J Virol.* 2005;79(4):2001-9. doi: 10.1128/JVI.79.4.2001-2009.2005.
- Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, et al. Bats are natural reservoirs of SARS-like coronaviruses. *Science.* 2005;310(5748):676-9. doi: 10.1126/science.1118391.
- Woo PC, Lau SK, Li KS, Poon RW, Wong BH, Tsoi HW, et al. Molecular diversity of coronaviruses in bats. *Virology.* 2006;351(1):180-7. doi: 10.1016/j.virol.2006.02.041.
- Mihindukulasuriya KA, Wu G, St Leger J, Nordhausen RW, Wang D. Identification of a novel coronavirus from a beluga whale by using a panviral microarray. *J Virol.* 2008;82(10):5084-8. doi: 10.1128/JVI.02722-07.
- Hamre D, Procknow JJ. A new virus isolated from the human respiratory tract. *Proc Soc Exp Biol Med.* 1966;121(1):190-3. doi: 10.3181/00379727-121-30734.
- Almeida JD, Tyrrell DA. The morphology of three previously uncharacterized human respiratory viruses that grow in organ culture. *J Gen Virol.* 1967;1(2):175-8. doi: 10.1099/0022-1317-1-2-175.
- Bradburne AF, Bynoe ML, Tyrrell DA. Effects of a "new" human respiratory virus in volunteers. *Br Med J.* 1967;3(5568):767-9. doi: 10.1136/bmj.3.5568.767.
- McIntosh K, Dees JH, Becker WB, Kapikian AZ, Chanock RM. Recovery in tracheal organ cultures of novel viruses from patients with respiratory disease. *Proc Natl Acad Sci U S A.* 1967;57(4):933-40. doi: 10.1073/pnas.57.4.933.
- Burks JS, DeVald BL, Jankovsky LD, Gerdes JC. Two coronaviruses isolated from central nervous system tissue of two multiple sclerosis patients. *Science.* 1980;209(4459):933-4. doi: 10.1126/science.7403860.
- Murray RS, Brown B, Brian D, Cabirac GF. Detection of coronavirus RNA and antigen in multiple sclerosis brain. *Ann Neurol.* 1992;31(5):525-33. doi: 10.1002/ana.410310511.
- Stewart JN, Mounir S, Talbot PJ. Human coronavirus gene expression in the brains of multiple sclerosis patients. *Virology.* 1992;191(1):502-5. doi: 10.1016/0042-6822(92)90220-j.
- Arbour N, Day R, Newcombe J, Talbot PJ. Neuroinvasion by human respiratory coronaviruses. *J Virol.* 2000;74(19):8913-21. doi: 10.1128/jvi.74.19.8913-8921.2000.
- Dessau RB, Lisby G, Frederiksen JL. Coronaviruses in brain tissue from patients with multiple sclerosis. *Acta Neuropathol.* 2001;101(6):601-4. doi: 10.1007/s004010000331.
- Gilden DH. Infectious causes of multiple sclerosis. *Lancet Neurol.* 2005;4(3):195-202. doi: 10.1016/S1474-4422(05)01017-3.
- Hansen GH, Delmas B, Besnardeau L, Vogel LK, Laude H, Sjöström H, et al. The coronavirus transmissible gastroenteritis virus causes infection after receptor-mediated endocytosis and acid-dependent fusion with an intracellular compartment. *J Virol.* 1998;72(1):527-34. doi: 10.1128/JVI.72.1.527-534.1998.
- Pertman S. Pathogenesis of coronavirus-induced infections. Review of pathological and immunological aspects. *Adv Exp Med Biol.* 1998;440:503-13.
- Weiss SR, Navas-Martin S. Coronavirus pathogenesis and the emerging pathogen severe acute respiratory syndrome coronavirus. *Microbiol Mol Biol Rev.* 2005;69(4):635-64. doi: 10.1128/MMBR.69.4.635-664.2005.
- He B, Zhang Y, Xu L, Yang W, Yang F, Feng Y, et al. Identification of diverse alphacoronaviruses and genomic characterization of a novel severe acute respiratory syndrome-like coronavirus from bats in China. *J Virol.* 2014;88(12):7070-82. doi: 10.1128/JVI.00631-14.
- Shi Z, Hu Z. A review of studies on animal reservoirs of the SARS coronavirus. *Virus Res.* 2008;133(1):74-87. doi: 10.1016/j.virusres.2007.03.012.
- Lau SK, Woo PC, Li KS, Huang Y, Tsoi HW, Wong BH, et al. Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. *Proc Natl Acad Sci U S A.* 2005;102(39):14040-5. doi: 10.1073/pnas.0506735102.
- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727-33. doi: 10.1056/NEJMoa2001017.
- WHO. Coronavirus disease 2019. Available from: <https://www.who.int/es/emergencias/diseases/novel-coronavirus-2019> (accessed May 2020).
- Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol.* 2020;5(4):536-44. doi: 10.1038/s41564-020-0695-z.
- Gralinski LE, Menachery VD. Return of the Coronavirus: 2019-nCoV. *Viruses.* 2020;12(2):135. doi: 10.3390/v12020135.
- Lauber C, Gorbalenya AE. Partitioning the genetic diversity of a virus family: approach and evaluation through a case study of picornaviruses. *J Virol.* 2012;86(7):3890-904. doi: 10.1128/JVI.07173-11.
- King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ. Virus taxonomy: classification and nomenclature of viruses. Ninth Report of the International Committee on Taxonomy of Viruses. Elsevier; 2011.
- Bárcena M, Oostergetel GT, Bartelink W, Faas FG, Verkleij A, Rottier PJ, et al. Cryo-electron tomography of mouse hepatitis virus: Insights into the structure of the coronavirus. *Proc Natl Acad Sci U S A.* 2009;106(2):582-7. doi: 10.1073/pnas.0805270106.
- Neuman BW, Adair BD, Yoshioka C, Quispe JD, Orca G, Kuhn P, et al. Supramolecular architecture of severe acute respiratory syndrome coronavirus revealed by electron cryomicroscopy. *J Virol.* 2006;80(16):7918-28. doi: 10.1128/JVI.00645-06.
- Maier HJ, Bickerton E, Britton P. Coronaviruses: methods and protocols. Springer; 2015.
- Belouzard S, Chu VC, Whittaker GR. Activation of the SARS coronavirus spike protein via sequential proteolytic cleavage at two distinct sites. *Proc Natl Acad Sci U S A.* 2009;106(14):5871-6. doi: 10.1073/pnas.0809524106.
- Kubo H, Yamada YK, Taguchi F. Localization of neutralizing epitopes and the receptor-binding site within the amino-terminal 330 amino acids of the murine coronavirus spike protein. *J Virol.* 1994;68(9):5403-10. doi: 10.1128/JVI.68.9.5403-5410.1994.
- Zeng F, Chow KY, Leung FC. Estimated timing of the last common ancestor of the SARS coronavirus. *N Engl J Med.* 2003;349(25):2469-70. doi: 10.1056/NEJM200312183492523.
- Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell.* 2020;181(2):281-92.e6. doi: 10.1016/j.cell.2020.02.058.
- Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol.* 2020;5(4):562-9. doi: 10.1038/s41564-020-0688-y.
- Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science.* 2020;367(6483):1260-3. doi: 10.1126/science.abb2507.
- Chan JF, Kok KH, Zhu Z, Chu H, To KK, Yuan S, et al. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect.* 2020;9(1):221-36. doi: 10.1080/22221751.2020.1719902.
- Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature.* 2020;579(7798):270-3. doi: 10.1038/s41586-020-2012-7.
- Zhang Z, Wu Q, Zhang T. Pangolin homology associated with 2019-nCoV. *bioRxiv.* 2020. doi: 10.1101/2020.02.19.950253.

42. Cui J, Li F, Shi ZL. Origin and evolution of pathogenic coronaviruses. *Nat Rev Microbiol.* 2019;17(3):181-92. doi: 10.1038/s41579-018-0118-9.
43. Fischer H, Tschachler E, Eckhart L. Pangolins Lack IFIH1/MDA5, a Cytoplasmic RNA Sensor That Initiates Innate Immune Defense Upon Coronavirus Infection. *Front Immunol.* 2020;11:939. doi: 10.3389/fimmu.2020.00939.
44. Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor Recognition by the Novel Coronavirus from Wuhan: an Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. *J Virol.* 2020;94(7):e00127-20. doi: 10.1128/JVI.00127-20.
45. Guo L, Ren L, Yang S, Xiao M, Chang D, Yang F, et al. Profiling Early Humoral Response to Diagnose Novel Coronavirus Disease (COVID-19). *Clin Infect Dis.* 2020;71(15):778-85. doi: 10.1093/cid/ciaa310.
46. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus associated with human respiratory disease in China. *Nature.* 2020;579(7798):265-9. doi: 10.1038/s41586-020-2008-3.
47. Hu D, Zhu C, Ai L, He T, Wang Y, Ye F, et al. Genomic characterization and infectivity of a novel SARS-like coronavirus in Chinese bats. *Emerg Microbes Infect.* 2018;7(1):154. doi: 10.1038/s41426-018-0155-5.
48. McIntosh K, Becker WB, Chanock RM. Growth in suckling-mouse brain of "IBV-like" viruses from patients with upper respiratory tract disease. *Proc Natl Acad Sci U S A.* 1967;58(6):2268-73. doi: 10.1073/pnas.58.6.2268.
49. Gámbaro F, Behillil S, Baidaliuk A, Donati F, Albert M, Alexandru A, et al. Introductions and early spread of SARS-CoV-2 in France, 24 January to 23 March 2020. *Euro Surveill.* 2020;25(26):2001200. doi: 10.2807/1560-7917.ES.2020.25.26.2001200.
50. Deslandes A, Berti V, Tandjaoui-Lambotte Y, Alloui C, Carbonnelle E, Zahar JR, et al. SARS-CoV-2 was already spreading in France in late December 2019. *Int J Antimicrob Agents.* 2020;55(6):106006. doi: 10.1016/j.ijantimicag.2020.106006.
51. GISAIID. Available from: <https://www.gisaid.org/> (accessed May 2020).
52. Lauring AS, Andino R. Quasispecies theory and the behavior of RNA viruses. *PLoS Pathog.* 2010;6(7):e1001005. doi: 10.1371/journal.ppat.1001005.
53. Domingo E, Baranowski E, Ruiz-Jarabo CM, Martín-Hernández AM, Sáiz JC, Escarmis C. Quasispecies structure and persistence of RNA viruses. *Emerg Infect Dis.* 1998;4(4):521-7. doi: 10.3201/eid0404.980402.
54. Domingo E, Martin V, Perales C, Grande-Pérez A, García-Arriaza J, Arias A. Viruses as quasispecies: biological implications. *Curr Top Microbiol Immunol.* 2006;299:51-82. doi: 10.1007/3-540-26397-7_3.
55. Duffy S, Shackleton LA, Holmes EC. Rates of evolutionary change in viruses: patterns and determinants. *Nat Rev Genet.* 2008;9(4):267-76. doi: 10.1038/nrg2323.
56. Holland J, Spindler K, Horodyski F, Grabau E, Nichol S, VandePol S. Rapid evolution of RNA genomes. *Science.* 1982;215(4540):1577-85. doi: 10.1126/science.7041255.
57. Batschelet E, Domingo E, Weissmann C. The proportion of revertant and mutant phage in a growing population, as a function of mutation and growth rate. *Gene.* 1976;1(1):27-32. doi: 10.1016/0378-1119(76)90004-4.
58. Steinhauer DA, Holland JJ. Rapid evolution of RNA viruses. *Annu Rev Microbiol.* 1987;41:409-33. doi: 10.1146/annurev.mi.41.100187.002205.
59. Shen Z, Xiao Y, Kang L, Ma W, Shi L, Zhang L, et al. Genomic Diversity of Severe Acute Respiratory Syndrome-Coronavirus 2 in Patients With Coronavirus Disease 2019. *Clin Infect Dis.* 2020;71(15):713-20. doi: 10.1093/cid/ciaa203.
60. Goldmann DA. Transmission of viral respiratory infections in the home. *Pediatr Infect Dis J.* 2000;19(10 Suppl):S97-102. doi: 10.1097/00006454-200010001-00002.
61. Phan T. Genetic diversity and evolution of SARS-CoV-2. *Infect Genet Evol.* 2020;81:104260. doi: 10.1016/j.meegid.2020.104260.
62. Ni M, Chen C, Qian J, Xiao HX, Shi WF, Luo Y, et al. Intra-host dynamics of Ebola virus during 2014. *Nat Microbiol.* 2016;1(11):16151. doi: 10.1038/nmicrobiol.2016.151.
63. Zhao Z, Li H, Wu X, Zhong Y, Zhang K, Zhang YP, et al. Moderate mutation rate in the SARS coronavirus genome and its implications. *BMC Evol Biol.* 2004;4:21. doi: 10.1186/1471-2148-4-21.
64. Marra MA, Jones SJ, Astell CR, Holt RA, Brooks-Wilson A, Butterfield YS, et al. The Genome sequence of the SARS-associated coronavirus. *Science.* 2003;303(5624):1399-404. doi: 10.1126/science.1085953.
65. Braun MJ, Clements JE, Gonda MA. The visna virus genome: evidence for a hypervariable site in the env gene and sequence homology among lentivirus envelope proteins. *J Virol.* 1987;61(12):4046-54. doi: 10.1128/JVI.61.12.4046-4054.1987.
66. Chinese SARS Molecular Epidemiology Consortium. Molecular evolution of the SARS coronavirus during the course of the SARS epidemic in China. *Science.* 2004;303(5664):1666-9. doi: 10.1126/science.1092002.
67. Brown EG, Tetro JA. Comparative analysis of the SARS coronavirus genome: a good start to a long journey. *Lancet.* 2003;361(9371):1756-7. doi: 10.1016/S0140-6736(03)13444-7.
68. Phan LT, Nguyen TV, Luong QC, Nguyen TV, Nguyen HT, Le HQ, et al. Importation and Human-to-Human Transmission of a Novel Coronavirus in Vietnam. *N Engl J Med.* 2020;382(9):872-4. doi: 10.1056/NEJMc2001272.
69. Zumla A, Hui DS. Infection control and MERS-CoV in health-care workers. *Lancet.* 2014;383(9932):1869-71. doi: 10.1016/S0140-6736(14)60852-7.
70. WHO. WHO statement on the Fifth Meeting of the IHR Emergency Committee concerning MERS-CoV. May 14, 2014. Available from: <https://www.who.int/mediacentre/news/statements/2014/mers-20140514/en/> (accessed May 2020).
71. Sizon J, Yu MW, Talbot PJ. Survival of human coronaviruses 229E and OC43 in suspension and after drying on surfaces: a possible source of hospital-acquired infections. *J Hosp Infect.* 2000;46(1):55-60. doi: 10.1053/jhin.2000.0795.
72. Falsey AR, McCann RM, Hall WJ, Criddle MM, Formica MA, Wycoff D, et al. The "common cold" in frail older persons: impact of rhinovirus and coronavirus in a senior daycare center. *J Am Geriatr Soc.* 1997;45(6):706-11. doi: 10.1111/j.1532-5415.1997.tb01474.x.
73. Seto WH, Conly JM, Pessoa-Silva CL, Malik M, Eremin S. Infection prevention and control measures for acute respiratory infections in healthcare settings: an update. *East Mediterr Health J.* 2013;19 Suppl 1:S39-47.
74. Health Protection Agency (HPA) UK Novel Coronavirus Investigation team. Evidence of person-to-person transmission within a family cluster of novel coronavirus infections. United Kingdom, February 2013. *Euro Surveill.* 2013;18(11):20427. doi: 10.2807/ese.18.11.20427-en.
75. Pebody RG, Chand MA, Thomas HL, Green HK, Boddington NL, Carvalho C, et al. The United Kingdom public health response to an imported laboratory confirmed case of a novel coronavirus in September 2012. *Euro Surveill.* 2012;17(40):20292.
76. Buchholz U, Müller MA, Nitsche A, Sanewski A, Wevering N, Bauer-Balci T, et al. Contact investigation of a case of human novel coronavirus infection treated in a German hospital, October-November 2012. *Euro Surveill.* 2013;18(8):20406.
77. Bean B, Moore BM, Sterner B, Peterson LR, Gerding DN, Balfour HH Jr. Survival of influenza viruses on environmental surfaces. *J Infect Dis.* 1982;146(1):47-51. doi: 10.1093/infdis/146.1.47.
78. Ryan MA, Christian RS, Wohlrahe J. Handwashing and respiratory illness among young adults in military training. *Am J Prev Med.* 2001;21(2):79-83. doi: 10.1016/s0749-3797(01)00323-3.
79. van Dorp L, Richard D, Tan CCS, Shaw LP, Acman M, Balloux F. No evidence for increased transmissibility from recurrent mutations in SARS-CoV-2. *Nat Commun.* 2020;11(1):5986. doi: 10.1038/s41467-020-19818-2.
80. Buxton Bridges C, Katz JM, Seto WH, Chan PK, Tsang D, Ho W, et al. Risk of influenza A (H5N1) infection among health care workers exposed to patients with influenza A (H5N1), Hong Kong. *J Infect Dis.* 2000;181(1):344-8. doi: 10.1086/315213.
81. Evans ME, Hall KL, Berry SE. Influenza control in acute care hospitals. *Am J Infect Control.* 1997;25(4):357-62. doi: 10.1016/s0196-6553(97)90029-8.
82. Cunney RJ, Bialachowski A, Thornley D, Smaill FM, Pennie RA. An outbreak of influenza A in a neonatal intensive care unit. *Infect Control Hosp Epidemiol.* 2000;21(7):449-54. doi: 10.1086/501786.
83. Sugaya N, Kusumoto N, Suzuki Y, Nerome R, Nerome K. Large sequential outbreaks caused by influenza A (H3N2) and B viruses in an

- institution for the mentally handicapped. *J Med Virol.* 1996;50(2):120-5. doi: 10.1002/(SICI)1096-9071(199610)50:2<120::AID-JMV4>3.0.CO;2-C.
84. Bean B, Rhame FS, Hughes RS, Weiler MD, Peterson LR, Gerding DN. Influenza B: hospital activity during a community epidemic. *Diagn Microbiol Infect Dis.* 1983;1(3):177-83. doi: 10.1016/0732-8893(83)90016-0.
85. Paules CI, Marston HD, Fauci AS. Coronavirus Infections-More Than Just the Common Cold. *JAMA.* 2020;323(8):707-8. doi: 10.1001/jama.2020.0757.
86. Hall CB. The spread of influenza and other respiratory viruses: complexities and conjectures. *Clin Infect Dis.* 2007;45(3):353-9. doi: 10.1086/519433.
87. Langmuir AD. Changing concepts of airborne infection of acute contagious diseases: a reconsideration of classic epidemiologic theories. *Ann N Y Acad Sci.* 1980;353:35-44. doi: 10.1111/j.1749-6632.1980.tb18903.x.
88. van den Brand JM, Smits SL, Haagmans BL. Pathogenesis of Middle East respiratory syndrome coronavirus. *J Pathol.* 2015;235(2):175-84. doi: 10.1002/path.4458.
89. Burke JP. Infection control - a problem for patient safety. *N Engl J Med.* 2003;348(7):651-6. doi: 10.1056/NEJMhpr020557.
90. Li W, Zhang C, Sui J, Kuhn JH, Moore MJ, Luo S, et al. Receptor and viral determinants of SARS-coronavirus adaptation to human ACE2. *EMBO J.* 2005;24(8):1634-43. doi: 10.1038/sj.emboj.7600640.
91. Qu XX, Hao P, Song XJ, Jiang SM, Liu YX, Wang PG, et al. Identification of two critical amino acid residues of the severe acute respiratory syndrome coronavirus spike protein for its variation in zoonotic tropism transition via a double substitution strategy. *J Biol Chem.* 2005;280(33):29588-95. doi: 10.1074/jbc.M500662200.
92. Bell DM; World Health Organization Working Group on International and Community Transmission of SARS. Public health interventions and SARS spread, 2003. *Emerg Infect Dis.* 2004;10(11):1900-6. doi: 10.3201/eid1011.040729.
93. Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease outbreak controllable. *Proc Natl Acad Sci U S A.* 2004;101(16):6146-51. doi: 10.1073/pnas.0307506101.
94. Mumford JD. Environmental risk evaluation in quarantine decision making. In: Anderson K, McRae C, Wilson D. *The economics of quarantine and the SPS agreement.* University of Adelaide Press; 2001:353-84.
95. Breukers A, Mourits M, van der Werf W, Lansink AO. Costs and benefits of controlling quarantine diseases: a bio-economic modeling approach. *Agric Econ.* 2008;38(2):137-49. doi: 10.1111/j.1574-0862.2008.00288.x.
96. Mumford JD. Economic issues related to quarantine in international trade. *Eur Rev Agric Econ.* 2002;29(3):329-48.
97. James S, Anderson K. On the need for more economic assessment of quarantine policies. *Aust J Agric Resour Econ.* 1998;42(4):425-44. doi: 10.22004/ag.econ.117286.
98. Leung GM, Chung PH, Tsang T, Lim W, Chan SK, Chau P, et al. SARS-CoV antibody prevalence in all Hong Kong patient contacts. *Emerg Infect Dis.* 2004;10(9):1653-6. doi: 10.3201/eid1009.040155.
99. Riley S, Fraser C, Donnelly CA, Ghani AC, Abu-Raddad LJ, Hedley AJ, et al. Transmission dynamics of the etiological agent of SARS in Hong Kong: impact of public health interventions. *Science.* 2003;300(5627):1961-6. doi: 10.1126/science.1086478.
100. Guan Y, Zheng BJ, He YQ, Liu XL, Zhuang ZX, Cheung CL, et al. Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science.* 2003;302(5643):276-8. doi: 10.1126/science.1087139.
101. Centers for Disease Control and Prevention (CDC). Prevalence of IgG antibody to SARS-associated coronavirus in animal traders--Guangdong Province, China, 2003. *MMWR Morb Mortal Wkly Rep.* 2003;52(41):986-7.
102. Yu IT, Li Y, Wong TW, Tam W, Chan AT, Lee JH, et al. Evidence of airborne transmission of the severe acute respiratory syndrome virus. *N Engl J Med.* 2004;350(17):1731-9. doi: 10.1056/NEJMoa032867.
103. de Wit E, Rasmussen AL, Falzarano D, Bushmaker T, Feldmann F, Brining DL, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) causes transient lower respiratory tract infection in rhesus macaques. *Proc Natl Acad Sci U S A.* 2013;110(41):16598-603. doi: 10.1073/pnas.1310744110.
104. Gralinski LE, Baric RS. Molecular pathology of emerging coronavirus infections. *J Pathol.* 2015;235(2):185-95. doi: 10.1002/path.4454.
105. Coleman CM, Frieman MB. Coronaviruses: important emerging human pathogens. *J Virol.* 2014;88(10):5209-12. doi: 10.1128/JVI.03488-13.
106. Woo PC, Lau SK, Lam CS, Lau CC, Tsang AK, Lau JH, et al. Discovery of seven novel Mammalian and avian coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of gammacoronavirus and deltacoronavirus. *J Virol.* 2012;86(7):3995-4008. doi: 10.1128/JVI.06540-11.
107. Lau SK, Woo PC, Yip CC, Fan RY, Huang Y, Wang M, et al. Isolation and characterization of a novel Betacoronavirus subgroup A coronavirus, rabbit coronavirus HKU14, from domestic rabbits. *J Virol.* 2012;86(10):5481-96. doi: 10.1128/JVI.06927-11.
108. Lau SK, Poon RW, Wong BH, Wang M, Huang Y, Xu H, et al. Coexistence of different genotypes in the same bat and serological characterization of Roussetus bat coronavirus HKU9 belonging to a novel Betacoronavirus subgroup. *J Virol.* 2010;84(21):11385-94. doi: 10.1128/JVI.01121-10.
109. Lau SK, Woo PC, Li KS, Huang Y, Wang M, Lam CS, et al. Complete genome sequence of bat coronavirus HKU2 from Chinese horseshoe bats revealed a much smaller spike gene with a different evolutionary lineage from the rest of the genome. *Virology.* 2007;367(2):428-39. doi: 10.1016/j.virol.2007.06.009.
110. Benvenuto D, Giovanetti M, Ciccozzi A, Spoto S, Angeletti S, Ciccozzi M. The 2019-new coronavirus epidemic: Evidence for virus evolution. *J Med Virol.* 2020;92(4):455-9. doi: 10.1002/jmv.25688.
111. Kim YI, Kim SG, Kim SM, Kim EH, Park SJ, Yu KM, et al. Infection and Rapid Transmission of SARS-CoV-2 in Ferrets. *Cell Host Microbe.* 2020;27(5):704-9.e2. doi: 10.1016/j.chom.2020.03.023.
112. Richard M, Kok A, de Meulder D, Bestebroer TM, Lamers MM, Okba NMA, et al. SARS-CoV-2 is transmitted via contact and via the air between ferrets. *Nat Commun.* 2020;11(1):3496. doi: 10.1038/s41467-020-17367-2.
113. Munster VJ, Feldmann F, Williamson BN, van Doremalen N, Pérez-Pérez L, Schulz J, et al. Respiratory disease and virus shedding in rhesus macaques inoculated with SARS-CoV-2. *bioRxiv [Preprint].* 2020:2020.03.21.001628. doi: 10.1101/2020.03.21.001628.
114. World Health Organization. Modes of transmission of virus causing COVID-19: implications for IPC precaution recommendations: scientific brief, 29 March 2020. Available from: [https://apps.who.int/iris/handle/10665/331616?locale-attribute=es&\(accessed May 2020\).](https://apps.who.int/iris/handle/10665/331616?locale-attribute=es&(accessed May 2020).)
115. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. *JAMA.* 2020;323(16):1610-2. doi: 10.1001/jama.2020.3227.
116. Hasonry HJ, Macnaughton MR. Prevalence of human coronavirus antibody in the population of southern Iraq. *J Med Virol.* 1982;9(3):209-16. doi: 10.1002/jmv.1890090308.
117. Buchholz UJ, Bukreyev A, Yang L, Lamirande EW, Murphy BR, Subbarao K, et al. Contributions of the structural proteins of severe acute respiratory syndrome coronavirus to protective immunity. *Proc Natl Acad Sci U S A.* 2004;101(26):9804-9. doi: 10.1073/pnas.0403492101.
118. ter Meulen J, van den Brink EN, Poon LL, Marissen WE, Leung CS, Cox F, et al. Human monoclonal antibody combination against SARS coronavirus: synergy and coverage of escape mutants. *PLoS Med.* 2006;3(7):e237. doi: 10.1371/journal.pmed.0030237.
119. Hsueh PR, Kao CL, Lee CN, Chen LK, Ho MS, Sia C, et al. SARS antibody test for serosurveillance. *Emerg Infect Dis.* 2004;10(9):1558-62. doi: 10.3201/eid1009.040101.
120. Peiris JS, Chu CM, Cheng VC, Chan KS, Hung IF, Poon LL, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet.* 2003;361(9371):1767-72. doi: 10.1016/s0140-6736(03)13412-5.

121. Grant PR, Garson JA, Tedder RS, Chan PK, Tam JS, Sung JJ. Detection of SARS coronavirus in plasma by real-time RT-PCR. *N Engl J Med*. 2003;349(25):2468-9. doi: 10.1056/NEJM200312183492522.
122. Wu HS, Chiu SC, Tseng TC, Lin SF, Lin JH, Hsu YH, et al. Serologic and molecular biologic methods for SARS-associated coronavirus infection, Taiwan. *Emerg Infect Dis*. 2004;10(2):304-10. doi: 10.3201/eid1002.030731.
123. World Health Organization. Communicable disease surveillance & response. Use of laboratory methods for SARS diagnosis. Available from: <https://www.who.int/health-topics/severe-acute-respiratory-syndrome/technical-guidance/laboratory/use-of-laboratory-methods-for-sars-diagnosis>(accessed May 2020).
124. Yam WC, Chan KH, Poon LL, Guan Y, Yuen KY, Seto WH, et al. Evaluation of reverse transcription-PCR assays for rapid diagnosis of severe acute respiratory syndrome associated with a novel coronavirus. *J Clin Microbiol*. 2003;41(10):4521-4. doi: 10.1128/jcm.41.10.4521-4524.2003.
125. Centers for Disease Control and Prevention (CDC). Revised U.S. surveillance case definition for severe acute respiratory syndrome (SARS) and update on SARS cases--United States and worldwide, December 2003. *MMWR Morb Mortal Wkly Rep*. 2003;52(49):1202-6.
126. Peiris JS, Yuen KY, Osterhaus AD, Stöhr K. The severe acute respiratory syndrome. *N Engl J Med*. 2003;349(25):2431-41. doi: 10.1056/NEJMra032498.
127. Li G, Chen X, Xu A. Profile of specific antibodies to the SARS-associated coronavirus. *N Engl J Med*. 2003;349(5):508-9. doi: 10.1056/NEJM200307313490520.
128. Woo PC, Lau SK, Wong BH, Chan KH, Chu CM, Tsoi HW, et al. Longitudinal profile of immunoglobulin G (IgG), IgM, and IgA antibodies against the severe acute respiratory syndrome (SARS) coronavirus nucleocapsid protein in patients with pneumonia due to the SARS coronavirus. *Clin Diagn Lab Immunol*. 2004;11(4):665-8. doi: 10.1128/CDLI.11.4.665-668.2004.
129. Shi Y, Wan Z, Li L, Li P, Li C, Ma Q, et al. Antibody responses against SARS-coronavirus and its nucleocapsid in SARS patients. *J Clin Virol*. 2004;31(1):66-8. doi: 10.1016/j.jcv.2004.05.006.
130. Sanna PP, Burton DR. Role of antibodies in controlling viral disease: lessons from experiments of nature and gene knockouts. *J Virol*. 2000;74(21):9813-7. doi: 10.1128/jvi.74.21.9813-9817.2000.
131. Amanat F, Stadlbauer D, Strohmaier S, Nguyen THO, Chromikova V, McMahon M, et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat Med*. 2020;26(7):1033-6. doi: 10.1038/s41591-020-0913-5.
132. Liu W, Fontanet A, Zhang PH, Zhan L, Xin ZT, Baril L, et al. Two-year prospective study of the humoral immune response of patients with severe acute respiratory syndrome. *J Infect Dis*. 2006;193(6):792-5. doi: 10.1086/500469.
133. Callow KA, Parry HF, Sergeant M, Tyrrell DA. The time course of the immune response to experimental coronavirus infection of man. *Epidemiol Infect*. 1990;105(2):435-46. doi: 10.1017/s0950268800048019.
134. Choe PG, Perera RAPM, Park WB, Song KH, Bang JH, Kim ES, et al. MERS-CoV Antibody Responses 1 Year after Symptom Onset, South Korea, 2015. *Emerg Infect Dis*. 2017;23(7):1079-1084. doi: 10.3201/eid2307.170310.
135. Payne DC, Iblan I, Rha B, Alqasrawi S, Haddadin A, Al Nsour M, et al. Persistence of Antibodies against Middle East Respiratory Syndrome Coronavirus. *Emerg Infect Dis*. 2016;22(10):1824-6. doi: 10.3201/eid2210.160706.
136. Haveri A, Smura T, Kuivanen S, Österlund P, Hepojoki J, Ikonen N, et al. Serological and molecular findings during SARS-CoV-2 infection: the first case study in Finland, January to February 2020. *Euro Surveill*. 2020;25(11):2000266. doi: 10.2807/1560-7917.ES.2020.25.11.2000266.
137. Cowling BJ, Chan KH, Fang VJ, Lau LLH, So HC, Fung ROP, et al. Comparative epidemiology of pandemic and seasonal influenza A in households. *N Engl J Med*. 2010;362(23):2175-84. doi: 10.1056/NEJMoa0911530.
138. Liu L, Xie J, Sun J, Han Y, Zhang C, Fan H, et al. Longitudinal profiles of immunoglobulin G antibodies against severe acute respiratory syndrome coronavirus components and neutralizing activities in recovered patients. *Scand J Infect Dis*. 2011;43(6-7):515-21. doi: 10.3109/00365548.2011.560184.
139. Zhang W, Du RH, Li B, Zheng XS, Yang XL, Hu B, et al. Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. *Emerg Microbes Infect*. 2020;9(1):386-389. doi: 10.1080/22221751.2020.1729071.
140. Amanat F, Meade P, Strohmaier S, Krammer F. Cross-reactive antibodies binding to H4 hemagglutinin protect against a lethal H4N6 influenza virus challenge in the mouse model. *Emerg Microbes Infect*. 2019;8(1):155-168. doi: 10.1080/22221751.2018.1564369.
141. Wohlbold TJ, Podolsky KA, Chromikova V, Kirkpatrick E, Falconieri V, Meade P, et al. Broadly protective murine monoclonal antibodies against influenza B virus target highly conserved neuraminidase epitopes. *Nat Microbiol*. 2017;2(10):1415-1424. doi: 10.1038/s41564-017-0011-8.
142. Stadlbauer D, Amanat F, Chromikova V, Jiang K, Strohmaier S, Arunkumar GA, et al. SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a Serological Assay, Antigen Production, and Test Setup. *Curr Protoc Microbiol*. 2020;57(1):e100. doi: 10.1002/cpmc.100.
143. Huang LR, Chiu CM, Yeh SH, Huang WH, Hsueh PR, Yang WZ, et al. Evaluation of antibody responses against SARS coronavirus nucleocapsid or spike proteins by immunoblotting or ELISA. *J Med Virol*. 2004;73(3):338-46. doi: 10.1002/jmv.20096.
144. Kellam P, Barclay W. The dynamics of humoral immune responses following SARS-CoV-2 infection and the potential for reinfection. *J Gen Virol*. 2020;101(8):791-7. doi: 10.1099/jgv.0.001439.
145. Brekke OH, Sandlie I. Therapeutic antibodies for human diseases at the dawn of the twenty-first century. *Nat Rev Drug Discov*. 2003;2(1):52-62. doi: 10.1038/nrd984.
146. Wong VW, Dai D, Wu AK, Sung JJ. Treatment of severe acute respiratory syndrome with convalescent plasma. *Hong Kong Med J*. 2003;9(3):199-201.
147. Subbarao K, McAuliffe J, Vogel L, Fahle G, Fischer S, Tatti K, et al. Prior infection and passive transfer of neutralizing antibody prevent replication of severe acute respiratory syndrome coronavirus in the respiratory tract of mice. *J Virol*. 2004;78(7):3572-7. doi: 10.1128/jvi.78.7.3572-3577.2004.
148. Zhang Z, Xie YW, Hong J, Zhang X, Kwok SY, Huang X, et al. Purification of severe acute respiratory syndrome hyperimmune globulins for intravenous injection from convalescent plasma. *Transfusion*. 2005;45(7):1160-4. doi: 10.1111/j.1537-2995.2005.00179.x.
149. Sui J, Li W, Roberts A, Matthews LJ, Murakami A, Vogel L, et al. Evaluation of human monoclonal antibody 80R for immunoprophylaxis of severe acute respiratory syndrome by an animal study, epitope mapping, and analysis of spike variants. *J Virol*. 2005;79(10):5900-6. doi: 10.1128/JVI.79.10.5900-5906.2005.
150. Traggiai E, Becker S, Subbarao K, Kolesnikova L, Uematsu Y, Gismondo MR, et al. An efficient method to make human monoclonal antibodies from memory B cells: potent neutralization of SARS coronavirus. *Nat Med*. 2004;10(8):871-5. doi: 10.1038/nm1080.
151. ter Meulen J, Bakker AB, van den Brink EN, Weverling GJ, Martina BE, Haagmans BL, et al. Human monoclonal antibody as prophylaxis for SARS coronavirus infection in ferrets. *Lancet*. 2004;363(9427):2139-41. doi: 10.1016/S0140-6736(04)16506-9.
152. Zwick MB, Labrijn AF, Wang M, Spenlehauer C, Saphire EO, Binley JM, et al. Broadly neutralizing antibodies targeted to the membrane-proximal external region of human immunodeficiency virus type 1 glycoprotein gp41. *J Virol*. 2001;75(22):10892-905. doi: 10.1128/JVI.75.22.10892-10905.2001.
153. Xu D, Zhang Z, Wang FS. SARS-associated coronavirus quasispecies in individual patients. *N Engl J Med*. 2004;350(13):1366-7. doi: 10.1056/NEJMc032421.
154. Liu J, Lim SL, Ruan Y, Ling AE, Ng LF, Drosten C, et al. SARS transmission pattern in Singapore reassessed by viral sequence variation analysis. *PLoS Med*. 2005;2(2):e43. doi: 10.1371/journal.pmed.0020043.
155. Poon LL, Leung CS, Chan KH, Yuen KY, Guan Y, Peiris JS. Recurrent mutations associated with isolation and passage of SARS coronavirus in cells from non-human primates. *J Med Virol*. 2005;76(4):435-40. doi: 10.1002/jmv.20379.
156. Liu L, Wei Q, Lin Q, Fang J, Wang H, Kwok H, et al. Anti-spike IgG causes severe acute lung injury by skewing macrophage responses during acute SARS-CoV infection. *JCI Insight*. 2019;4(4):e123158. doi: 10.1172/jci.insight.123158.

157. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science*. 2020;368(6493):860-8. doi: 10.1126/science.abb5793.
158. Eickhoff TC, Sherman IL, Serfling RE. Observations on excess mortality associated with epidemic influenza. *JAMA*. 1961;176:776-82. doi: 10.1001/jama.1961.03040220024005.
159. Blumenfeld HL, Kilbourne ED, LOURIA DB, Rogers DE. Studies on influenza in the pandemic of 1957-1958. I. An epidemiologic, clinical and serologic investigation of an intrahospital epidemic, with a note on vaccination efficacy. *J Clin Invest*. 1959;38(1 Part 2):199-212. doi: 10.1172/JCI103789.
160. Martin WJ. Recent changes in the death rate from influenza. *Br Med J*. 1950;1(4648):267. doi: 10.1136/bmj.1.4648.267.
161. Rosenwald MS. History's deadliest pandemics, from ancient Rome to modern America. Available from: <https://www.spokesman.com/stories/2020/apr/15/historys-deadliest-pandemics-from-ancient-rome-to/> (accessed May 2020).
162. Harper SA, Bradley JS, Englund JA, File TM, Gravenstein S, Hayden FG, et al. Seasonal influenza in adults and children--diagnosis, treatment, chemoprophylaxis, and institutional outbreak management: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis*. 2009;48(8):1003-32. doi: 10.1086/598513.
163. Iuliano AD, Roguski KM, Chang HH, Muscatello DJ, Palekar R, Tempia S, et al. Estimates of global seasonal influenza-associated respiratory mortality: a modelling study. *Lancet*. 2018;391(10127):1285-300. doi: 10.1016/S0140-6736(17)33293-2.
164. GBD 2017 Influenza Collaborators. Mortality, morbidity, and hospitalisations due to influenza lower respiratory tract infections, 2017: an analysis for the Global Burden of Disease Study 2017. *Lancet Respir Med*. 2019;7(1):69-89. doi: 10.1016/S2213-2600(18)30496-X.
165. Simonsen L, Spreeuwenberg P, Lustig R, Taylor RJ, Fleming DM, Kroneman M, et al. Global mortality estimates for the 2009 Influenza Pandemic from the GLAMOR project: a modeling study. *PLoS Med*. 2013;7(11):e1001558. doi: 10.1371/journal.pmed.1001558.
166. Thompson WW, Shay DK, Weintraub E, Brammer L, Cox N, Anderson LJ, et al. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA*. 2003;289(2):179-86. doi: 10.1001/jama.289.2.179.
167. Hayward AC, Harling R, Wetten S, Johnson AM, Munro S, Smedley J, et al. Effectiveness of an influenza vaccine programme for care home staff to prevent death, morbidity, and health service use among residents: cluster randomised controlled trial. *BMJ*. 2006;333(7581):1241. doi: 10.1136/bmj.39010.581354.55.
168. Longini IM Jr, Koopman JS, Monto AS, Fox JP. Estimating household and community transmission parameters for influenza. *Am J Epidemiol*. 1982;115(5):736-51. doi: 10.1093/oxfordjournals.aje.a113356.
169. Gani R, Hughes H, Fleming D, Griffin T, Medlock J, Leach S. Potential impact of antiviral drug use during influenza pandemic. *Emerg Infect Dis*. 2005;11(9):1355-62. doi: 10.3201/eid1209.041344.
170. Ferguson NM, Cummings DA, Fraser C, Cajka JC, Cooley PC, Burke DS. Strategies for mitigating an influenza pandemic. *Nature*. 2006;442(7101):448-52. doi: 10.1038/nature04795.
171. Ferguson NM, Fraser C, Donnelly CA, Ghani AC, Anderson RM. Public health. Public health risk from the avian H5N1 influenza epidemic. *Science*. 2004;304(5673):968-9. doi: 10.1126/science.1096898.
172. Leung NH, Xu C, Ip DK, Cowling BJ. Review Article: The Fraction of Influenza Virus Infections That Are Asymptomatic: A Systematic Review and Meta-analysis. *Epidemiology*. 2015;26(6):862-72. doi: 10.1097/EDE.0000000000000340.
173. Saad-Roy CM, Wingreen NS, Levin SA, Grenfell BT. Dynamics in a simple evolutionary-epidemiological model for the evolution of an initial asymptomatic infection stage. *Proc Natl Acad Sci U S A*. 2020;117(21):11541-50. doi: 10.1073/pnas.1920761117.
174. Argente DO, Hsieh CT, Lee M. The Cost of Privacy: Welfare Effects of the Disclosure of Covid-19 Cases. *NBER Work Pap Ser*. 2020. doi: 10.3386/w27220.
175. Ing AJ, Cocks C, Green JP. COVID-19: in the footsteps of Ernest Shackleton. *Thorax*. 2020;75(8):693-4. doi: 10.1136/thoraxjnl-2020-215091.
176. Zhou G, Zhao Q. Perspectives on therapeutic neutralizing antibodies against the Novel Coronavirus SARS-CoV-2. *Int J Biol Sci*. 2020;16(10):1718-23. doi: 10.7150/ijbs.45123.
177. Duan K, Liu B, Li C, Zhang H, Yu T, Qu J, et al. Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proc Natl Acad Sci U S A*. 2020;117(17):9490-6. doi: 10.1073/pnas.2004168117.
178. World Health Organization Writing Group, Bell D, Nicoll A, Fukuda K, Horby P, Monto A, et al. Non-pharmaceutical interventions for pandemic influenza, international measures. *Emerg Infect Dis*. 2006;12(1):81-7. doi: 10.3201/eid1201.051370.
179. Hayward AC, Frangou EB, Birmingham A, Wang L, Copas A, Edmunds WJ, et al. Comparative community burden and severity of seasonal and pandemic influenza: results of the Flu Watch cohort study. *Lancet Respir Med*. 2014;2(6):445-54. doi: 10.1016/S2213-2600(14)70034-7.
180. Dijkman R, Jebbink MF, El Idrissi NB, Pijrc K, Müller MA, Kuijpers TW, et al. Human coronavirus NL63 and 229E seroconversion in children. *J Clin Microbiol*. 2008;46(7):2368-73. doi: 10.1128/JCM.00533-08.
181. Piedra PA, Gaglani MJ, Kozinetz CA, Herschler G, Riggs M, Griffith M, et al. Herd immunity in adults against influenza-related illnesses with use of the trivalent-live attenuated influenza vaccine (CAIV-T) in children. *Vaccine*. 2005;23(13):1540-8. doi: 10.1016/j.vaccine.2004.09.025.
182. Webster RG. Immunity to influenza in the elderly. *Vaccine*. 2000;18(16):1686-9. doi: 10.1016/S0264-410X(99)00507-1.
183. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet*. 2020;395(10224):565-74. doi: 10.1016/S0140-6736(20)30251-8.
184. Su S, Wong G, Shi W, Liu J, Lai ACK, Zhou J, et al. Epidemiology, Genetic Recombination, and Pathogenesis of Coronaviruses. *Trends Microbiol*. 2016;24(6):490-502. doi: 10.1016/j.tim.2016.03.003.
185. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med*. 2020;382(19):1787-99. doi: 10.1056/NEJMoa2001282.
186. Pelegrin M, Naranjo-Gomez M, Piechaczyk M. Antiviral Monoclonal Antibodies: Can They Be More Than Simple Neutralizing Agents? *Trends Microbiol*. 2015;23(10):653-65. doi: 10.1016/j.tim.2015.07.005.
187. Schmaljohn AL, McClain D. Alphaviruses (Togaviridae) and Flaviviruses (Flaviviridae). In: Baron S, editor. *Medical Microbiology*. 4th ed. Galveston (TX): University of Texas Medical Branch at Galveston; 1996:Chapter 54.
188. Brenner BG, Gryllis C, Wainberg MA. Role of antibody-dependent cellular cytotoxicity and lymphokine-activated killer cells in AIDS and related diseases. *J Leukoc Biol*. 1991;50(6):628-40. doi: 10.1002/jlb.50.6.628.
189. Bernasconi NL, Traggiai E, Lanzavecchia A. Maintenance of serological memory by polyclonal activation of human memory B cells. *Science*. 2002;298(5601):2199-202. doi: 10.1126/science.1076071.
190. Kurosaki T, Kometani K, Ise W. Memory B cells. *Nat Rev Immunol*. 2015;15(3):149-59. doi: 10.1038/nri3802.
191. Diamond MS, Shrestha B, Marri A, Mahan D, Engle M. B cells and antibody play critical roles in the immediate defense of disseminated infection by West Nile encephalitis virus. *J Virol*. 2003;77(4):2578-86. doi: 10.1128/jvi.77.4.2578-2586.2003.
192. Yang Y, Yang M, Yuan J, Wang F, Wang Z, Li J, et al. Laboratory Diagnosis and Monitoring the Viral Shedding of SARS-CoV-2 Infection. *Innovation*. 2020;1(3):100061. Doi: 10.1016/j.xinn.2020.100061.
193. Ludert JE, Alcalá AC, Liprandi F. Primer pair p289-p290, designed to detect both noroviruses and sapoviruses by reverse transcription-PCR, also detects rotaviruses by cross-reactivity. *J Clin Microbiol*. 2004;42(2):835-6. doi: 10.1128/jcm.42.2.835-836.2004.
194. Amar RL, Estes MK. Diagnosis of noncultivable gastroenteritis viruses, the human caliciviruses. *Clin Microbiol Rev*. 2001;14(1):15-37. doi: 10.1128/CMR.14.1.15-37.2001.
195. Vinjé J, Vennema H, Maunula L, von Bonsdorff CH, Hoehne M, Schreier E, et al. International collaborative study to compare reverse transcriptase PCR assays for detection and genotyping of noroviruses. *J Clin Microbiol*. 2003;41(4):1423-33. doi: 10.1128/jcm.41.4.1423-1433.2003.

196. Vijgen L, Keyaerts E, Moës E, Maes P, Duson G, Van Ranst M. Development of one-step, real-time, quantitative reverse transcriptase PCR assays for absolute quantitation of human coronaviruses OC43 and 229E. *J Clin Microbiol.* 2005;43(11):5452-6. doi: 10.1128/JCM.43.11.5452-5456.2005.
197. Drake JW, Holland JJ. Mutation rates among RNA viruses. *Proc Natl Acad Sci U S A.* 1999;96(24):13910-3. doi: 10.1073/pnas.96.24.13910.
198. Padmanabhan R, Mishra AK, Raoult D, Fournier PE. Genomics and metagenomics in medical microbiology. *J Microbiol Methods.* 2013;95(3):415-24. doi: 10.1016/j.mimet.2013.10.006.
199. Miller RR, Montoya V, Gardy JL, Patrick DM, Tang P. Metagenomics for pathogen detection in public health. *Genome Med.* 2013;5(9):81. doi: 10.1186/gm485.
200. Liais E, Croville G, Mariette J, Delverdier M, Lucas MN, Klopp C, et al. Novel avian coronavirus and fulminating disease in guinea fowl, France. *Emerg Infect Dis.* 2014;20(1):105-8. doi: 10.3201/eid2001.130774.
201. Tahamtan A, Ardebili A. Real-time RT-PCR in COVID-19 detection: issues affecting the results. *Expert Rev Mol Diagn.* 2020;20(5):453-4. doi: 10.1080/14737159.2020.1757437.
202. Xi M, Wei Q, Qihua F, Ming G. Understanding the Influence Factors in Viral Nucleic Acid Test of 2019 novel Coronavirus (2019-nCoV). *Chin J Lab Med.* 2020;12:E002.
203. Wang Y, Kang H, Liu X, Tong Z. Combination of RT-qPCR testing and clinical features for diagnosis of COVID-19 facilitates management of SARS-CoV-2 outbreak. *J Med Virol.* 2020;92(6):538-9. doi: 10.1002/jmv.25721.
204. Rainer TH, Chan PK, Ip M, Lee N, Hui DS, Smit D, et al. The spectrum of severe acute respiratory syndrome-associated coronavirus infection. *Ann Intern Med.* 2004;140(8):614-9. doi: 10.7326/0003-4819-140-8-200404200-00008.
205. Reuters. Tanzania suspends laboratory head after president questions coronavirus tests. Available from: <https://www.reuters.com/article/us-health-coronavirus-tanzania/tanzania-suspends-laboratory-head-after-president-questions-coronavirus-tests-idUSKBN2G295> (accessed May 2020).
206. Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-Negative Rate of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-CoV-2 Tests by Time Since Exposure. *Ann Intern Med.* 2020;173(4):262-7. doi: 10.7326/M20-1495.
207. Gao X, Zhou H, Wu C, Xiao Y, Ren L, Paranhos-Baccalà G, et al. Antibody against nucleocapsid protein predicts susceptibility to human coronavirus infection. *J Infect.* 2015;71(5):599-602. doi: 10.1016/j.jinf.2015.07.002.
208. Singh K, Vasoo S, Stevens J, Schreckenberger P, Trenholme G. Pitfalls in diagnosis of pandemic (novel) A/H1N1 2009 influenza. *J Clin Microbiol.* 2010;48(4):1501-3. doi: 10.1128/JCM.02483-09.
209. Blyth CC, Iredell JR, Dwyer DE. Rapid-test sensitivity for novel swine-origin influenza A (H1N1) virus in humans. *N Engl J Med.* 2009;361(25):2493. doi: 10.1056/NEJMc0909049.
210. Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, et al. Correlation of Chest CT and RT-PCR Testing for Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology.* 2020;296(2):E32-E40. doi: 10.1148/radiol.2020200642.
211. Huang P, Liu T, Huang L, Liu H, Lei M, Xu W, et al. Use of Chest CT in Combination with Negative RT-PCR Assay for the 2019 Novel Coronavirus but High Clinical Suspicion. *Radiology.* 2020;295(1):22-3. doi: 10.1148/radiol.2020200330.
212. Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for Typical Coronavirus Disease 2019 (COVID-19) Pneumonia: Relationship to Negative RT-PCR Testing. *Radiology.* 2020;296(2):E41-E45. doi: 10.1148/radiol.2020200343.
213. Shi H, Han X, Zheng C. Evolution of CT Manifestations in a Patient Recovered from 2019 Novel Coronavirus (2019-nCoV) Pneumonia in Wuhan, China. *Radiology.* 2020;295(1):20. doi: 10.1148/radiol.2020200269.
214. Pan Y, Guan H, Zhou S, Wang Y, Li Q, Zhu T, et al. Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63 patients in Wuhan, China. *Eur Radiol.* 2020;30(6):3306-9. doi: 10.1007/s00330-020-06731-x.
215. Ho PL, Chau PH, Yip PS, Ooi GC, Khong PL, Ho JC, et al. A prediction rule for clinical diagnosis of severe acute respiratory syndrome. *Eur Respir J.* 2005;26(3):474-9. doi: 10.1183/09031936.05.1076704.
216. Roberts A, Paddock C, Vogel L, Butler E, Zaki S, Subbarao K. Aged BALB/c mice as a model for increased severity of severe acute respiratory syndrome in elderly humans. *J Virol.* 2005;79(9):5833-8. doi: 10.1128/JVI.79.9.5833-5838.2005.
217. Papenburg J, Baz M, Hamelin MÈ, Rhéaume C, Carbonneau J, Ouakki M, et al. Household transmission of the 2009 pandemic A/H1N1 influenza virus: elevated laboratory-confirmed secondary attack rates and evidence of asymptomatic infections. *Clin Infect Dis.* 2010;51(9):1033-41. doi: 10.1086/656582.
218. Miller E, Hoschler K, Hardelid P, Stanford E, Andrews N, Zambon M. Incidence of 2009 pandemic influenza A H1N1 infection in England: a cross-sectional serological study. *Lancet.* 2010;375(9720):1100-8. doi: 10.1016/S0140-6736(09)62126-7.
219. Zambon M, Hays J, Webster A, Newman R, Keene O. Diagnosis of influenza in the community: relationship of clinical diagnosis to confirmed virological, serologic, or molecular detection of influenza. *Arch Intern Med.* 2001;161(17):2116-22. doi: 10.1001/archinte.161.17.2116.
220. Call SA, Vollenweider MA, Hornung CA, Simel DL, McKinney WP. Does this patient have influenza? *JAMA.* 2005;293(8):987-97. doi: 10.1001/jama.293.8.987.
221. Hohdatsu T, Okada S, Koyama H. Characterization of monoclonal antibodies against feline infectious peritonitis virus type II and antigenic relationship between feline, porcine, and canine coronaviruses. *Arch Virol.* 1991;117(1-2):85-95. doi: 10.1007/BF01310494.
222. Motokawa K, Hohdatsu T, Aizawa C, Koyama H, Hashimoto H. Molecular cloning and sequence determination of the peplomer protein gene of feline infectious peritonitis virus type I. *Arch Virol.* 1995;140(3):469-80. doi: 10.1007/BF01718424.
223. Takano T, Ishihara Y, Matsuoka M, Yokota S, Matsuoka-Kobayashi Y, Doki T, et al. Use of recombinant nucleocapsid proteins for serological diagnosis of feline coronavirus infection by three immunochromatographic tests. *J Virol Methods.* 2014;196:1-6. doi: 10.1016/j.jviromet.2013.10.014.
224. Ferris MM, Stepp PC, Ranno KA, Mahmoud W, Ibbitson E, Jarvis J, et al. Evaluation of the Virus Counter® for rapid baculovirus quantitation. *J Virol Methods.* 2011;171(1):111-6. doi: 10.1016/j.jviromet.2010.10.010.
225. Driskell JD, Jones CA, Tompkins SM, Tripp RA. One-step assay for detecting influenza virus using dynamic light scattering and gold nanoparticles. *Analyst.* 2011;136(15):3083-90. doi: 10.1039/c1an15303j.
226. Mironov GG, Chechik AV, Ozer R, Bell JC, Berezovski MV. Viral quantitative capillary electrophoresis for counting intact viruses. *Anal Chem.* 2011;83(13):5431-5. doi: 10.1021/ac201006u.
227. Schwille P, Bieschke J, Oehlenschläger F. Kinetic investigations by fluorescence correlation spectroscopy: the analytical and diagnostic potential of diffusion studies. *Biophys Chem.* 1997;66(2-3):211-28. doi: 10.1016/s0301-4622(97)00061-6.
228. Almázan F, González JM, Pénzes Z, Izeta A, Calvo E, Plana-Durán J, et al. Engineering the largest RNA virus genome as an infectious bacterial artificial chromosome. *Proc Natl Acad Sci U S A.* 2000;97(10):5516-21. doi: 10.1073/pnas.97.10.5516.
229. Yount B, Curtis KM, Baric RS. Strategy for systematic assembly of large RNA and DNA genomes: transmissible gastroenteritis virus model. *J Virol.* 2000;74(22):10600-11. doi: 10.1128/jvi.74.22.10600-10611.2000.
230. Thiel V, Herold J, Schelle B, Siddell SG. Infectious RNA transcribed in vitro from a cDNA copy of the human coronavirus genome cloned in vaccinia virus. *J Gen Virol.* 2001;82(Pt 6):1273-81. doi: 10.1099/0022-1317-82-6-1273.
231. Casais R, Thiel V, Siddell SG, Cavanagh D, Britton P. Reverse genetics system for the avian coronavirus infectious bronchitis virus. *J Virol.* 2001;75(24):12359-69. doi: 10.1128/JVI.75.24.12359-12369.2001.
232. Kouadio IK, Aljunid S, Kamigaki T, Hammad K, Oshitani H. Infectious diseases following natural disasters: prevention and control measures. *Expert Rev Anti Infect Ther.* 2012;10(1):95-104. doi: 10.1586/eri.11.155.
233. Sehulster L, Chinn RY; CDC; HICPAC. Guidelines for environmental infection control in health-care facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *MMWR Recomm Rep.* 2003;52(RR-10):1-42.

234. Siegel JD, Rhinehart E, Jackson M, Chiarello L; Health Care Infection Control Practices Advisory Committee. 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Health Care Settings. *Am J Infect Control*. 2007;35(10 Suppl 2):S65-164. doi: 10.1016/j.ajic.2007.10.007.
235. Chretien JH, Esswein JG. How frequent is bacterial superinfection of the pharynx in infectious mononucleosis? Observations on incidence, recognition, and management with antibiotics. *Clin Pediatr (Phila)*. 1976;15(5):424-7. doi: 10.1177/000992287601500505.
236. Hansen NS, Byberg S, Hervig Jacobsen L, Bjerregaard-Andersen M, et al. Effect of early measles vaccine on pneumococcal colonization: A randomized trial from Guinea-Bissau. *PLoS One*. 2017;12(5):e0177547. doi: 10.1371/journal.pone.0177547.
237. Alter MJ. Epidemiology of viral hepatitis and HIV co-infection. *J Hepatol*. 2006;44(1 Suppl):S6-9. doi: 10.1016/j.jhep.2005.11.004.
238. Pawlowski A, Jansson M, Sköld M, Rottenberg ME, Källenius G. Tuberculosis and HIV co-infection. *PLoS Pathog*. 2012;8(2):e1002464. doi: 10.1371/journal.ppat.1002464.
239. Rockstroh JK, Spengler U. HIV and hepatitis C virus co-infection. *Lancet Infect Dis*. 2004;4(7):437-44. doi: 10.1016/S1473-3099(04)01059-X.
240. Wilson J, Loveday H. Does glove use increase the risk of infection?. *Nursing Times*. 2014;110(39):12-5.
241. Heal JS, Blom AW, Titcomb D, Taylor A, Bowker K, Hardy JR. Bacterial contamination of surgical gloves by water droplets spilt after scrubbing. *J Hosp Infect*. 2003;53(2):136-9. doi: 10.1053/jhin.2002.1352.
242. Misteli H, Weber WP, Reck S, Rosenthal R, Zwahlen M, Fueglistaler P, et al. Surgical glove perforation and the risk of surgical site infection. *Arch Surg*. 2009;144(6):553-8; discussion 558. doi: 10.1001/archsurg.2009.60.
243. Piro S, Samud M, Badi S, Al Ssabi L. Hospital-acquired malaria transmitted by contaminated gloves. *J Hosp Infect*. 2001;47(2):156-8. doi: 10.1053/jhin.2000.0907.
244. Ye D, Shan J, Huang Y, Li J, Li C, Liu X, et al. A gloves-associated outbreak of imipenem-resistant *Acinetobacter baumannii* in an intensive care unit in Guangdong, China. *BMC Infect Dis*. 2015;15:179. doi: 10.1186/s12879-015-0917-9.
245. Burke FJ. Use of non-sterile gloves in clinical practice. *J Dent*. 1990;18(2):79-89. doi: 10.1016/0300-5712(90)90089-w.
246. Go! & Express Live. EC COVID-19 testing. Available from: <https://www.goexpress.co.za/2020/04/27/the-eastern-capes-covid-19-breakdown-and-coronavirus-hotspots/ec-covid-19-testing> (accessed May 2020).
247. Daily Express News. Available from: https://www.express.com.pk/epaper/PoPupwindow.aspx?newsID=1107450831&Issue=NP_KHI&Date=20200521 (accessed May 2020).