

Assessment of knowledge and awareness of community pharmacists toward epilepsy

Muhammed Mansour¹, Aisha E. Alkadi^{1*}  , Heba A. Alnayif¹ and Souad Qtannish²

¹ Department of Pharmaceutics, Faculty of Pharmacy, University of Tripoli, Tripoli, Libya

² Ali Omer Asker Hospital, Ministry of Health, Tripoli, Libya

*Author to whom correspondence should be addressed

Received: 30-04-2022, **Revised:** 27-06-2022, **Accepted:** 25-07-2022, **Published:** 30-09-2022

Copyright © 2022. This open-access article is distributed under the *Creative Commons Attribution License*, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Mansour et al. (2022) Assessment of knowledge and awareness of community pharmacists toward epilepsy
Mediterr J Pharm Pharm Sci. 2 (3): 17-23. [Article number: 78]. <https://doi.org/10.5281/zenodo.7115139>

Keywords: Antiepileptic drug, community pharmacist, epilepsy, knowledge, Libya

Abstract: Community pharmacist plays an essential role in educating epileptic patients about their disease and medications. Improving the patient's awareness may lead to improved compliance and decreased drug-drug interaction and ultimately improved their quality of life. Pharmacists can detect the emergence of health problems and can help prevent the progression of comorbidities. Considering the complexity of treating epilepsy and the lack of information about pharmacists' contributions to epilepsy management, pharmacists performed pharmaceutical counselling, pharmaco-therapeutic follow-up and systematic measurement and evaluation of findings and increased medication adherence of patients with epilepsy. This study aimed to assess the community pharmacist's knowledge of epilepsy and their treatment by antiepileptic drugs in Libya. The design of the study is a cross-sectional study. The knowledge was collected through a questionnaire which included 35 questions divided into three sections: demographic data, general information about epilepsy and information about epilepsy and antiepileptic drugs. The questionnaire was prepared and validated by consultant training in Ali Omer Asker Hospital in Tripoli for two months. The findings indicate that over 200 patients are altered viewed, of whom, only epileptic patients were observed. The majority of the participants were female, qualification degree BSc and years of experience from one to five years. Unfortunately, some of the participants (40.0%) had poor knowledge and about 60.0% of them had good knowledge. This study indicates that the importance of community pharmacists in Libya requires more improvement to achieve the existing function and that the impact of continuous study of everything related to diseases and medicines is important to obtain a qualified pharmacist who can become an effective agent for change.

Introduction

Epilepsy is a neurological disorder caused by an abnormal electrical discharge in the brain, affecting around 50 million people worldwide [1]. About 80% of the people with epilepsy live in low- and middle-income countries. A three-quarter of the people with epilepsy may not receive the treatment they need. It is estimated up to 70.0% of the people living with epilepsy could live seizure-free if properly diagnosed, and treated with regular follow-up [1, 2]. In Libya, epilepsy has a prevalence rate of 230 cases per 100,000 people [3]. Epilepsy can be caused by genetic or acquired factors such as brain injury, stroke, brain tumours, and infections of the

brain or birth defects. Seizures are divided into two main types - generalized and partial. This is based on the international classification of epileptic seizures proposed by the Commission on Classification and Terminology of the International League against Epilepsy (ILAE) with signs and symptoms of loss of awareness or consciousness, a staring spell and disturbances of movement, sensation (including vision, hearing and taste) mood or other cognitive functions, anxiety or psychosis and depression [4]. Thus, epilepsy can be treated by anti-epileptic drugs (AEDs) including phenytoin, carbamazepine, levetiracetam and valproate. Effective of the treatment of types of epilepsy but AEDs also affect patient health as hepatotoxicity, renal stone, polycystic ovary syndrome, aplastic anaemia, rash, and decrease in hormones and vitamins of the body as vitamin B12, vitamin D are common in patients with epilepsy on AEDs. Monitoring of vitamin D levels should be considered as a part of the routine management of patients with epilepsy and lifestyle modifications and neurosurgery [5].

Pharmacist plays an essential role in educating epileptic patients about their disease and their medications. Improving the patient's knowledge and awareness may lead to improved compliance and decreased drug-drug interaction, resulting in improved clinical outcomes and decreased costs and ultimately improving their quality of life. Pharmacists involved in the review of prescriptions were able to prevent errors in dose and frequency of administration of AEDs and prevent problems related to the use of medicines. Identified therapeutic problems and proposed changes in prescriptions to a health team when the pharmacotherapy was not appropriate. Pharmacists can detect the emergence of health problems and can help prevent the progression of comorbidities. Considering the complexity of treating epilepsy and the lack of information about pharmacists' contributions to epilepsy management. Pharmacists performed pharmaceutical counselling, pharmacotherapeutic follow-up and systematic measurement as well as evaluation of findings and increased medication adherence of people with epilepsy. Pharmacists are also guided to ensure the adequacy of the pharmacotherapy through therapeutic drug monitoring. All of these successful scenarios demonstrate that including pharmacists in care staff produces effective results for the success of pharmacotherapy and the quality of life of people with epilepsy. Therefore, the present project was started by attending a training session at the epilepsy unit Ali Omer Asker Hospital by training to get acquainted with epilepsy disease and its treatment through patient observation, under the supervision of a specialist at this unit. This training allowed us to know all types of medication used to treat different types of epilepsy and prepared us to conduct the second part of the work, and that is assessing the knowledge and awareness of Libyan community pharmacists about epilepsy and its treatment with antiepileptic drugs [6]. Thus, this study aimed to identify clinical services performed by pharmacists for people with epilepsy and the services' impact on the health of these patients.

Materials and methods

The study was planned as a self-designed cross-sectional questionnaire study. The knowledge was collected through a pretested validated questionnaire. The questionnaire included 35 questions divided into three parts: the first part is demographic data (five questions), and the second part is general information about epilepsy (15 questions divided into multiple choice questions and yes or no). The third part is concentrated on data about epilepsy and antiepileptic drugs (15 questions). The questionnaire was prepared with the help of consultant training in Ali Omer Asker Hospital in the Andalus district in Tripoli for two months. Over 200 patients were altered viewed, of whom, only epileptic patients were observed. Some cases in addition to epilepsy with other complications such as diabetes mellites, hypertension, arteriosclerosis, kidney and liver diseases, heart failure, lung sensitivity, and pregnancy. Some cases of epilepsy were psychosis and depression. In addition to cases that have been completely cured of epilepsy. Throughout training, several epileptic seizure attacks were observed, some of which will be found in the results. Questionnaires were distributed online and offline for pharmacists working in different pharmacies during the beginning of 2021 and data was collected.

Ethical approval for the study was obtained from the ethics committee of the administration of the University of Tripoli, Tripoli, Libya before starting the study (2021/001). Analysis of data as descriptive statistics was conducted by using SPSS package 21.

Results

The majority of the participants were female subjects (66.0%) with a qualification degree of bachelor of pharmaceutical (79.0%) and duration of experience from one to five years (45.9%). The demographic data of the pharmacists who participated in the present study is presented in **Table 1**.

Table 1: Pharmacists demographic data of Libyan pharmacists

Total	Percentage
Gender	
Male	34.0%
Female	66.0%
Qualification degree	
Bachelor of pharmacy	79.0%
Mater of Sciences	21.0%
Years of experience	
One year	28.1%
One to five years	45.9%
Five to ten years	21.2%
More than ten years	04.8%

In **Table 2**, answers to the Q-1 that 53.3% of pharmacists believed that epilepsy is a mental disorder and a disorder caused by abnormal electrical discharges in the brain, whereas also 46.7% stated that epilepsy is a disorder caused by abnormal electrical discharges in the brain. While epilepsy is a disorder caused by abnormal electrical discharges in the brain. In response to Q-2, 80.0% of pharmacists know the situation will increase seizures, and only 20.0% answered incorrectly. 53.3% of answers to Q-3, believe that there is no correct answer to this question while the other half their answers differed on this question, from a legal point of view, epilepsy patients are prevented from driving a car but not everyone applies to this clause, so the answers to this question differ. 80.0% of the answers to Q-4 agreed that the patient does not stop taking the medicine without consulting his physician, even if the seizure has stopped. More than 66.7% of the answers to Q-5 are correct.

When answering Q-6 and 7, the majority of the responders think that epilepsy cannot be treated completely and the chance of epilepsy does not increase with age but these answers are incorrect (70.0%). 46.7% of pharmacists respond to Q-8, that violent sports are appropriate for epilepsy patients but this type of sport causes stress and pressure on patient health and, thus, the chance of a seizure increases, so epilepsy patients must have the physician's consult before doing this kind of sport. Only 26.7% of the answers to Q-9 were all people with epilepsy who lost consciousness during the seizure were answered correctly (73.3%). 60.0% of answers to Q-10 and 11 were corrected by pharmacists. In Q-10, brain damage is one of the causes of epilepsy, but not always. More than 80.0% of the pharmacists answered correctly for Q-12 about epilepsy and depression. Answers to Q-13, about pregnant epilepsy was 93.7%. A 73.3% of pharmacists responded to Q-14 which related to epilepsy as a disorder of the transmission of electrical signals within the brain. Answers to Q-15 related to careful of community pharmacists on the patient's health, as they could not prescribe any medicine without a prescription to preserve the patient's health from side effects (93.3%).

Table 2: General information about Libyan epileptic patients

Questions	Correct answer (%)	incorrect answer (%)	Corrected answer
MCQ			
Q-1: What is epilepsy? A. A mental disorder B. A disorder caused by abnormal electrical discharges in the brain C. A disorder that you can catch that causes people to shake all over D. Both A and B are correct	44.7%	55.3%	B
Q-2: In epilepsy patients, which of the following situations will increase seizures: A. Not taking his seizure medication B. Not getting enough sleep C. Stress D. All of the above	74.5%	25.5%	D
Q-3: If patient is still having seizures, he can drive a car under the following conditions: A. If he drives only when someone else is in a car B. If he “double-up” on his medication just before driving C. If he pulls over when he felt a seizure coming on D. None of the above	41.6%	58.4%	D
Q-4: If the patient stops having seizures while taking his seizure medicine, he can do the following: A. Stop his medicine because he does not need it B. Reduce his medicine without asking his doctor C. Continue taking medicine as ordered D. Stop taking his medicine every day and take double doses when he “feels a seizure coming on	81.2%	18.8%	C
Yes / No			
Q-5: Blood samples can be used to measure the concentrations of anti-epileptic drugs in the system.	75.0%	25.0%	Yes
Q-6: Epilepsy can be treated completely.	30.2%	69.8%	Yes
Q-7: The chance of epilepsy increase with age.	39.6%	60.4%	Yes
Q-8: Violent sports s such as boxing is appropriate for epilepsy to dispose of excess shipments.	63.4%	36.6%	No
Q-9: All people with epilepsy loss consciousness during a seizure.	66.7%	33.3%	No
Q-10: Epilepsy can always be caused by brain damage.	54.4%	45.6%	No
Q-11: An epileptic seizure can be described as a temporary lack of oxygen to the brain.	69.1%	30.9%	No
Q-12: Does a patient with epilepsy get depression?	80.5%	19.5%	Yes
Q-13: To prevent birth defects, women with epilepsy should stop taking their medicine by themselves, if they become pregnant.	85.8%	14.2%	No
Q-14: Are anti-epileptic drugs making a behavioural change in epilepsy children?	69.6%	30.4%	Yes
Q-15: As a pharmacist, do you have the ability to issue anti-epileptic drugs without prescription?	87.1%	12.9%	No

In **Table 3**, when answering Q-16 about (75.0%) of pharmacist's answers yes on the question if valproate is the drug of choice for juvenile myoclonic epilepsy. While 26.4% of pharmacist's answered are no. In answers to Q-17, the majority of pharmacists about 85.0% agreed that an anti-epileptic drug can be described as an abnormality in the function of nerve cells in the brain which is the correct answer about the epileptic seizure, whereas 16.6% did not agree. In response to Q-18 and 19, more than half of the responder (55.3%) suggested that vigabatrin causes aggressiveness and 66.0% answered not all seizures can affect both sides of the brain, they may even affect one part of the brain. Regarding Q-20, only 35.6% of questioned pharmacists agreed that the antiepileptic drug (valproate) can cause a decrease in vitamin D. 40.0% of pharmacists answered properly

on Q-21 regarding high level vitamin B-12 which may lead to a decrease of potassium and then may lead to heart failure. When the answer on which of these common anti-epileptic drugs cause psychosis and anorexia in Q-22, 30.0% answered carbamazepine while 45.0% answered levetiracetam which is the correct answer. About 50.0% agreed that carbamazepine act as a mood stabilizer in Q-23. In the Q-24, about 60.0% of pharmacists agreed that phenytoin causes side effects like hepatotoxicity, aplastic anaemia and intractable bleeding. 67.0% of the pharmacists believed that mothers taking phenobarbital during pregnancy may get deficient in vitamin K-dependent clotting factor at birth. In **Table 3**, when answering Q-26, the antiepileptic drug may cause impaired sexual function for males 50.0% answers carbamazepine which is correct. Q-27, about the adverse effects of valproate that bothers the women, 68.2% answered weight gain and hair loss while 22.3% answered weight gain only. 53.9% agreed valproate can cause polycystic ovary syndrome while others disagreed in Q-28. In Q-29, 52.1% answered levetiracetam is the safest drug during pregnancy while 29.5% answered valproate which could lead to the risk of disabilities such as spina bifida is 10.0%, while in Q-30, 55.2% answered carbamazepine and valproate are safe during lactation.

Table 3: Specific information about epilepsy and antiepileptic drugs

Questions	Correct	Incorrect	Answer
Yes / No			
Q-16: Drug of choice for juvenile myoclonic epilepsy is valproate.	73.4%	26.6%	Yes
Q-17: An epileptic seizure is described as abnormality in function of nerve cells in brain.	84.0%	16.0%	Yes
Q-18: Vigabatrin (Sabriel) cause aggressiveness.	55.0%	45.0%	Yes
Q-19: All seizure can affect both sides of the brain.	65.7%	34.3%	No
Q-20: Anti-epileptic drug as (valproate) can cause: A. Decrease vitamin D B. Increase vitamin B-12 C. Increase vitamin D D. B & C are correct	35.9%	64.1%	A
Q-21: Increase of vitamin B-12 in the body of epileptic patients may lead to: A. Increase in potassium and heart failure B. Decrease of potassium and heart failure C. All of the above D. None of the above	37.0%	63.0%	B
MCQ			
Q-22: Which of these common anti-epileptic drugs cause psychosis and anorexia? A. Levetiracetam B. Carbamazepine C. Valproate D. Lamotrigine	44.8%	55.2%	A
Q-23: Anti-epileptic drug act as mood stabilizer: A. Carbamazepine B. Levetiracetam C. Lamotrigine D. None of above	47.6%	52.4%	A
Q-24: Phenytoin cause side effect like: A. Hepatotoxicity B. Aplastic anaemia C. Intractable bleeding D. All of the above	60.4%	39.6%	D
Q-25: Mothers taking Phenobarbital during pregnancy may get: A. increase in Vitamin D B. Deficiency in Vitamin K dependent clotting factor at birth C. increase in Vitamin K dependent clotting factor at birth D. All of the above	68.1%	31.9%	B
Q-26: Which of these anti-epileptic drugs may cause impaired sexual function for male? A. Carbamazepine B. Oxcarbazepine C. Levetiracetam D. A & B are correct	48.6%	51.4%	A

<p>Q-27: Adverse Effect of valproate that bothers women is:</p> <p>A. Weight loss B. Weight gain C. Hair loss D. B & C are correct</p>	68.0%	32.0%	D
<p>Q-28: Which of these anti-epileptic drugs can cause polycystic ovary syndrome?</p> <p>A. Carbamazepine B. Levetiracetam C. Valproate D. Primidone</p>	53.6%	46.4%	C
<p>Q-29: Which of the anti-epileptic drug is the safest during pregnancy?</p> <p>A. Carbamazepine B. Valproate C. Levetiracetam D. All of the above</p>	51.7%	48.3%	C
<p>Q-30: Which of these anti-epileptic drugs safe on lactation?</p> <p>A. Levetiracetam+lamotrigine B. Carbamazepine+lamotrigine C. Carbamazepine+valproate D. Levetiracetam+risperidone</p>	54.9%	45.1%	C

Discussion

Libyan pharmacists are knowledgeable regarding certain aspects of epilepsy. The answers to the general question of epilepsy by community pharmacists were acceptable, because every pharmacist, with continuous study and research, must have sufficient information and how to treat them, to increase efficiency and experience that enables them to deal with the patient [7]. The participants showed poor knowledge of antiepileptic drugs with about 50.0%. It is found that few community pharmacists preferred to leave some of the questions unanswered instead of giving an incorrect answer, which may be due to the large group of them being satisfied with dispensing epilepsy drugs as they are with the prescription and thus we conclude that pharmacists must take sufficient information on some diseases and cure it [8, 9]. These findings indicate the need for a better education of pharmacists regarding epilepsy and its treatment. Valproate can cause a decrease in vitamin D and the patient must constantly conduct the necessary tests to be sure that is in the normal range [8]. Thus, the patient must be given the appropriate dose and be careful not to increase it in the body. Vitamins are necessary, but increasing it can lead to risks. In contrast, patient on carbamazepine, may see the patient accepting his condition and the treatment he is taking [9]. Unlike oxcarbazepine, which can cause an improvement in the sexual function [10]. Accordingly, it should not be given to women who want to have children or to females at young ages, except in cases where the treatment is not beneficial only with this drug [11]. About 35.0% of babies exposed to drug delays in their early development such as talking, and walking, have low intellectual abilities, poor language skills and memory problems [12]. Carbamazepine and valproate are safe during lactation, and can be excreted with breast milk, but in small quantities that do not harm the fetus therefore it is assumed reduce the dose to the maximum extent to make the patient stable and maintain the health of the fetus [13]. More is needed to investigate the actual factors behind this knowledge gap and propose interventions to improve the pharmacist's knowledge and practice to enhance the quality of health care provided for the patients. A recent study was conducted at Khartoum University, Sudan, about the knowledge of pharmacists about anti-epileptic drugs in a developing country, the results indicated that the majority have poor knowledge and only a few have good knowledge. Further research is needed to investigate the actual factors behind this knowledge gap and to propose interventions to improve pharmacist's knowledge and practice to improve the quality of healthcare provided to the patients [14].

Conclusion: Most community pharmacists have poor knowledge about anti-epileptic drugs. Further study is needed to reveal the reasons that led to this gap. Continuous training to enable the stakeholders to implement the needed interventions to improve the pharmacist's knowledge and subsequently improve the quality of health care provided to patients in Libya.

Acknowledgments: The authors would like to thank all the participants for their support of this study.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

Author contributions: AEA & HAA collected data and contributed to analysis of data with drafting the manuscript. MA, AEA & HAA contributed to the conception and compilation of data. SQ contributed to collecting and contributing to the analysis of the data. All authors have approved the final version of the manuscript and agreed to be accountable for its contents.

Conflict of interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical issues: Including plagiarism, informed consent, data fabrication or falsification and double publication or submission were completely observed by the authors.

Author declarations: The authors confirm that all relevant ethical guidelines have been followed and any necessary IRB and/or ethics committee approvals have been obtained.

References

1. Beghi E (2020) The epidemiology of epilepsy. *Neuroepidemiology*. 54 (2): 185-191. doi: 10.1159/000503831
2. Halvorsen KH, landmark CJ, Granas AG (2016) Prevalence of different combinations of antiepileptic drugs and CNS drugs in elderly home care service and nursing home patients in Norway. *Epilepsy Research and Treatment*. 2016: 1-8. doi: 10.1155/2016/5153093
3. Taher YA, Al-Gamati MA, Samud AM, El-Taher FE (2018) Knowledge and attitudes toward epilepsy among Libyan parents resident in Tripoli. *Libyan Journal of Medical Sciences*. 2 (3): 102-107. doi: 10.4103/LJMS.LJMS3718
4. Brodie MJ, Zuberi SM, Scheffer IE, Fisher RS (2018) The 2017 ILAE classification of seizure types and the epilepsies: what do people with epilepsy and their caregivers need to know? *Epileptic Disorders*. 20 (2): 77-87. doi: 10.1684/epd.2018.0957
5. Schachter SC (2020) Anticonvulsant agents: Principal considerations on effectiveness, side effects, and interactions. *Neuropsychopharmacotherapy*. 1-8. In: Riederer P, Laux G, Mulsant B, Le W, Nagatsu T (eds). Springer, Cham. Nature Switzerland AG. doi: 10.1007/978-3-319-56015-1_280-1
6. Reis TM, Campos M, Nagai MM, Pereira L (2016) Contributions of pharmacists in the treatment of epilepsy: a systematic review. *The American Journal of Pharmacy Benefits*. 8 (3): e55-e60. doi: Nil.
7. Chowdhury A, Brodie MJ (2016) Pharmacological outcomes in juvenile myoclonic epilepsy: support for sodium valproate. *Epilepsy Research*. 119: 62-66. doi: 10.1016/j.epilepsyres.2015.11.012
8. Sharma A, Sinha S, Narang A, Chouhan DK, Gupta S (2020) Waddling gait: A complication of valproate therapy and a thought beyond vitamin D deficiency. *Sultan Qaboos University Medical Journal*. 20 (1): e104. doi: 10.18295/squmj.2020.20.01.016
9. Kumar A, Maini K, Kadian R (2020) Levetiracetam. *StatPearls* [Internet]. Bookshelf ID: NBK499890PMID: 29763065.
10. Ozek SU, Ak HC, Ucler S (2020) Sexual dysfunction related with carbamazepine treatment: how much do we question?. *Neurological Sciences*. 33: 328-330. doi: 10.14744/DAJPNS.2020.00099
11. Iliescu, CATRINEL, Tarta-Arsene O, Craiu DANA (2017) Valproic acid, polycystic ovary syndrome and the adolescent with epilepsy. *Farmacia*. 65 (1): 1-4. doi: Nil.
12. Bansal R, Suri V, Chopra S, Aggarwal N, Sikka P, Saha SC, Goyal MK, Kumar P (2018) Levetiracetam use during pregnancy in women with epilepsy: Preliminary observations from a tertiary care center in Northern India. *Indian Journal of Pharmacology*. 50 (1): 39-43. doi: 10.4103/ijp.IJP_692_17
13. Uguz F, Sharma V (2016) Mood stabilizers during breastfeeding: a systematic review of the recent literature. *Bipolar Disorders*. 18 (4): 325-333. doi: 10.1111/bdi.12398
14. Ali MA, Mahgoub EAA, Nimir M, Ali KM (2020) Knowledge of pharmacists about anti-epileptic drugs in a developing country. *Current Drug Safety*. 15 (1): 32-37. doi: 10.2174/1574886314666190729114946